

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number
WO 01/96580 A2

(51) International Patent Classification⁷: **C12N 15/82**
(21) International Application Number: **PCT/EP01/06833**
(22) International Filing Date: **18 June 2001 (18.06.2001)**
(25) Filing Language: **English**
(26) Publication Language: **English**
(30) Priority Data:
00870132.8 16 June 2000 (16.06.2000) EP
60/258,415 27 December 2000 (27.12.2000) US
01870053.4 16 March 2001 (16.03.2001) EP

(71) Applicants and
(72) Inventors: **SCHMULLING, Thomas** [DE/DE];
Gertrud-Balmer-strasse 8, 72074 Tübingen (DE).
WERNER, Tomás [DE/DE]; Haldenstrasse 2, 72074
Tübingen (DE).

(74) Agents: **DE CLERCQ, Ann et al.**; De Clercq, Brants &
Partners, E. Gevaertdreef 10 a, B-9830 Sint-Martens-Latem
(BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR MODIFYING PLANT MORPHOLOGY, BIOCHEMISTRY AND PHYSIOLOGY

(57) Abstract: The present invention relates to methods for stimulating root growth and/or enhancing the formation of lateral or adventitious roots and/or altering root geotropism comprising expression of a plant cytokinin oxidase or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts. The invention also relates to novel plant cytokinin oxidase proteins, nucleic acid sequences encoding cytokinin oxidase proteins as well as to vectors, host cells, transgenic cells and plants comprising said sequences. The invention also relates to the use of said sequences for improving root-related characteristics including increasing yield and/or enhancing early vigor and/or modifying root/shoot ratio and/or improving resistance to lodging and/or increasing drought tolerance and/or promoting in vitro propagation of explants and/or modifying cell fate and/or plant development and/or plant morphology and/or plant biochemistry and/or plant physiology. The invention also relates to the use of said sequences in the above-mentioned methods. The invention also relates to methods for identifying and obtaining proteins and compounds interacting with cytokinin oxidase proteins. The invention also relates to the use of said compounds as a plant growth regulator or herbicide.



WO 01/96580 A2

METHOD FOR MODIFYING PLANT MORPHOLOGY, BIOCHEMISTRY AND PHYSIOLOGY

FIELD OF THE INVENTION

5 The present invention generally relates to a method for modifying plant morphological, biochemical and physiological properties or characteristics, such as one or more developmental processes and/or environmental adaptive processes, including but not limited to the modification of initiation or stimulation or enhancement of root growth, and/or adventitious root formation, and/or lateral root formation, and/or root geotropism,
10 and/or shoot growth, and/or apical dominance, and/or branching, and/or timing of senescence, and/or timing of flowering, and/or flower formation, and/or seed development, and/or seed yield, said method comprising expressing a cytokinin degradation control protein, in particular cytokinin oxidase, in the plant, operably under the control of a regulatable promoter sequence such as a cell-specific promoter, tissue-specific promoter, or organ-specific promoter sequence. Preferably, the characteristics
15 modified by the present invention are cytokinin-mediated and/or auxin-mediated characteristics. The present invention extends to genetic constructs which are useful for performing the inventive method and to transgenic plants produced therewith having altered morphological and/or biochemical and/or physiological properties compared to
20 their otherwise isogenic counterparts.

BACKGROUND OF THE INVENTION

Roots are an important organ of higher plants. Their main functions are anchoring of the plant in the soil and uptake of water and nutrients (N-nutrition, minerals, etc.). Thus, root
25 growth has a direct or indirect influence on growth and yield of aerial organs, particularly under conditions of nutrient limitation. Roots are also relevant for the production of secondary plant products, such as defense compounds and plant hormones.

Roots are also storage organs in a number of important staple crops. Sugar beet is the most important plant for sugar production in Europe (260 Mill t/year; 38 % of world
30 production). Manioc (cassava), yams and sweet potato (batate) are important starch producers (app. 150 Mill t/year each). Their content in starch can be twice as high as that of potato. Roots are also the relevant organ for consumption in a number of vegetables (e.g. carrots, radish), herbs (e.g. ginger, kukuma) and medicinal plants (e.g.

ginseng). In addition, some of the secondary plant products found in roots are of economic importance for the chemical and pharmaceutical industry. An example is yams, which contain basic molecules for the synthesis of steroid hormones. Another example is shikonin, which is produced by the roots of *Lithospermum erythrorhizon* in hairy root cultures. Shikonin is used for its anti-inflammatory, anti-tumour and wound-healing properties.

Moreover, improved root growth of crop plants will also enhance competitiveness with weedy plants and will improve growth in arid areas, by increasing water accessibility and uptake.

Improved root growth is also relevant for ecological purposes, such as bioremediation and prevention/arrest of soil erosion.

Root architecture is an area that has remained largely unexplored through classical breeding, because of difficulties with assessing this trait in the field. Thus, biotechnology could have significant impact on the improvement of this trait, because it does not rely on large-scale screenings in the field. Rather, biotechnological approaches require a basic understanding of the molecular components that determine a specific characteristic of the plant. Today, this knowledge is only fragmentary, and as a consequence, biotechnology was so far unable to realize a break-through in this area.

A well-established regulator of root growth is auxin. Application of indole-3-acetic acid (IAA) to growing plants stimulates lateral root development and lateral root elongation (Torrey, Am J Bot 37: 257-264, 1950; Blakely *et al.*, Bot Gaz 143: 341-352, 1982; Muday and Haworth, Plant Physiol Biochem 32: 193-203, 1994). Roots exposed to a range of concentrations of IAA initiated increasing numbers of lateral roots (Kerk *et al.*, Plant Physiol, 122: 925-932, 2000). Furthermore, when roots that had produced laterals in response to a particular concentration of exogenous auxin were subsequently exposed to a higher concentration of IAA, numerous supernumerary lateral roots spaced between existing ones were formed (Kerk *et al.*, Plant Physiol, 122: 925-932, 2000). Conversely, growth of roots on agar containing auxin-transport inhibitors, including NPA, decreases the number of lateral roots (Muday and Haworth, Plant Physiol Biochem 32: 193-203, 1994).

Arabidopsis mutants containing increased levels of endogenous IAA have been isolated (Boerjan *et al.*, Plant Cell 7: 1405-141, 1995; Celenza *et al.*, Gene Dev 9: 2131-2142, 1995; King *et al.*, Plant Cell 7: 2023-2037, 1995; Lehman *et al.*, Cell 85: 183-194, 1996).

They are now known to be alleles of a single locus located on chromosome 2. These mutant seedlings have excess adventitious and lateral roots, which is in accordance with the above-described effects of external auxin application.

5 The stimulatory effect of auxins on adventitious and lateral root formation suggests that overproduction of auxins in transgenic plants is a valid strategy for increasing root growth. Yet, it is also questionable whether this would yield a commercial product with improved characteristics. Apart from its stimulatory effect on adventitious and lateral root formation, auxin overproduction triggers other effects, such as reduction in leaf number, abnormal leaf morphology (narrow, curled leaves), aborted inflorescences, increased
10 apical dominance, adventitious root formation on the stem, most of which are undesirable from an agronomic perspective (Klee *et al.*, *Genes Devel* 1: 86-96, 1987; Kares *et al.*, *Plant Mol Biol* 15: 225-236, 1990). Therefore, the major problem with approaches that rely on increased auxin synthesis is a problem of containment, namely to confine the effects of auxin to the root. This problem of containment is not likely overcome by using
15 tissue-specific promoters: auxins are transported in the plant and their action is consequently not confined to the site of synthesis. Another issue is whether auxins will always enhance the total root biomass. For agar-grown plants, it has been noticed that increasing concentrations progressively stimulated lateral root formation but concurrently inhibited the outgrowth of these roots (Kerk *et al.*, *Plant Physiol*, 122: 925-932, 2000).
20 The above-mentioned problems related to containment of auxin effects and to maintenance of root outgrowth are solved by the embodiments in the patent claim.

SUMMARY OF THE INVENTION

25 The present invention relates to a genetic construct comprising a gene encoding a protein with cytokinin oxidase activity from *Arabidopsis thaliana*. This gene is expressed under control of a regulated promoter. This promoter may be regulated by endogenous tissue-specific or environment-specific factors or, alternatively, it may be induced by application of specific chemicals.

The present invention also relates to a cell or plant containing the genetic construct.

30 The present invention also relates to a method to modify root architecture and biomass by expression of a cytokinin oxidase gene under control of a promoter that is specific to the root or to certain tissues or cell types of the root.

DETAILED DESCRIPTION OF THE INVENTION

To by-pass above-mentioned problems associated with increasing auxin biosynthesis, it was decided to follow an alternative approach. We reasoned that down-regulation of biological antagonists of auxins could evoke similar or even superior effects on root growth as compared to increasing auxin levels. Hormone actions and interactions are extremely complex, but we hypothesized that cytokinins could function as auxin antagonists with respect to root growth. Hormone studies on plant tissue cultures have shown that the ratio of auxin versus cytokinin is more important for organogenesis than the absolute levels of each of these hormones, which indeed indicates that these hormones function as antagonists – at least in certain biological processes. Furthermore, lateral root formation is inhibited by exogenous application of cytokinins. Interestingly, also root elongation is negatively affected by cytokinin treatment, which suggests that cytokinins control both root branching and root outgrowth.

Together, current literature data indicate that increasing cytokinin levels negatively affects root growth, but the mechanisms underlying this process are not understood. The sites of cytokinin synthesis in the plant are root tips and young tissues of the shoot. Endogenous concentrations of cytokinins are in the nM range. However, as their quantification is difficult, rather large tissue amounts need to be extracted and actual local concentrations are not known. Also the subcellular compartmentation of cytokinins is not known. It is generally thought that the free base and ribosides are localized in the cytoplasm and nucleus, while glucosides are localized in the vacuole. There exist also different cytokinins with slightly different chemical structure. As a consequence, it is not known whether the effects of exogenous cytokinins should be ascribed to a raise in total cytokinin concentration or rather to the competing out of other forms of plant-borne cytokinins (which differ either in structure, cellular or subcellular location) for receptors, translocators, transporters, modifying enzymes ...

In order to test the hypothesis that cytokinin levels in the root indeed exceed the level optimal for root growth, novel genes encoding cytokinin oxidases (which are cytokinin metabolizing enzymes) were cloned from *Arabidopsis thaliana* (designated *AtCKX*) and were subsequently expressed under a strong constitutive promoter in transgenic tobacco and *Arabidopsis*. Transformants showing *AtCKX* mRNA expression and increased cytokinin oxidase activity also manifested enhanced formation and growth of roots.

Negative effects on shoot growth were also observed. The latter is in accordance with the constitutive expression of the cytokinin oxidase gene in these plants, illustrating the importance of confined expression of the cytokinin oxidase gene for general plant growth properties. Containment of cytokinin oxidase activity can be achieved by using cell-,
5 tissue- or organ-specific promoters, since cytokinin degradation is a process limited to the tissues or cells that express the CKX protein, this in contrast to approaches relying on hormone synthesis, as explained above.

The observed negative effects of cytokinin oxidase expression on shoot growth demonstrate that cytokinin oxidases are interesting targets for the design of or screening
10 for growth-promoting chemicals. Such chemicals should inhibit cytokinin oxidase activity, should preferably not be transported to the root and should be rapidly degraded in soil, so that application of these chemicals will not inhibit root growth. Cytokinins also delay leaf senescence, which means that positive effects will include both growth and maintenance of photosynthetic tissues. In addition, the observation that cytokinins delay
15 senescence, enhance greening (chlorophyll content) of leaves and reduce shoot apical dominance shows that strategies based on suppressing CKX activity (such as antisense, ribozyme, and cosuppression technology) in the aerial parts of the plant could result in delayed senescence, enhanced leaf greening and increased branching.

Similarly, the observed positive effects of cytokinin oxidase expression on root growth
20 demonstrate that cytokinin oxidases are interesting targets for the design of or screening for herbicides. Such herbicides should inhibit cytokinin oxidase activity, should preferably not be transported to the shoot, and should be soluble and relatively stable in a solvent that can be administered to the root through the soil.

These effects of cytokinin oxidase overexpression on plant development and
25 architecture were hitherto unknown and, as a consequence, the presented invention and its embodiments could not be envisaged.

The observed negative effects on shoot growth demonstrate that manipulation of cytokinin oxidases can also be used for obtaining dwarfing phenotypes. Dwarfing
30 phenotypes are particularly useful in commercial crops such as cereals and fruit trees for example.

Preferable embodiments of the invention relate to the positive effect of cytokinin oxidase expression on plant growth and architecture, and in particular on root growth and architecture. The cytokinin oxidase gene family contains at least six members in

Arabidopsis (see examples below) and the present inventors have shown that there are quantitative differences in the effects achieved with some of these genes in transgenic plants. It is anticipated that functional homologs of the described *Arabidopsis* cytokinin oxidases can be isolated from other organisms, given the evidence for the presence of cytokinin oxidase activity in many green plants (Hare and van Staden, *Physiol Plant* 91:128-136, 1994; Jones and Schreiber, *Plant Growth Reg* 23:123-134, 1997), as well as in other organisms (Armstrong, in *Cytokinins: Chemistry, Activity and Function*. Eds Mok and Mok, CRC Press, pp139-154, 1994). Therefore, the sequence of the cytokinin oxidase, functional in the invention, need not to be identical to those described herein.

This invention is particularly useful for cereal crops and monocot crops in general and cytokinin oxidase genes from for example wheat or maize may be used as well (Morris et al., 1999; Rinaldi and Comandini, 1999). It is envisaged that other genes with cytokinin oxidase activity or with any other cytokinin metabolizing activity (see Zažímalová *et al.*, *Biochemistry and Molecular Biology of Plant Hormones*, Hooykaas, Hall and Libbenga (Eds.), Elsevier Science, pp141-160, 1997) can also be used for the purpose of this invention. Similarly, genes encoding proteins that would increase endogenous cytokinin metabolizing activity can also be used for the purpose of this invention. In principle, similar phenotypes could also be obtained by interfering with genes that function downstream of cytokinin such as receptors or proteins involved in signal transduction pathways of cytokinin.

For the purpose of this invention, it should be understood that the term 'root growth' encompasses all aspects of growth of the different parts that make up the root system at different stages of its development, both in monocotyledonous and dicotyledonous plants. It is to be understood that enhanced growth of the root can result from enhanced growth of one or more of its parts including the primary root, lateral roots, adventitious roots, etc. all of which fall within the scope of this invention.

According to a first embodiment, the present invention relates to a method for stimulating root growth and/or enhancing the formation of lateral and/or adventitious roots and/or altering root geotropism comprising expression of a plant cytokinin oxidase or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

In the context of the present invention it should be understood that the term "expression" and/or 'overexpression' are used interchangeably and both relate to an "enhanced

and/or ectopic expression" of a plant cytokinin oxidase or any other protein that reduces the level of active cytokinins in plants. It should be clear that herewith an enhanced expression of the plant cytokinin oxidase as well as "de novo" expression of plant cytokinin oxidases or of said other proteins is meant. Alternatively, said other protein
5 enhances the cytokinin metabolizing activity of a plant cytokinin oxidase.

It further should be understood that in the context of the present invention the expression "lateral and/or adventitious roots" can mean "lateral and adventitious roots" but also "lateral or adventitious roots". The enhancement can exist in the formation of lateral roots or in the formation of adventitious roots as well as in the formation of both types of non-
10 primary roots, but not necessarily.

According to a further embodiment, the present invention relates to a method for stimulating root growth and/or enhancing the formation of lateral or adventitious roots and/or altering root geotropism and/or increasing yield and/or enhancing early vigor and/or modifying root/shoot ratio and/or improving resistance to lodging and/or
15 increasing drought tolerance and/or promoting in vitro propagation of explants, comprising expression of a plant cytokinin oxidase or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

According to a preferred embodiment, the present invention relates to a method for stimulating root growth resulting in an increase of root mass by overexpression of a
20 cytokinin oxidase, preferably a cytokinin oxidase according to the invention, or another protein that reduces the level of active cytokinins in plants or plant parts, preferably in roots.

Higher root biomass production due to overexpression of growth promoting sequences has a direct effect on the yield and an indirect effect of production of compounds
25 produced by root cells or transgenic root cells or cell cultures of said transgenic root cells. One example of an interesting compound produced in root cultures is shikonin, the yield of which can be advantageously enhanced by said methods.

According to a more specific embodiment, the present invention relates to a method for stimulating root growth or for enhancing the formation of lateral and adventitious roots or
30 for altering root geotropism comprising expression of a nucleic acid encoding a plant cytokinin oxidase selected from the group consisting of:

- (a) nucleic acids comprising a DNA sequence as given in any of SEQ ID NOs 27, 1, 3, 5, 7, 9, 11, 25, 26, 28 to 31, 33 or 34, or the complement thereof,

(b) nucleic acids comprising the RNA sequences corresponding to any of SEQ ID NOs 27, 1, 3, 5, 7, 9, 11, 25, 26, 28 to 31, 33 or 34, or the complement thereof,

(c) nucleic acids specifically hybridizing to any of SEQ ID NOs 27, 1, 3, 5, 7, 9, 11, 25, 26, 28 to 31, 33 or 34, or to the complement thereof,

(d) nucleic acids encoding a protein comprising the amino acid sequence as given in any of SEQ ID NOs 2, 4, 6, 8, 10, 12, 32 or 35, or the complement thereof,

(e) nucleic acids as defined in any of (a) to (d) characterized in that said nucleic acid is DNA, genomic DNA, cDNA, synthetic DNA or RNA wherein T is replaced by U,

(f) nucleic acids which are degenerated to a nucleic acid as given in any of SEQ ID NOs 27, 1, 3, 5, 7, 9, 11, 25, 26, 28 to 31, 33 or 34, or which are degenerated to a nucleic acid as defined in any of (a) to (e) as a result of the genetic code,

(g) nucleic acids which are diverging from a nucleic acid encoding a protein as given in any of SEQ ID NOs 2, 4, 6, 8, 10, 12 or 35 or which are diverging from a nucleic acid as defined in any of (a) to (e), due to the differences in codon usage between the organisms,

(h) nucleic acids encoding a protein as given in SEQ ID NOs 2, 4, 6, 8, 10, 12 or 35 or nucleic acids as defined in (a) to (e) which are diverging due to the differences between alleles,

(i) nucleic acids encoding a protein as given in any of SEQ ID NOs 2, 4, 6, 8, 10, 12 or 35,

(j) functional fragments of nucleic acids as defined in any of (a) to (i) having the biological activity of a cytokinin oxidase, and

(k) nucleic acids encoding a plant cytokinin oxidase,

or comprising expression, preferably in roots, of a nucleic acid encoding a protein that reduces the level of active cytokinins in plants or plant parts.

In the present invention, nucleic acids encoding novel *Arabidopsis thaliana* cytokinin oxidases have been isolated and for the first time, the present inventors surprisingly could show that the expression of cytokinin oxidases in transgenic plants or in transgenic plant parts resulted in the above-mentioned root-related features. Preferably, the expression of the cytokinin oxidase(s) should take place in roots, preferably under the control of a root-specific promoter. One example of such a root-specific promoter is provided in SEQ ID NO 36.

It should be clear that, although the invention is supported in the examples section by several new AtCKX genes and proteins, the inventive concept also relates to the use of other cytokinin oxidases isolated from and expressed in other plants, preferably in the roots of said other plants to obtain similar effects in plants as described in the examples section.

Therefore, the present invention more generally relates to the use of a nucleic acid encoding a plant cytokinin oxidase or encoding a protein that reduces the level of active cytokinins in plants or plant parts for stimulating root growth or for enhancing the formation of lateral or adventitious roots or for altering root geotropism. Preferred cytokinin oxidases to be used are encoded by the nucleic acids encoding the cytokinin oxidases as defined above and are encoded by the novel nucleic acids of the invention as defined hereunder.

The invention relates to an isolated nucleic acid encoding a novel plant protein having cytokinin oxidase activity selected from the group consisting of:

- (a) a nucleic acid comprising a DNA sequence as given in any of SEQ ID NOs 29, 3, 5, 9, 26, 27, 31, 33 or 34, or the complement thereof,
- (b) a nucleic acid comprising the RNA sequences corresponding to any of SEQ ID NOs 29, 3, 5, 9, 26, 27, 31, 33 or 34, or the complement thereof,
- (c) a nucleic acid specifically hybridizing to a nucleic acid as given in any of SEQ ID NOs 29, 3, 5, 9, 26, 27, 31, 33 or 34, or the complement thereof,
- (d) a nucleic acid encoding a protein with an amino acid sequence comprising the polypeptide as given in SEQ ID NO 32 and which is at least 70% similar, preferably at least 75%, 80% or 85%, more preferably at least 90% or 95%, most preferably at least 99% similar to the amino acid sequence as given in SEQ ID NO 4,
- (e) a nucleic acid encoding a protein with an amino acid sequence which is at least 35% similar, preferably 37%, 40%, 45%, 47% or 50%, similar, more preferably 55%, 60%, 65%, 70%, 75% or 80% similar, most preferably 85%, 90% or 95% similar to the amino acid sequence as given in SEQ ID NO 6,
- (f) a nucleic acid encoding a protein with an amino acid sequence which is at least 35% similar, preferably 37%, 40%, 45%, 47% or 50%, similar, more preferably 55%, 60%, 65%, 70%, 75% or 80% similar, most preferably 85%, 90% or 95% similar to the amino acid sequence as given in SEQ ID NO 10 or 35,

(g) a nucleic acid encoding a protein comprising the amino acid sequence as given in any of SEQ ID NOs 4, 6, 10, 32 or 35,

(h) a nucleic acid which is degenerated to a nucleic acid as given in any of SEQ ID NOs 29, 3, 5, 9, 26, 27, 33 or 34 or which is degenerated to a nucleic acid as defined in any of (a) to (g) as a result of the genetic code,

(i) a nucleic acid which is diverging from a nucleic acid encoding a protein as given in any of SEQ ID NOs 4, 6, 10 or 35 or which is diverging from a nucleic acid as defined in any of (a) to (g) due to the differences in codon usage between the organisms,

(j) a nucleic acid encoding a protein as given in SEQ ID NOs 4, 6, 10 or 35, or a nucleic acid as defined in (a) to (g) which is diverging due to the differences between alleles,

(k) a nucleic acid encoding an immunologically active fragment of a cytokinin oxidase encoded by a nucleic acid as given in any of SEQ ID NOs 29, 3, 5, 9, 26, 27, 31, 33 or 34, or an immunologically active fragment of a nucleic acid as defined in any of (a) to (j),

(l) a nucleic acid encoding a functional fragment of a cytokinin oxidase encoded by a nucleic acid as given in any of SEQ ID NOs 29, 3, 5, 9, 26, 27, 31, 33 or 34, or a functional fragment of a nucleic acid as defined in any of (a) to (j), wherein said fragment has the biological activity of a cytokinin oxidase, and

(m) a nucleic acid encoding a protein as defined in SEQ ID NO 4, 6, 10 or 35,

provided that said nucleic acid is not the nucleic acid as deposited under any of the following Genbank accession numbers: AC005917, AB024035, and AC023754

The invention also relates to an isolated nucleic acid of the invention which is DNA, cDNA, genomic DNA or synthetic DNA, or RNA wherein T is replaced by U.

The invention also relates to a nucleic acid molecule of at least 15 nucleotides in length hybridizing specifically with or specifically amplifying a nucleic acid of the invention.

According to another embodiment, the invention also relates to a vector comprising a nucleic acid of the invention. In a preferred embodiment, said vector is an expression vector wherein the nucleic acid is operably linked to one or more control sequences allowing the expression of said sequence in prokaryotic and/or eukaryotic host cells.

It should be understood that for expression of the cytokinin oxidase genes of the invention in monocots, a nucleic acid sequence corresponding to the cDNA sequence

should be used to avoid mis-splicing of introns in monocots. Preferred cDNA sequences to be expressed in monocots have a nucleic acid sequence as represented in any of SEQ ID NOs 25 to 30 and 34.

The invention also relates to a host cell containing any of the nucleic acid molecules or
5 vectors of the invention. Said host cell is chosen from the group comprising bacterial, insect, fungal, plant or animal cells.

Another embodiment of the invention relates to an isolated polypeptide encodable by a nucleic acid of the invention, or a homologue or a derivative thereof, or an immunologically active or a functional fragment thereof. Preferred polypeptides of the
10 invention comprise the amino acid sequences as represented in any of SEQ ID NOs 2, 4, 6, 8, 10, 12, 32 and 35, or a homologue or a derivative thereof, or an immunologically active and/or functional fragment thereof. In an even more preferred embodiment, the invention relates to a polypeptide which has an amino acid sequence as given in SEQ ID
15 NO 2, 4, 6, 8, 10, 12 or 35, or a homologue or a derivative thereof, or an immunologically active and/or functional fragment thereof. Preferred functional fragments thereof are those fragments which are devoid of their signal peptide.

According to yet another embodiment, the invention relates to a method for producing a polypeptide of the invention comprising culturing a host cell of the invention under conditions allowing the expression of the polypeptide and recovering the produced
20 polypeptide from the culture.

The invention also relates to an antibody specifically recognizing a polypeptide of the invention or a specific epitope thereof.

The invention further relates to a method for the production of transgenic plants, plant cells or plant tissues comprising the introduction of a nucleic acid molecule of the
25 invention in an expressible format or a vector of the invention in said plant, plant cell or plant tissue.

The invention also relates to a method for the production of altered plants, plant cells or plant tissues comprising the introduction of a polypeptide of the invention directly into a cell, a tissue or an organ of said plant.

30 According to another embodiment, the invention relates to a method for effecting the expression of a polypeptide of the invention comprising the introduction of a nucleic acid molecule of the invention operably linked to one or more control sequences or a vector

of the invention stably into the genome of a plant cell. The invention further relates to the method as described above further comprising regenerating a plant from said plant cell.

The invention also relates to a transgenic plant cell comprising a nucleic acid sequence of the invention which is operably linked to regulatory elements allowing transcription and/or expression of said nucleic acid in plant cells or obtainable by a method as explained above.

According to another preferred embodiment, the invention relates to a transgenic plant cell as described here above wherein the nucleic acid of the invention is stably integrated into the genome of said plant cell.

The invention further relates to a transgenic plant or plant tissue comprising plant cells as herein described and also to a harvestable part of said transgenic plant, preferably selected from the group consisting of seeds, leaves, fruits, stem cultures, roots, tubers, rhizomes and bulbs. The invention also relates to the progeny derived from any of said transgenic plants or plant parts.

According to another embodiment, the invention relates to a method for stimulating root growth comprising expression of a nucleic acid of the invention or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

A plant cell or tissue culture is an artificially produced culture of plants cells or plant tissues that is grown in a special medium, either liquid or solid, which provides these plant cells or tissues with all requirements necessary for growth and/or production of certain compounds. Plant cell and/or tissue cultures can be used for the rapid propagation of plants and for the production of transgenic plant to name a few examples.

Root formation can be difficult for some explants or under some conditions in said cultures and expression of a cytokinin oxidase gene in said cultured plant cells or

tissue(s) can be used to enhance root formation. Plant cell and/or tissue culture can also be used for the industrial production of valuable compounds. Possible production compounds are pharmaceuticals, pesticides, pigments, cosmetics, perfumes, food additives, etc. An example of such a product is shikonin, which is produced by the roots of the plant *Lithospermum erythrorhizon*. An example of a plant tissue culture is a hairy

root culture, which is an artificially produced mass of hairy roots. Roots of *L. erythrorhizon* are difficult to collect in large numbers and by preparing hairy root cultures, the end product shikonin could be industrially prepared at a faster rate than would normally occur. As disclosed herein, expression of cytokinin oxidases enhances root

growth and development and can therefore be used advantageously in said plant cell and tissue culture procedures. Therefore, according to another embodiment of this invention, a method is provided for stimulating root growth and development comprising expression of a nucleic acid encoding a plant cytokinin oxidase, preferably a cytokinin oxidase of the invention, in a transgenic plant cell or tissue culture comprising said transgenic plant cells.

The invention further relates to a method for enhancing the formation of lateral or adventitious roots comprising expression of a nucleic acid of the invention or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

The invention also relates to method for altering root geotropism comprising altering the expression of a nucleic acid of the invention or comprising expression of another protein that that reduces the level of active cytokinins in plants or plant parts.

The invention also relates to methods for enhancing early vigor and/or for modifying root/shoot ratio and/or for improving resistance to lodging and/or for increasing drought tolerance and/or for promoting *in vitro* propagation of explants comprising expression of a nucleic acid of the invention comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

The invention further relates to methods for increasing the root size or the size of the root meristem comprising expression of a nucleic acid of the invention or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts, preferably in roots.

According to yet another embodiment, the invention relates to a method for increasing the size of the shoot meristem comprising downregulation of expression of a nucleic acid of the invention, preferably in shoots.

According to a preferred embodiment the invention relates to a method for delaying leaf senescence comprising downregulation of expression of any of the cytokinin oxidases of the invention in leaves, preferably in senescing leaves. Also the invention relates to a method for altering leaf senescence comprising expression of one of the cytokinin oxidases in senescing leaves.

The invention also relates to methods for increasing leaf thickness comprising expression of a nucleic acid of the invention or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts, preferably in leaves.

The invention also relates to a method for reducing the vessel size comprising expression of a nucleic acid of the invention or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts, preferably in vessels.

The invention further relates to a method for increasing the vessel size comprising
5 downregulation of expression of a nucleic acid of the invention in plants or plant parts.

According to another embodiment, the invention relates to a method for improving standability of seedlings comprising expression of a nucleic acid of the invention or comprising expression of another protein that reduces the level of active cytokinins in seedlings.

10 Furthermore, the invention relates to any of the above described methods, said method leading to an increase in yield.

The invention further relates to any of the methods of the invention wherein said expression of said nucleic acid occurs under the control of a strong constitutive promoter. In a preferred embodiment the invention relates to any of the methods of the
15 invention wherein said expression of said nucleic acid occurs under the control of a promoter that is preferentially expressed in roots. In Table 5 a non-exhaustive list of root specific promoters is included. A preferred promoter to be used in the methods of the invention is the root *clavata* homolog promoter, having a sequence as given in SEQ ID NO 36.

20 According to yet another embodiment, the invention relates to a method for modifying cell fate and/or modifying plant development and/or modifying plant morphology and/or modifying plant biochemistry and/or modifying plant physiology and/or modifying the cell cycle progression rate comprising the modification of expression in particular cells, tissues or organs of a plant, of a nucleic acid of the invention.

25 The invention also relates to a method for obtaining enhanced growth, and/or increased yield and/or altered senescence of a plant cell, tissue and/or organ and/or increased frequency of formation of lateral organs in a plant, comprising the ectopic expression of a nucleic acid of the invention.

The invention also relates to a method for promoting and extending cell division activity
30 in cells in adverse growth conditions and/or in stress, comprising the ectopic expression of a nucleic acid sequence of the invention.

According to yet another embodiment, the invention relates to a method for identifying and obtaining proteins interacting with a polypeptide of the invention comprising a screening assay wherein a polypeptide of the invention is used.

5 In a more preferred embodiment, the invention relates to a method for identifying and obtaining proteins interacting with a polypeptide of the invention comprising a two-hybrid screening assay wherein a polypeptide of the invention as a bait and a cDNA library as prey are used.

10 The invention further relates to a method for modulating the interaction between a polypeptide of the invention and interacting protein partners obtainable by a method as described above.

In a further embodiment, the invention relates to a method for identifying and obtaining compounds interacting with a polypeptide of the invention comprising the steps of:

- a) providing a two-hybrid system wherein a polypeptide of the invention and an interacting protein partner obtainable by a method as described above,
- 15 b) interacting said compound with the complex formed by the expressed polypeptides as defined in a), and,
- c) performing (real-time) measurement of interaction of said compound with said polypeptide or the complex formed by the expressed polypeptides as defined in a).

20 The invention further relates to a method for identifying compounds or mixtures of compounds which specifically bind to a polypeptide of the invention, comprising:

- a) combining a polypeptide of the invention with said compound or mixtures of compounds under conditions suitable to allow complex formation, and,
- b) detecting complex formation, wherein the presence of a complex identifies a
- 25 compound or mixture which specifically binds said polypeptide.

The invention also relates to a method as described above wherein said compound or mixture inhibits the activity of said polypeptide of the invention and can be used for the rational design of chemicals.

30 According to another embodiment, the invention relates to the use of a compound or mixture identified by means of a method as described above as a plant growth regulator or herbicide.

The invention also relates to a method for production of a plant growth regulator or herbicide composition comprising the steps of the compound screening methods

described above and formulating the compounds obtained from said steps in a suitable form for the application in agriculture or plant cell or tissue culture.

The invention also relates to a method for increasing branching comprising expression of a nucleic acid of the invention in plants or plant parts, preferably in stems or axillary buds.

The invention also relates to a method for improving lodging resistance comprising expression of a nucleic acid of the invention in plants or plant parts, preferably in stems or axillary buds.

The invention also relates to a method for the design of or screening for growth-promoting chemicals or herbicides comprising the use of a nucleic acid of the invention or a vector of the invention.

According to another embodiment, the invention relates to the use of a nucleic acid molecule of the invention, a vector of the invention or a polypeptide of the invention for increasing yield.

The invention also relates to the use of a nucleic acid molecule of the invention, a vector of the invention or a polypeptide of the invention for stimulating root growth.

The invention also relates to the use of a nucleic acid molecule of the invention, a vector of the invention or a polypeptide of the invention for enhancing the formation of lateral or adventitious roots.

The invention also relates to the use of a nucleic acid molecule of the invention, a vector of the invention or a polypeptide of the invention for altering root geotropism.

The invention further relates to the use of a nucleic acid molecule of the invention, a vector of the invention or a polypeptide of the invention for enhancing early vigor and/or for modifying root/shoot ratio and/or for improving resistance to lodging and/or for increasing drought tolerance and/or for promoting *in vitro* propagation of explants.

The invention also relates to the use of a nucleic acid molecule of the invention, a recombinant vector of the invention or a polypeptide of the invention for modifying plant development and/or for modifying plant morphology and/or for modifying plant biochemistry and/or for modifying plant physiology.

According to yet another embodiment, the invention relates to a diagnostic composition comprising at least a nucleic acid molecule of the invention, a vector of the invention, a polypeptide of the invention or an antibody of the invention.

Another embodiment of the current invention relates to the use of a transgenic rootstock that has an enhanced root growth and development due to expression of a cytokinin oxidase in grafting procedures with a scion to produce a plant or tree with improved agricultural or horticultural characteristics. The scion may be transgenic or non-transgenic. Specific characteristics envisaged by this embodiment are those conferred by root systems and include improved anchoring of the plant/tree in the soil and/or improved uptake of water resulting for example in improved drought tolerance, and/or improved nutrient uptake from the soil and/or improved transport of organic substances throughout the plant and/or enhanced secretion of substances into the soil such as for example phyto siderophores, and/or improved respiration and/or improved disease resistance and/or enhanced yield. An advantage of using *AtCKX* transformed rootstocks for grafting, in addition to their enhanced root system, is the delayed senescence of leaves on the graft, as disclosed herein (see Figure 12 A). Preferred plants or trees for this particular embodiment include plants or trees that do not grow well on their own roots and are grafted in cultivated settings such as commercially profitable varieties of grapevines, citrus, apricot, almond, plum, peach, apple, pear, cherry, walnut, fig, hazel and loquat.

As mentioned *supra*, auxins and cytokinins act as antagonists in certain biological processes. For example, the cytokinin/auxin ratio regulates the production of roots and shoots with a high concentration of auxin resulting in organized roots and a high concentration of cytokinins resulting in shoot production. As disclosed in this invention, expression of cytokinin oxidases in tobacco and Arabidopsis results in enhanced root development consistent with enhanced auxin effects. Auxins are also involved in the development of fruit. Treatment of female flower parts with auxin results in the development of parthenocarpic fruit in some plant species. Parthenocarpic fruit development has been genetically engineered in several horticultural crop plants through increased biosynthesis of auxins in the female reproductive organs (WO0105985).

Therefore, according to another embodiment, this invention relates to a method for inducing the parthenocarpic trait in plants, said method consisting of downregulating the expression of one or more cytokinin oxidases or of another protein that reduces the level of active cytokinins in plants or plant parts, preferably in the female reproductive organs such as the placenta, ovules and tissues derived therefrom. The DefH9 promoter region

from *Antirrhinum majus* or one of its homologues, which confer high expression specificity in placenta and ovules, can be used for this purpose.

5 DEFINITIONS AND ELABORATIONS TO THE EMBODIMENTS

Those skilled in the art will be aware that the invention described herein is subject to variations and modifications other than those specifically described. It is to be understood that the invention described herein includes all such variations and
 10 modifications. The invention also includes all such steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any or more of said steps or features.

The present invention is applicable to any plant, in particular a monocotyledonous plants and dicotyledonous plants including a fodder or forage legume, ornamental plant, food
 15 crop, tree, or shrub selected from the list comprising *Acacia spp.*, *Acer spp.*, *Actinidia spp.*, *Aesculus spp.*, *Agathis australis*, *Albizia amara*, *Alsophila tricolor*, *Andropogon spp.*, *Arachis spp.*, *Areca catechu*, *Astelia fragrans*, *Astragalus cicer*, *Baikiaea plurijuga*, *Betula spp.*, *Brassica spp.*, *Bruguiera gymnorrhiza*, *Burkea africana*, *Butea frondosa*, *Cadaba farinosa*, *Calliandra spp.*, *Camellia sinensis*, *Canna indica*, *Capsicum spp.*, *Cassia spp.*,
 20 *Centroema pubescens*, *Chaenomeles spp.*, *Cinnamomum cassia*, *Coffea arabica*, *Colophospermum mopane*, *Coronillia varia*, *Cotoneaster serotina*, *Crataegus spp.*, *Cucumis spp.*, *Cupressus spp.*, *Cyathea dealbata*, *Cydonia oblonga*, *Cryptomeria japonica*, *Cymbopogon spp.*, *Cynthea dealbata*, *Cydonia oblonga*, *Dalbergia monetaria*, *Davallia divaricata*, *Desmodium spp.*, *Dicksonia squarosa*, *Diheteropogon amplexans*,
 25 *Dioclea spp.*, *Dolichos spp.*, *Dorycnium rectum*, *Echinochloa pyramidalis*, *Ehrhartia spp.*, *Eleusine coracana*, *Eragrestis spp.*, *Erythrina spp.*, *Eucalyptus spp.*, *Euclea schimperi*, *Eulalia villosa*, *Fagopyrum spp.*, *Feijoa sellowiana*, *Fragaria spp.*, *Flemingia spp.*, *Freycinetia banksii*, *Geranium thunbergii*, *Ginkgo biloba*, *Glycine javanica*, *Gliricidia spp.*, *Gossypium hirsutum*, *Grevillea spp.*, *Guibourtia coleosperma*, *Hedysarum spp.*,
 30 *Hemarthia altissima*, *Heteropogon contortus*, *Hordeum vulgare*, *Hyparrhenia rufa*, *Hypericum erectum*, *Hyperthelia dissoluta*, *Indigo incamata*, *Iris spp.*, *Leptarrhena pyrolifolia*, *Lespedeza spp.*, *Lettuca spp.*, *Leucaena leucocephala*, *Loudetia simplex*, *Lotonus bainesii*, *Lotus spp.*, *Macrotyloma axillare*, *Malus spp.*, *Manihot esculenta*, *Medicago sativa*, *Metasequoia glyptostroboides*, *Musa sapientum*, *Nicotianum spp.*,

Onobrychis spp., *Ornithopus* spp., *Oryza* spp., *Peltophorum africanum*, *Pennisetum* spp., *Persea gratissima*, *Petunia* spp., *Phaseolus* spp., *Phoenix canariensis*, *Phormium cookianum*, *Photinia* spp., *Picea glauca*, *Pinus* spp., *Pisum sativum*, *Podocarpus totara*, *Pogonarthria fleckii*, *Pogonarthria squarrosa*, *Populus* spp., *Prosopis cineraria*,
 5 *Pseudotsuga menziesii*, *Pterolobium stellatum*, *Pyrus communis*, *Quercus* spp., *Rhaphiolepis umbellata*, *Rhopalostylis sapida*, *Rhus natalensis*, *Ribes grossularia*, *Ribes* spp., *Robinia pseudoacacia*, *Rosa* spp., *Rubus* spp., *Salix* spp., *Schyzachyrium sanguineum*, *Sciadopitys verticillata*, *Sequoia sempervirens*, *Sequoiadendron giganteum*, *Sorghum bicolor*, *Spinacia* spp., *Sporobolus fimbriatus*, *Stiburus*
 10 *alopecuroides*, *Stylosanthos humilis*, *Tadehagi* spp., *Taxodium distichum*, *Themeda triandra*, *Trifolium* spp., *Triticum* spp., *Tsuga heterophylla*, *Vaccinium* spp., *Vicia* spp., *Vitis vinifera*, *Watsonia pyramidata*, *Zantedeschia aethiopica*, *Zea mays*, amaranth, artichoke, asparagus, broccoli, brussel sprout, cabbage, canola, carrot, cauliflower, celery, collard greens, flax, kale, lentil, oilseed rape, okra, onion, potato, rice, soybean,
 15 straw, sugarbeet, sugar cane, sunflower, tomato, squash, and tea, amongst others, or the seeds of any plant specifically named above or a tissue, cell or organ culture of any of the above species.

Throughout this specification, unless the context requires otherwise the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the
 20 inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

As used herein, the term "derived from" shall be taken to indicate that a particular integer or group of integers has originated from the species specified, but has not necessarily been obtained directly from the specified source.

25 The terms "protein(s)", "peptide(s)" or "oligopeptide(s)", when used herein refer to amino acids in a polymeric form of any length. Said terms also include known amino acid modifications such as disulphide bond formation, cysteinylolation, oxidation, glutathionylation, methylation, acetylation, farnesylation, biotinylation, stearylolation, formylation, lipoic acid addition, phosphorylation, sulphation, ubiquitination,
 30 myristoylation, palmitoylation, geranylgeranylation, cyclization (e.g. pyroglutamic acid formation), oxidation, deamidation, dehydration, glycosylation (e.g. pentoses, hexosamines, N-acetylhexosamines, deoxyhexoses, hexoses, sialic acid etc.) and

acylation as well as non-naturally occurring amino acid residues, L-amino acid residues and D-amino acid residues.

"Homologues" of a protein of the invention are those peptides, oligopeptides, polypeptides, proteins and enzymes which contain amino acid substitutions, deletions and/or additions relative to the said protein with respect to which they are a homologue, without altering one or more of its functional properties, in particular without reducing the activity of the resulting. For example, a homologue of said protein will consist of a bioactive amino acid sequence variant of said protein. To produce such homologues, amino acids present in the said protein can be replaced by other amino acids having similar properties, for example hydrophobicity, hydrophilicity, hydrophobic moment, antigenicity, propensity to form or break α -helical structures or β -sheet structures, and so on. An overview of physical and chemical properties of amino acids is given in Table 1.

Substitutional variants of a protein of the invention are those in which at least one residue in said protein amino acid sequence has been removed and a different residue inserted in its place. Amino acid substitutions are typically of single residues, but may be clustered depending upon functional constraints placed upon the polypeptide; insertions will usually be of the order of about 1-10 amino acid residues. and deletions will range from about 1-20 residues. Preferably, amino acid substitutions will comprise conservative amino acid substitutions, such as those described *supra*.

Table 1. Properties of naturally occurring amino acids.

Charge properties / hydrophobicity	Side group	Amino Acid
nonpolar hydrophobic	Aliphatic aliphatic, S-containing aromatic imino	ala, ile, leu, val met phe, trp pro
polar uncharged	Aliphatic amide aromatic hydroxyl sulfhydryl	gly asn, gln tyr ser, thr cys

positively charged	Basic	arg, his, lys
negatively charged	Acidic	asp, glu

Insertional amino acid sequence variants of a protein of the invention are those in which one or more amino acid residues are introduced into a predetermined site in said protein. Insertions can comprise amino-terminal and/or carboxy-terminal fusions as well as intra-sequence insertions of single or multiple amino acids. Generally, insertions within the amino acid sequence will be smaller than amino or carboxyl terminal fusions, of the order of about 1 to 10 residues. Examples of amino- or carboxy-terminal fusion proteins or peptides include the binding domain or activation domain of a transcriptional activator as used in a two-hybrid system, phage coat proteins, (histidine)₆-tag, glutathione S-transferase, protein A, maltose-binding protein, dihydrofolate reductase, Tag•100 epitope (EETARFQPGYRS), c-myc epitope (EQKLISEEDL), FLAG[®]-epitope (DYKDDDK), lacZ, CMP (calmodulin-binding peptide), HA epitope (YPYDVPDYA), protein C epitope (EDQVDPRLIDGK) and VSV epitope (YTDIEMNRLGK).

Deletional variants of a protein of the invention are characterised by the removal of one or more amino acids from the amino acid sequence of said protein.

Amino acid variants of a protein of the invention may readily be made using peptide synthetic techniques well known in the art, such as solid phase peptide synthesis and the like, or by recombinant DNA manipulations. The manipulation of DNA sequences to produce variant proteins which manifest as substitutional, insertional or deletional variants are well known in the art. For example, techniques for making substitution mutations at predetermined sites in DNA having known sequence are well known to those skilled in the art, such as by M13 mutagenesis, T7-Gen in vitro mutagenesis kit (USB, Cleveland, OH), QuickChange Site Directed mutagenesis kit (Stratagene, San Diego, CA), PCR-mediated site-directed mutagenesis or other site-directed mutagenesis protocols.

In the current invention "identity" and/or "similarity" percentages between DNA sequences and/or proteins are calculated using computer programs known in the art such as the DNASTAR/MegAlign programs in combination with the Clustal method.

"Derivatives" of a protein of the invention are those peptides, oligopeptides, polypeptides, proteins and enzymes which comprise at least about five contiguous

amino acid residues of said polypeptide but which retain the biological activity of said protein. A "derivative" may further comprise additional naturally-occurring, altered glycosylated, acylated or non-naturally occurring amino acid residues compared to the amino acid sequence of a naturally-occurring form of said polypeptide. Alternatively or in addition, a derivative may comprise one or more non-amino acid substituents compared to the amino acid sequence of a naturally-occurring form of said polypeptide, for example a reporter molecule or other ligand, covalently or non-covalently bound to the amino acid sequence such as, for example, a reporter molecule which is bound thereto to facilitate its detection.

With "immunologically active" is meant that a molecule or specific fragments thereof such as specific epitopes or haptens are recognized by, i.e. bind to antibodies. Specific epitopes may be determined using, for example, peptide scanning techniques as described in Geysen *et al.* (1996) (Geysen, H.M., Rodda, S.J. and Mason, T.J. (1986). A priori delineation of a peptide which mimics a discontinuous antigenic determinant. *Mol. Immunol.* 23, 709-715.).

The term "fragment of a sequence" or "part of a sequence" means a truncated sequence of the original sequence referred to. The truncated sequence (nucleic acid or protein sequence) can vary widely in length; the minimum size being a sequence of sufficient size to provide a sequence with at least a comparable function and/or activity or the original sequence referred to (e. g. "functional fragment"), while the maximum size is not critical. In some applications, the maximum size usually is not substantially greater than that required to provide the desired activity and/or function(s) of the original sequence. Typically, the truncated amino acid sequence will range from about 5 to about 60 amino acids in length. More typically, however, the sequence will be a maximum of about 50 amino acids in length, preferably a maximum of about 60 amino acids. It is usually desirable to select sequences of at least about 10, 12 or 15 amino acids, up to a maximum of about 20 or 25 amino acids.

Functional fragments can also include those comprising an epitope which is specific for the proteins according to the invention. Preferred functional fragments have a length of at least, for example, 5, 10, 25, 100, 150 or 200 amino acids.

It should thus be understood that functional fragments can also be immunologically active fragments or not.

In the context of the current invention are embodied homologues, derivatives and/or immunologically active and/or functional fragments of the cytokinin oxidases as defined supra. Particularly preferred homologues, derivatives and/or immunologically active and/or functional fragments of the cytokinin oxidase proteins which are contemplated for use in the current invention are derived from plants, more specifically from *Arabidopsis thaliana*, even more specifically said cytokinin oxidases are the *Arabidopsis thaliana* (At)CKX, or are capable of being expressed therein. The present invention clearly contemplates the use of functional homologues or derivatives and/or immunologically active fragments of the AtCKX proteins and is not to be limited in application to the use of a nucleotide sequence encoding one of said AtCKX proteins.

Any of said proteins, polypeptides, peptides and fragments thereof can be produced in a biological system, e.g. a cell culture. Alternatively any of said proteins, polypeptides, peptides and fragments thereof can be chemically manufactured e.g. by solid phase peptide synthesis. Said proteins or fragments thereof can be part of a fusion protein as is the case in e.g. a two-hybrid assay which enables e.g. the identification of proteins interacting with a cytokinin oxidase according to the invention.

The proteins or fragments thereof are furthermore useful e.g. to modulate the interaction between a cytokinin oxidase according to the invention and interacting protein partners obtained by a method of the invention. Chemically synthesized peptides are particularly useful e.g. as a source of antigens for the production of antisera and/or antibodies.

"Antibodies" include monoclonal, polyclonal, synthetic or heavy chain camel antibodies as well as fragments of antibodies such as Fab, Fv or scFv fragments. Monoclonal antibodies can be prepared by the techniques as described in e.g. Liddle and Cryer (1991) which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized animals. Furthermore, antibodies or fragments thereof to a molecule or fragments thereof can be obtained by using methods as described in e.g. Harlow and Lane (1988). In the case of antibodies directed against small peptides such as fragments of a protein of the invention, said peptides are generally coupled to a carrier protein before immunization of animals. Such protein carriers include keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA), ovalbumin and Tetanus toxoid. The carrier protein enhances the immune response of the animal and provides epitopes for T-cell receptor binding sites. The term "antibodies" furthermore includes derivatives thereof such as labelled antibodies. Antibody labels include alkaline phosphatase,

PKH2, PKH26, PKH67, fluorescein (FITC), Hoechst 33258, R-phycoerythrin (PE), rhodamine (TRITC), Quantum Red, Texas Red, Cy3, biotin, agarose, peroxidase and gold spheres. Tools in molecular biology relying on antibodies against a protein include protein gel blot analysis, screening of expression libraries allowing gene identification, protein quantitative methods including ELISA and RIA, immunoaffinity purification of proteins, immunoprecipitation of proteins (see e.g. Example 6) and immunolocalization. Other uses of antibodies and especially of peptide antibodies include the study of proteolytic processing (Loffler et al. 1994, Woulfe et al. 1994), determination of protein active sites (Lerner 1982), the study of precursor and post-translational processing (Baron and Baltimore 1982, Lerner et al. 1981, Semier et al. 1982), identification of protein domains involved in protein-protein interactions (Murakami et al. 1992) and the study of exon usage in gene expression (Tamura et al. 1991).

Embodied in the current invention are antibodies specifically recognizing a cytokinin oxidase or homologue, derivative or fragment thereof as defined supra. Preferably said cytokinin oxidase is a plant cytokinin oxidase, more specifically one of the *Arabidopsis thaliana* cytokinin oxidases (AtCKX).

The terms "gene(s)", "polynucleotide(s)", "nucleic acid(s)", "nucleic acid sequence(s)", "nucleotide sequence(s)", or "nucleic acid molecule(s)", when used herein refer to nucleotides, either ribonucleotides or deoxyribonucleotides or a combination of both, in a polymeric form of any length. Said terms furthermore include double-stranded and single-stranded DNA and RNA. Said terms also include known nucleotide modifications such as methylation, cyclization and 'caps' and substitution of one or more of the naturally occurring nucleotides with an analog such as inosine. Modifications of nucleotides include the addition of acridine, amine, biotin, cascade blue, cholesterol, Cy3®, Cy5®, Cy5.5® Dabcyl, digoxigenin, dinitrophenyl, Edans, 6-FAM, fluorescein, 3'-glyceryl, HEX, IRD-700, IRD-800, JOE, phosphate psoralen, rhodamine, ROX, thiol (SH), spacers, TAMRA, TET, AMCA-S®, SE, BODIPY®, Marina Blue®, Pacific Blue®, Oregon Green®, Rhodamine Green®, Rhodamine Red®, Rhodol Green® and Texas Red®. Polynucleotide backbone modifications include methylphosphonate, 2'-OMe-methylphosphonate RNA, phosphorothiorate, RNA, 2'-OMeRNA. Base modifications include 2-amino-dA, 2-aminopurine, 3'-(ddA), 3'dA(cordycepin), 7-deaza-dA, 8-Br-dA, 8-oxo-dA, N⁶-Me-dA, abasic site (dSpacer), biotin dT, 2'-OMe-5Me-C, 2'-OMe-propynyl-C, 3'-(5-Me-dC), 3'-(ddC), 5-Br-dC, 5-I-dC, 5-Me-dC, 5-F-dC, carboxy-dT, convertible dA,

convertible dC, convertible dG, convertible dT, convertible dU, 7-deaza-dG, 8-Br-dG, 8-oxo-dG, O⁶-Me-dG, S6-DNP-dG, 4-methyl-indole, 5-nitroindole, 2'-OMe-inosine, 2'-dl, O⁶-phenyl-dl, 4-methyl-indole, 2'-deoxynebularine, 5-nitroindole, 2-aminopurine, dP(purine analogue), dK(pyrimidine analogue), 3-nitropyrrole, 2-thio-dT, 4-thio-dT, biotin-dT, carboxy-dT, O⁴-Me-dT, O⁴-triazol dT, 2'-OMe-propynyl-U, 5-Br-dU, 2'-dU, 5-F-dU, 5-I-dU, O⁴-triazol dU. Said terms also encompass peptide nucleic acids (PNAs), a DNA analogue in which the backbone is a pseudopeptide consisting of N-(2-aminoethyl)-glycine units rather than a sugar. PNAs mimic the behaviour of DNA and bind complementary nucleic acid strands. The neutral backbone of PNA results in stronger binding and greater specificity than normally achieved. In addition, the unique chemical, physical and biological properties of PNA have been exploited to produce powerful biomolecular tools, antisense and antigene agents, molecular probes and biosensors.

The present invention also advantageously provides nucleic acid sequences of at least approximately 15 contiguous nucleotides of a nucleic acid according to the invention and preferably from 15 to 50 nucleotides. These sequences may, advantageously be used as probes to specifically hybridise to sequences of the invention as defined above or primers to initiate specific amplification or replication of sequences of the invention as defined above, or the like. Such nucleic acid sequences may be produced according to techniques well known in the art, such as by recombinant or synthetic means. They may also be used in diagnostic kits or the like for detecting the presence of a nucleic acid according to the invention. These tests generally comprise contacting the probe with the sample under hybridising conditions and detecting the presence of any duplex or triplex formation between the probe and any nucleic acid in the sample.

Advantageously, the nucleic acid sequences, according to the invention may be produced using such recombinant or synthetic means, such as for example using PCR cloning mechanisms which generally involve making a pair of primers, which may be from approximately 15 to 50 nucleotides to a region of the gene which is desired to be cloned, bringing the primers into contact with mRNA, cDNA or genomic DNA from a cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the amplified region or fragment and recovering the amplified DNA. Generally, such techniques as defined herein are well known in the art, such as described in Sambrook et al. (Molecular Cloning: a Laboratory Manual, 1989).

A "coding sequence" or "open reading frame" or "ORF" is defined as a nucleotide sequence that can be transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate control sequences or regulatory sequences, i.e. when said coding sequence or ORF is present in an expressible format. Said coding
5 sequence of ORF is bounded by a 5' translation start codon and a 3' translation stop codon. A coding sequence or ORF can include, but is not limited to RNA, mRNA, cDNA, recombinant nucleotide sequences, synthetically manufactured nucleotide sequences or genomic DNA. Said coding sequence or ORF can be interrupted by intervening nucleic acid sequences.

10 Genes and coding sequences essentially encoding the same protein but isolated from different sources can consist of substantially divergent nucleic acid sequences. Reciprocally, substantially divergent nucleic acid sequences can be designed to effect expression of essentially the same protein. Said nucleic acid sequences are the result of e.g. the existence of different alleles of a given gene, of the degeneracy of the genetic
15 code or of differences in codon usage. Thus, as indicated in Table 2, amino acids such as methionine and tryptophan are encoded by a single codon whereas other amino acids such as arginine, leucine and serine can each be translated from up to six different codons. Differences in preferred codon usage are illustrated in Table 3 for *Agrobacterium tumefaciens* (a bacterium), *A. thaliana*, *M. sativa* (two dicotyledonous
20 plants) and *Oryza sativa* (a monocotyledonous plant). To extract one example, the codon GGC (for glycine) is the most frequently used codon in *A. tumefaciens* (36.2 %), is the second most frequently used codon in *O. sativa* but is used at much lower frequencies in *A. thaliana* and *M. sativa* (9 % and 8.4 %, respectively). Of the four possible codons encoding glycine (see Table 2), said GGC codon is most preferably
25 used in *A. tumefaciens* and *O. sativa*. However, in *A. thaliana* this is the GGA (and GGU) codon whereas in *M. sativa* this is the GGU (and GGA) codon.

DNA sequences as defined in the current invention can be interrupted by intervening sequences. With "intervening sequences" is meant any nucleic acid sequence which disrupts a coding sequence comprising said inventive DNA sequence or which disrupts
30 the expressible format of a DNA sequence comprising said inventive DNA sequence. Removal of the intervening sequence restores said coding sequence or said expressible format. Examples of intervening sequences include introns and mobilizable DNA

sequences such as transposons. With "mobilizable DNA sequence" is meant any DNA sequence that can be mobilized as the result of a recombination event.

Table 2. Degeneracy of the genetic code.

Amino Acid	Three-letter code	One-letter code	Possible codons					
Alanine	Ala	A	GCA	GCC	GCG	GCU		
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU
Asparagine	Asn	N	AAC	AAU				
Aspartic Acid	Asp	D	GAC	GAU				
Cysteine	Cys	C	UGC	UGU				
Glutamic Acid	Glu	E	GAA	GAG				
Glutamine	Gln	Q	CAA	CAG				
Glycine	Gly	G	GGA	GGC	GGG	GGU		
Histidine	His	H	CAC	CAU				
Isoleucine	Ile	I	AUA	AUC	AUU			
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU
Lysine	Lys	K	AAA	AAG				
Methionine	Met	M	AUG					
Phenylalanine	Phe	F	UUC	UUU				
Proline	Pro	P	CCA	CCC	CCG	CCU		
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU
Threonine	Thr	T	ACA	ACC	ACG	ACU		
Tryptophan	Trp	W	UGG					
Tyrosine	Tyr	Y	UAC	UAU				
Valine	Val	V	GUA	GUC	GUG	GUU		
Possible "STOP" codons								
			UAA	UAG	UGA			

Table 3. Usage of the indicated codons in the different organisms given as frequency per thousand codons (<http://www.kazusa.or.jp/codon>).

Codon	<i>Agrobacterium tumefaciens</i>	<i>Arabidopsis thaliana</i>	<i>Medicago sativa</i>	<i>Oryza sativa</i>
UUU	13.9	22.5	24.1	11.3
UUC	24.3	20.7	16.9	26.3
UUA	3.5	12.9	10.4	4.7
UUG	13.2	21.0	22.4	11.8

UCU	7.0	24.6	19.8	10.1
UCC	14.8	10.8	7.7	16.9
UCA	7.4	17.8	17.2	9.7
UCG	18.2	8.9	3.2	10.8
UAU	12.3	15.2	16.6	9.2
UAC	10.3	13.7	14.0	20.6
UAA	0.9	0.9	1.2	0.9
UAG	0.6	0.5	0.8	0.8
UGU	3.0	10.8	10.6	5.0
UGC	7.4	7.2	5.8	14.3
UGA	1.8	1.0	0.8	1.3
UGG	12.2	12.7	10.0	12.8
CUU	19.1	24.3	28.3	14.6
CUC	25.7	15.9	12.0	28.0
CUA	5.2	10.0	8.8	5.7
CUG	31.6	9.9	8.5	22.1
CCU	7.7	18.3	23.2	11.8
CCC	10.6	5.3	5.3	12.5
CCA	8.9	16.1	22.6	12.2
CCG	20.7	8.3	3.6	16.7
CAU	10.6	14.0	14.6	9.2
CAC	9.1	8.7	9.1	14.6
CAA	11.2	19.7	23.2	11.9
CAG	24.9	15.2	12.3	24.6
CGU	12.2	8.9	10.1	6.8
CGC	25.5	3.7	4.2	15.9
CGA	8.2	6.2	4.2	4.2
CGG	13.2	4.8	1.8	9.7
AUU	15.4	22.0	29.4	13.8
AUC	36.9	18.5	14.7	25.5
AUA	6.2	12.9	11.7	7.2
AUG	24.7	24.5	21.7	24.4
ACU	6.4	17.8	20.8	10.3
ACC	20.9	10.3	11.7	18.6
ACA	9.1	15.9	18.9	10.0
ACG	18.8	7.6	2.8	10.8
AAU	13.5	22.7	25.0	12.9
AAC	18.7	20.9	18.7	25.1
AAA	13.6	31.0	32.2	12.0
AAG	24.4	32.6	35.1	39.4
AGU	5.7	14.0	12.6	7.3
AGC	15.8	11.1	8.8	16.9
AGA	5.3	18.7	13.6	7.7
AGG	6.5	10.9	11.7	14.9
GUU	16.6	27.3	34.7	15.0
GUC	29.3	12.7	9.9	22.8
GUA	6.1	10.1	10.0	5.7

GUG	19.7	17.5	16.5	25.0
GCU	17.4	28.0	34.6	19.8
GCC	35.8	10.3	11.4	33.2
GCA	19.5	17.6	25.9	15.6
GCG	31.7	8.8	3.4	25.3
GAU	25.8	36.8	40.0	21.5
GAC	28.0	17.3	15.5	31.6
GAA	29.9	34.4	35.9	17.1
GAG	26.3	32.2	27.4	41.1
GGU	16.5	22.2	28.7	16.3
GGC	36.2	9.0	8.4	34.7
GGA	12.5	23.9	27.3	15.0
GGG	11.3	10.2	7.4	16.6

"Hybridization" is the process wherein substantially homologous complementary nucleotide sequences anneal to each other. The hybridization process can occur entirely in solution, i.e. both complementary nucleic acids are in solution. Tools in molecular biology relying on such a process include PCR, subtractive hybridization and DNA sequence determination. The hybridization process can also occur with one of the complementary nucleic acids immobilized to a matrix such as magnetic beads, Sepharose beads or any other resin. Tools in molecular biology relying on such a process include the isolation of poly (A+) mRNA. The hybridization process can furthermore occur with one of the complementary nucleic acids immobilized to a solid support such as a nitrocellulose or nylon membrane or immobilized by e.g. photolithography to e.g. a silicious glass support (the latter known as nucleic acid arrays or microarrays or as nucleic acid chips). Tools in molecular biology relying on such a process include RNA and DNA gel blot analysis, colony hybridization, plaque hybridization and microarray hybridization. In order to allow hybridization to occur, the nucleic acid molecules are generally thermally or chemically (e.g. by NaOH) denatured to melt a double strand into two single strands and/or to remove hairpins or other secondary structures from single stranded nucleic acids. The stringency of hybridization is influenced by conditions such as temperature, salt concentration and hybridization buffer composition. High stringency conditions for hybridization include high temperature and/or low salt concentration (salts include NaCl and Na₃-citrate) and/or the inclusion of formamide in the hybridization buffer and/or lowering the concentration of compounds such as SDS (detergent) in the hybridization buffer and/or exclusion of compounds such as dextran sulfate or polyethylene glycol (promoting molecular crowding) from the

hybridization buffer. Conventional hybridization conditions are described in e.g. Sambrook et al. (1989) but the skilled craftsman will appreciate that numerous different hybridization conditions can be designed in function of the known or the expected homology and/or length of the nucleic acid sequence. Sufficiently low stringency hybridization conditions are particularly preferred to isolate nucleic acids heterologous to the DNA sequences of the invention defined supra. Elements contributing to said heterology include allelism, degeneration of the genetic code and differences in preferred codon usage as discussed supra.

Clearly, the current invention embodies the use of the inventive DNA sequences encoding a cytokinin oxidase, homologue, derivative or immunologically active and/or functional fragment thereof as defined higher in any method of hybridization. The current invention furthermore also relates to DNA sequences hybridizing to said inventive DNA sequences. Preferably said cytokinin oxidase is a plant cytokinin oxidase, more specifically the *Arabidopsis thaliana* (At)CKX.

To effect expression of a protein in a cell, tissue or organ, preferably of plant origin, either the protein may be introduced directly to said cell, such as by microinjection or ballistic means or alternatively, an isolated nucleic acid molecule encoding said protein may be introduced into said cell, tissue or organ in an expressible format.

Preferably, the DNA sequence of the invention comprises a coding sequence or open reading frame (ORF) encoding a cytokinin oxidase protein or a homologue or derivative thereof or an immunologically active and/or functional fragment thereof as defined supra. The preferred protein of the invention comprises the amino acid sequence of said cytokinin oxidase. Preferably said cytokinin oxidase is a plant cytokinin oxidase and more specifically a *Arabidopsis thaliana* (At)CKX.

With "vector" or "vector sequence" is meant a DNA sequence which can be introduced in an organism by transformation and can be stably maintained in said organism. Vector maintenance is possible in e.g. cultures of *Escherichia coli*, *A. tumefaciens*, *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*. Other vectors such as phagemids and cosmid vectors can be maintained and multiplied in bacteria and/or viruses. Vector sequences generally comprise a set of unique sites recognized by restriction enzymes, the multiple cloning site (MCS), wherein one or more non-vector sequence(s) can be inserted.

With "non-vector sequence" is accordingly meant a DNA sequence which is integrated in one or more of the sites of the MCS comprised within a vector.

"Expression vectors" form a subset of vectors which, by virtue of comprising the appropriate regulatory or control sequences enable the creation of an expressible format for the inserted non-vector sequence(s), thus allowing expression of the protein encoded by said non-vector sequence(s). Expression vectors are known in the art enabling protein expression in organisms including bacteria (e.g. *E. coli*), fungi (e.g. *S. cerevisiae*, *S. pombe*, *Pichia pastoris*), insect cells (e.g. baculoviral expression vectors), animal cells (e.g. COS or CHO cells) and plant cells (e.g. potato virus X-based expression vectors).

The current invention clearly includes any cytokinin oxidase, homologue, derivative and/or immunologically active and/or functional fragment thereof as defined supra. Preferably said cytokinin oxidase is a plant cytokinin oxidase, more specifically a *Arabidopsis thaliana* (At)CKX.

As an alternative to expression vector-mediated protein production in biological systems, chemical protein synthesis can be applied. Synthetic peptides can be manufactured in solution phase or in solid phase. Solid phase peptide synthesis (Merrifield 1963) is, however, the most common way and involves the sequential addition of amino acids to create a linear peptide chain. Solid phase peptide synthesis includes cycles consisting of three steps: (i) immobilization of the carboxy-terminal amino acid of the growing peptide chain to a solid support or resin; (ii) chain assembly, a process consisting of activation, coupling and deprotection of the amino acid to be added to the growing peptide chain; and (iii) cleavage involving removal of the completed peptide chain from the resin and removal of the protecting groups from the amino acid side chains. Common approaches in solid phase peptide synthesis include Fmoc/tBu (9-fluorenylmethyloxycarbonyl/t-butyl) and Boc (t-butyloxycarbonyl) as the amino-terminal protecting groups of amino acids. Amino acid side chain protecting groups include methyl (Me), formyl (CHO), ethyl (Et), acetyl (Ac), t-butyl (t-Bu), anisyl, benzyl (Bzl), trifluoroacetyl (Tfa), N-hydroxysuccinimide (ONSu, OSu), benzoyl (Bz), 4-methylbenzyl (Meb), thioanisyl, thiocresyl, benzyloxymethyl (Bom), 4-nitrophenyl (ONp), benzyloxycarbonyl (Z), 2-nitrobenzoyl (NBz), 2-nitrophenylsulphenyl (Nps), 4-toluenesulphonyl (Tosyl, Tos), pentafluorophenyl (Pfp), diphenylmethyl (Dpm), 2-chlorobenzyloxycarbonyl (Cl-Z), 2,4,5-trichlorophenyl, 2-bromobenzyloxycarbonyl (Br-Z), triphenylmethyl (Trityl, Trt), and 2,5,7,8-pentamethylchroman-6-sulphonyl (Pmc). During chain assembly, Fmoc or Boc are removed resulting

in an activated amino-terminus of the amino acid residue bound to the growing chain. The carboxy-terminus of the incoming amino acid is activated by conversion into a highly reactive ester, e.g. by HBTU. With current technologies (e.g. PerSeptive Biosystems 9050 synthesizer, Applied Biosystems Model 431A Peptide Synthesizer), linear peptides
5 of up to 50 residues can be manufactured. A number of guidelines is available to produce peptides that are suitable for use in biological systems including (i) limiting the use of difficult amino acids such as cys, met, trp (easily oxidized and/or degraded during peptide synthesis) or arg; (ii) minimize hydrophobic amino acids (can impair peptide solubility); and (iii) prevent an amino-terminal glutamic acid (can cyclize to
10 pyroglutamate).

By "expressible format" is meant that the isolated nucleic acid molecule is in a form suitable for being transcribed into mRNA and/or translated to produce a protein, either constitutively or following induction by an intracellular or extracellular signal, such as an environmental stimulus or stress (mitogens, anoxia, hypoxia, temperature, salt, light,
15 dehydration, etc) or a chemical compound such as IPTG (isopropyl- β -D-thiogalactopyranoside) or such as an antibiotic (tetracycline, ampicillin, rifampicin, kanamycin), hormone (e.g. gibberellin, auxin, cytokinin, glucocorticoid, brassinosteroid, ethylene, abscisic acid etc), hormone analogue (indolacetic acid (IAA), 2,4-D, etc) , metal (zinc, copper, iron, etc), or dexamethasone, amongst others. As will be known to
20 those skilled in the art, expression of a functional protein may also require one or more post-translational modifications, such as glycosylation, phosphorylation, dephosphorylation, or one or more protein-protein interactions, amongst others. All such processes are included within the scope of the term "expressible format".

Preferably, expression of a protein in a specific cell, tissue, or organ, preferably of plant
25 origin, is effected by introducing and expressing an isolated nucleic acid molecule encoding said protein, such as a cDNA molecule, genomic gene, synthetic oligonucleotide molecule, mRNA molecule or open reading frame, to said cell, tissue or organ, wherein said nucleic acid molecule is placed operably in connection with suitable regulatory or control sequences including a promoter, preferably a plant-expressible
30 promoter, and a terminator sequence.

Reference herein to a "promoter" is to be taken in its broadest context and includes the transcriptional regulatory sequences derived from a classical eukaryotic genomic gene, including the TATA box which is required for accurate transcription initiation, with or

without a CCAAT box sequence and additional regulatory or control elements (i.e. upstream activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner.

The term "promoter" also includes the transcriptional regulatory sequences of a classical prokaryotic gene, in which case it may include a -35 box sequence and/or a -10 box transcriptional regulatory sequences.

The term "promoter" is also used to describe a synthetic or fusion molecule, or derivative which confers, activates or enhances expression of a nucleic acid molecule in a cell, tissue or organ.

Promoters may contain additional copies of one or more specific regulatory elements, to further enhance expression and/or to alter the spatial expression and/or temporal expression of a nucleic acid molecule to which it is operably connected. Such regulatory elements may be placed adjacent to a heterologous promoter sequence to drive expression of a nucleic acid molecule in response to e.g. copper, glucocorticoids, dexamethasone, tetracycline, gibberellin, cAMP, abscisic acid, auxin, wounding, ethylene, jasmonate or salicylic acid or to confer expression of a nucleic acid molecule to specific cells, tissues or organs such as meristems, leaves, roots, embryo, flowers, seeds or fruits.

In the context of the present invention, the promoter preferably is a plant-expressible promoter sequence. Promoters that also function or solely function in non-plant cells such as bacteria, yeast cells, insect cells and animal cells are not excluded from the invention. By "plant-expressible" is meant that the promoter sequence, including any additional regulatory elements added thereto or contained therein, is at least capable of inducing, conferring, activating or enhancing expression in a plant cell, tissue or organ, preferably a monocotyledonous or dicotyledonous plant cell, tissue, or organ.

The terms "plant-operable" and "operable in a plant" when used herein, in respect of a promoter sequence, shall be taken to be equivalent to a plant-expressible promoter sequence.

Regulatable promoters as part of a binary viral plant expression system are also known to the skilled artisan (Yadav 1999 – WO9922003; Yadav 2000 – WO0017365).

In the present context, a "regulatable promoter sequence" is a promoter that is capable of conferring expression on a structural gene in a particular cell, tissue, or organ or group of cells, tissues or organs of a plant, optionally under specific conditions, however does

generally not confer expression throughout the plant under all conditions. Accordingly, a regulatable promoter sequence may be a promoter sequence that confers expression on a gene to which it is operably connected in a particular location within the plant or alternatively, throughout the plant under a specific set of conditions, such as following
5 induction of gene expression by a chemical compound or other elicitor.

Preferably, the regulatable promoter used in the performance of the present invention confers expression in a specific location within the plant, either constitutively or following induction, however not in the whole plant under any circumstances. Included within the scope of such promoters are cell-specific promoter sequences, tissue-specific promoter
10 sequences, organ-specific promoter sequences, cell cycle specific gene promoter sequences, inducible promoter sequences and constitutive promoter sequences that have been modified to confer expression in a particular part of the plant at any one time, such as by integration of said constitutive promoter within a transposable genetic element (*Ac*, *Ds*, *Spm*, *En*, or other transposon).

15 Similarly, the term "tissue-specific" shall be taken to indicate that expression is predominantly in a particular tissue or tissue-type, preferably of plant origin, albeit not necessarily exclusively in said tissue or tissue-type.

Similarly, the term "organ-specific" shall be taken to indicate that expression is predominantly in a particular organ, preferably of plant origin, albeit not necessarily
20 exclusively in said organ.

Similarly, the term "cell cycle specific" shall be taken to indicate that expression is predominantly cyclic and occurring in one or more, not necessarily consecutive phases of the cell cycle albeit not necessarily exclusively in cycling cells, preferably of plant origin.

25 Those skilled in the art will be aware that an "inducible promoter" is a promoter the transcriptional activity of which is increased or induced in response to a developmental, chemical, environmental, or physical stimulus. Similarly, the skilled craftsman will understand that a "constitutive promoter" is a promoter that is transcriptionally active throughout most, but not necessarily all parts of an organism, preferably a plant, during
30 most, but not necessarily all phases of its growth and development.

Those skilled in the art will readily be capable of selecting appropriate promoter sequences for use in regulating appropriate expression of the cytokinin oxidase protein from publicly-available or readily-available sources, without undue experimentation.

Placing a nucleic acid molecule under the regulatory control of a promoter sequence, or in operable connection with a promoter sequence, means positioning said nucleic acid molecule such that expression is controlled by the promoter sequence. A promoter is usually, but not necessarily, positioned upstream, or at the 5'-end, and within 2 kb of the start site of transcription, of the nucleic acid molecule which it regulates. In the construction of heterologous promoter/structural gene combinations it is generally preferred to position the promoter at a distance from the gene transcription start site that is approximately the same as the distance between that promoter and the gene it controls in its natural setting (i.e., the gene from which the promoter is derived). As is known in the art, some variation in this distance can be accommodated without loss of promoter function. Similarly, the preferred positioning of a regulatory sequence element with respect to a heterologous gene to be placed under its control is defined by the positioning of the element in its natural setting (i.e., the gene from which it is derived). Again, as is known in the art, some variation in this distance can also occur.

Examples of promoters suitable for use in gene constructs of the present invention include those listed in Table 4, amongst others. The promoters listed in Table 4 are provided for the purposes of exemplification only and the present invention is not to be limited by the list provided therein. Those skilled in the art will readily be in a position to provide additional promoters that are useful in performing the present invention.

In the case of constitutive promoters or promoters that induce expression throughout the entire plant, it is preferred that such sequences are modified by the addition of nucleotide sequences derived from one or more of the tissue-specific promoters listed in Table 4, or alternatively, nucleotide sequences derived from one or more of the above-mentioned tissue-specific inducible promoters, to confer tissue-specificity thereon. For example, the CaMV 35S promoter may be modified by the addition of maize *Adh1* promoter sequence, to confer anaerobically-regulated root-specific expression thereon, as described previously (Ellis *et al.*, 1987). Another example describes conferring root specific or root abundant gene expression by fusing the CaMV35S promoter to elements of the maize glycine-rich protein GRP3 gene (Feix and Wulff 2000 - WO0015662). Such modifications can be achieved by routine experimentation by those skilled in the art.

The term "terminator" refers to a DNA sequence at the end of a transcriptional unit which signals termination of transcription. Terminators are 3'-non-translated DNA sequences containing a polyadenylation signal, which facilitates the addition of polyadenylate

sequences to the 3'-end of a primary transcript. Terminators active in cells derived from viruses, yeasts, moulds, bacteria, insects, birds, mammals and plants are known and described in the literature. They may be isolated from bacteria, fungi, viruses, animals and/or plants.

Table 4. Exemplary plant-expressible promoters for use in the performance of the present invention

I: CELL-SPECIFIC, TISSUE-SPECIFIC, AND ORGAN-SPECIFIC PROMOTERS		
GENE SOURCE	EXPRESSION PATTERN	REFERENCE
α -amylase (<i>Amy32b</i>)	aleurone	Lanahan, M.B., <i>et al.</i> , <i>Plant Cell</i> 4:203-211, 1992; Skriver, K., <i>et al.</i> <i>Proc. Natl. Acad. Sci. (USA)</i> 88: 7266-7270, 1991
cathepsin β -like gene	aleurone	Cejudo, F.J., <i>et al.</i> <i>Plant Molecular Biology</i> 20:849-856, 1992.
<i>Agrobacterium rhizogenes</i> <i>rolB</i>	cambium	Nilsson <i>et al.</i> , <i>Physiol. Plant.</i> 100:456-462, 1997
AtPRP4	flowers	http://salus.medium.edu/mmg/tierney/html
chalcone synthase (<i>chsA</i>)	flowers	Van der Meer, <i>et al.</i> , <i>Plant Mol. Biol.</i> 15, 95-109, 1990.
LAT52	anther	Twell <i>et al.</i> <i>Mol. Gen Genet.</i> 217:240-245 (1989)
<i>apetala-3</i>	flowers	
chitinase	fruit (berries, grapes, etc)	Thomas <i>et al.</i> CSIRO Plant Industry, Urrbrae, South Australia, Australia; http://winetitles.com.au/gwrdc/csh95-1.html
<i>rbcS-3A</i>	green tissue (eg leaf)	Lam, E. <i>et al.</i> , <i>The Plant Cell</i> 2: 857-866, 1990.; Tucker <i>et al.</i> , <i>Plant Physiol.</i> 113: 1303-1308, 1992.
leaf-specific genes	leaf	Baszczynski, <i>et al.</i> , <i>Nucl. Acid Res.</i> 16: 4732, 1988.
AtPRP4	leaf	http://salus.medium.edu/mmg/tierney/html
chlorella virus adenine methyltransferase gene promoter	leaf	Mitra and Higgins, 1994, <i>Plant Molecular Biology</i> 26: 85-93
aldP gene promoter from rice	leaf	Kagaya <i>et al.</i> , 1995, <i>Molecular and General Genetics</i> 248: 668-674
<i>rbcS</i> promoter from rice or tomato	leaf	Kyozuka <i>et al.</i> , 1993, <i>Plant Physiology</i> 102: 991-1000
<i>Pinus cab-6</i>	leaf	Yamamoto <i>et al.</i> , <i>Plant Cell Physiol.</i> 35:773-778, 1994.
rubisco promoter	leaf	

cab (chlorophyll a/b-binding protein)	leaf	
SAM22	senescent leaf	Crowell, <i>et al.</i> , <i>Plant Mol. Biol.</i> 18: 459-466, 1992.
<i>ltp</i> gene (lipid transfer gene)		Fleming, <i>et al.</i> , <i>Plant J.</i> 2, 855-862.
<i>R. japonicum nif</i> gene	Nodule	United States Patent No. 4, 803, 165
<i>B. japonicum nifH</i> gene	Nodule	United States Patent No. 5, 008, 194
GmENOD40	Nodule	Yang, <i>et al.</i> , <i>The Plant J.</i> 3: 573-585.
PEP carboxylase (PEPC)	Nodule	Pathirana, <i>et al.</i> , <i>Plant Mol. Biol.</i> 20: 437-450, 1992.
leghaemoglobin (Lb)	Nodule	Gordon, <i>et al.</i> , <i>J. Exp. Bot.</i> 44: 1453-1465, 1993.
<i>Tungro bacilliform</i> virus gene	phloem	Bhattacharyya-Pakrasi, <i>et al.</i> , <i>The Plant J.</i> 4: 71-79, 1992.
pollen-specific genes	pollen; microspore	Albani, <i>et al.</i> , <i>Plant Mol. Biol.</i> 15: 605, 1990; Albani, <i>et al.</i> , <i>Plant Mol. Biol.</i> 16: 501, 1991)
Zm13	pollen	Guerrero <i>et al.</i> <i>Mol. Gen. Genet.</i> 224:161-168 (1993)
apg gene	microspore	Twell <i>et al.</i> <i>Sex. Plant Reprod.</i> 6:217-224 (1993)
maize pollen-specific gene	pollen	Hamilton, <i>et al.</i> , <i>Plant Mol. Biol.</i> 18: 211-218, 1992.
sunflower pollen-expressed gene	pollen	Baltz, <i>et al.</i> , <i>The Plant J.</i> 2: 713-721, 1992.
<i>B. napus</i> pollen-specific gene	pollen; anther; tapetum	Arnoldo, <i>et al.</i> , <i>J. Cell. Biochem.</i> , Abstract No. Y101, 204, 1992.
root-expressible genes	roots	Tingey, <i>et al.</i> , <i>EMBO J.</i> 6: 1, 1987.
tobacco auxin-inducible gene	root tip	Van der Zaal, <i>et al.</i> , <i>Plant Mol. Biol.</i> 16, 983, 1991.
β -tubulin	root	Oppenheimer, <i>et al.</i> , <i>Gene</i> 63: 87, 1988.
tobacco root-specific genes	root	Conkling, <i>et al.</i> , <i>Plant Physiol.</i> 93: 1203, 1990.
<i>B. napus</i> G1-3b gene	root	United States Patent No. 5, 401, 836
SbPRP1	roots	Suzuki <i>et al.</i> , <i>Plant Mol. Biol.</i> 21: 109-119, 1993.
AtPRP1; AtPRP3	roots; root hairs	http://salus.medium.edu/mmg/tierney/html
RD2 gene	root cortex	http://www2.cnsu.edu/ncsu/research
TobRB7 gene	root vasculature	http://www2.cnsu.edu/ncsu/research

AtPRP4	leaves; flowers; lateral root primordia	http://salus.medium.edu/mmg/tierney/html
seed-specific genes	seed	Simon, <i>et al.</i> , <i>Plant Mol. Biol.</i> 5: 191, 1985; Scofield, <i>et al.</i> , <i>J. Biol. Chem.</i> 262: 12202, 1987.; Baszczynski, <i>et al.</i> , <i>Plant Mol. Biol.</i> 14: 633, 1990.
Brazil Nut albumin	seed	Pearson, <i>et al.</i> , <i>Plant Mol. Biol.</i> 18: 235-245, 1992.
legumin	seed	Ellis, <i>et al.</i> , <i>Plant Mol. Biol.</i> 10: 203-214, 1988.
glutelin (rice)	seed	Takaiwa, <i>et al.</i> , <i>Mol. Gen. Genet.</i> 208: 15-22, 1986; Takaiwa, <i>et al.</i> , <i>FEBS Letts.</i> 221: 43-47, 1987.
zein	seed	Matzke <i>et al</i> <i>Plant Mol Biol</i> , 14(3):323-32 1990
napA	seed	Stalberg, <i>et al</i> , <i>Planta</i> 199: 515-519, 1996.
wheat LMW and HMW glutenin-1	endosperm	<i>Mol Gen Genet</i> 216:81-90, 1989; <i>NAR</i> 17:461-2, 1989
wheat SPA	seed	Albani <i>et al</i> , <i>Plant Cell</i> , 9: 171-184, 1997
wheat α , β , γ -gliadins	endosperm	<i>EMBO</i> 3:1409-15, 1984
barley <i>ltr1</i> promoter	endosperm	
barley B1, C, D, hordein	endosperm	<i>Theor Appl Gen</i> 98:1253-62, 1999; <i>Plant J</i> 4:343-55, 1993; <i>Mol Gen Genet</i> 250:750-60, 1996
barley DOF	endosperm	Mena <i>et al</i> , <i>The Plant Journal</i> , 116(1): 53-62, 1998
<i>blz2</i>	endosperm	EP99106056.7
synthetic promoter	endosperm	Vicente-Carbajosa <i>et al.</i> , <i>Plant J.</i> 13: 629-640, 1998.
rice prolamin NRP33	endosperm	Wu <i>et al</i> , <i>Plant Cell Physiology</i> 39(8) 885-889, 1998
rice α -globulin Glb-1	endosperm	Wu <i>et al</i> , <i>Plant Cell Physiology</i> 39(8) 885-889, 1998
rice OSH1	embryo	Sato <i>et al</i> , <i>Proc. Natl. Acad. Sci. USA</i> , 93: 8117-8122, 1996
rice α -globulin REB/OHP-1	endosperm	Nakase <i>et al.</i> <i>Plant Mol. Biol.</i> 33: 513-522, 1997
rice ADP-glucose PP	endosperm	<i>Trans Res</i> 6:157-68, 1997
maize ESR gene family	endosperm	<i>Plant J</i> 12:235-46, 1997
sorgum γ -kafirin	endosperm	<i>PMB</i> 32:1029-35, 1996
KNOX	embryo	Postma-Haarsma <i>et al</i> , <i>Plant Mol. Biol.</i> 39:257-71, 1999

rice oleosin	embryo and aleuron	Wu <i>et al.</i> , J. Biochem., 123:386, 1998
sunflower oleosin	seed (embryo and dry seed)	Cummins, <i>et al.</i> , Plant Mol. Biol. 19: 873-876, 1992
LEAFY	shoot meristem	Weigel <i>et al.</i> , Cell 69:843-859, 1992.
<i>Arabidopsis thaliana knat1</i>	shoot meristem	Accession number AJ131822
<i>Malus domestica kn1</i>	shoot meristem	Accession number Z71981
CLAVATA1	shoot meristem	Accession number AF049870
stigma-specific genes	stigma	Nasrallah, <i>et al.</i> , Proc. Natl. Acad. Sci. USA 85: 5551, 1988; Trick, <i>et al.</i> , Plant Mol. Biol. 15: 203, 1990.
class I patatin gene	tuber	Liu <i>et al.</i> , Plant Mol. Biol. 153:386-395, 1991.
PCNA rice	meristem	Kosugi <i>et al.</i> , Nucleic Acids Research 19:1571-1576, 1991; Kosugi S. and Ohashi Y, Plant Cell 9:1607-1619, 1997.
Pea TubA1 tubulin	Dividing cells	Stotz and Long, Plant Mol.Biol. 41, 601-614. 1999
<i>Arabidopsis cdc2a</i>	cycling cells	Chung and Parish, FEBS Lett, 3:362(2):215-9, 1995
<i>Arabidopsis Rop1A</i>	Anthers; mature pollen + pollen tubes	Li <i>et al.</i> 1998 Plant Physiol 118, 407-417.
<i>Arabidopsis AtDMC1</i>	Meiosis-associated	Klimyuk and Jones 1997 Plant J. 11, 1-14.
Pea PS-IAA4/5 and PS-IAA6	Auxin-inducible	Wong <i>et al.</i> 1996 Plant J. 9, 587-599.
Pea farnesyltransferase	Meristematic tissues; phloem near growing tissues; light- and sugar-repressed	Zhou <i>et al.</i> 1997 Plant J. 12, 921-930
Tobacco (<i>N. sylvestris</i>) cyclin B1;1	Dividing cells / meristematic tissue	Trehin <i>et al.</i> 1997 Plant Mol.Biol. 35, 667-672.
<i>Catharanthus roseus</i> Mitotic cyclins CYS (A-type) and CYM (B-type)	Dividing cells / meristematic tissue	Ito <i>et al.</i> 1997 Plant J. 11, 983-992
<i>Arabidopsis cyc1At</i> (=cyc B1;1) and cyc3aAt (A-type)	Dividing cells / meristematic tissue	Shaul <i>et al.</i> 1996 Proc.Natl.Acad.Sci.U.S.A 93, 4868-4872.
<i>Arabidopsis tef1</i> promoter box	Dividing cells / meristematic tissue	Regad <i>et al.</i> 1995 Mol.Gen.Genet. 248, 703-711.
<i>Catharanthus roseus cyc07</i>	Dividing cells / meristematic tissue	Ito <i>et al.</i> 1994 Plant Mol.Biol. 24, 863-878.

Table 4 (continued). Exemplary plant-expressible promoters for use in the performance of the present invention

II: EXEMPLARY CONSTITUTIVE PROMOTERS		
GENE SOURCE	EXPRESSION PATTERN	REFERENCE
Actin	constitutive	McElroy <i>et al</i> , Plant Cell, 2: 163-171, 1990
CAMV 35S	constitutive	Odell <i>et al</i> , Nature, 313: 810-812, 1985
CaMV 19S	constitutive	Nilsson <i>et al.</i> , Physiol. Plant. 100:456-462, 1997
GOS2	constitutive	de Pater <i>et al</i> , Plant J. 2:837-844, 1992
ubiquitin	constitutive	Christensen <i>et al</i> , Plant Mol. Biol. 18: 675-689, 1992
rice cyclophilin	constitutive	Buchholz <i>et al</i> , Plant Mol Biol. 25: 837-843, 1994
maize histone H3	constitutive	Lepetit <i>et al</i> , Mol. Gen. Genet. 231:276-285, 1992
alfalfa histone H3	constitutive	Wu <i>et al.</i> , Nucleic Acids Res. 17: 3057-3063, 1989; Wu <i>et al.</i> , Plant Mol. Biol. 11:641-649, 1988
actin 2	constitutive	An <i>et al</i> , Plant J. 10(1); 107-121, 1996

Table 4 (continued). Exemplary plant-expressible promoters for use in the performance of the present invention

III: EXEMPLARY STRESS-INDUCIBLE PROMOTERS		
NAME	STRESS	REFERENCE
P5CS (delta(1)-pyrroline-5-carboxylate syntase)	salt, water	Zhang et al. Plant Science. 129: 81-89, 1997
cor15a	cold	Hajela et al., Plant Physiol. 93: 1246-1252, 1990
cor15b	cold	Wlihelm et al., Plant Mol Biol. 23:1073-1077, 1993
cor15a (-305 to +78 nt)	cold, drought	Baker et al., Plant Mol Biol. 24: 701-713, 1994
rd29	salt, drought, cold	Kasuga et al., Nature Biotechnology 18:287-291, 1999
heat shock proteins, including artificial promoters containing the heat shock element (HSE)	heat	Barros et al., Plant Mol Biol 19: 665-75, 1992. Marrs et al., Dev Genet. 14: 27-41, 1993. Schoffl et al., Mol Gen Gent, 217: 246-53, 1989.
smHSP (small heat shock proteins)	heat	Waters et al, J Experimental Botany 47:325-338, 1996
wcs120	cold	Ouellet et al., FEBS Lett. 423: 324-328, 1998
ci7	cold	Kirch et al., Plant Mol Biol 33: 897-909, 1997
Adh	cold, drought, hypoxia	Dolferus et al., Plant Physiol 105: 1075-87, 1994
pws18	water: salt and drought	Joshee et al., Plant Cell Physiol 39: 64-72, 1998
ci21A	cold	Schneider et al., Plant Physiol 113: 335-45, 1997
Trg-31	drought	Chaudhary et al., Plant Mol Biol 30: 1247-57, 1996
osmotin	osmotic	Raghothama et al., Plant Mol Biol 23: 1117-28, 1993
Rab17	osmotic, ABA	Vilardell et al., Plant Mol Biol 17: 985-93, 1991
lapA	wounding, enviromental	WO99/03977 University of California/INRA

Table 4 (continued). Exemplary plant-expressible promoters for use in the performance of the present invention

5

IV: EXEMPLARY PATHOGEN-INDUCIBLE PROMOTERS		
NAME	PATHOGEN	REFERENCE
RB7	Root-knot nematodes (Meloidogyne spp.)	US5760386 - North Carolina State University; Opperman et al (1994) <i>Science</i> 263: 221-23.
PR-1, 2, 3, 4, 5, 8, 11	fungal, viral, bacterial	Ward et al (1991) <i>Plant Cell</i> 3: 1085-1094; Reiss et al 1996; Lebel et al (1998), <i>Plant J</i> , 16(2):223-33; Melchers et al (1994), <i>Plant J</i> , 5(4):469-80; Lawton et al (1992), <i>Plant Mol Biol</i> , 19(5):735-43.
HMG2	nematodes	WO9503690 - Virginia Tech Intellectual Properties Inc .
Abi3	Cyst nematodes (Heterodera spp.)	Unpublished
ARM1	nematodes	Barthels et al., (1997) <i>The Plant Cell</i> 9, 2119-2134. WO 98/31822 – Plant Genetic Systems
Att0728	nematodes	Barthels et al., (1997) <i>The Plant Cell</i> 9, 2119-2134. <i>PCT/EP98/07761</i>
Att1712	nematodes	Barthels et al., (1997) <i>The Plant Cell</i> 9, 2119-2134. <i>PCT/EP98/07761</i>
Gst1	Different types of pathogens	Strittmatter et al (1996) <i>Mol. Plant-Microbe Interact.</i> 9, 68-73.
LEMMI	nematodes	WO 92/21757 – Plant Genetic Systems
CLE	geminivirus	<i>PCT/EP99/03445 - CINESTAV</i>
PDF1.2	Fungal including <i>Alternaria brassicicola</i> and <i>Botrytis cinerea</i>	Manners et al (1998), <i>Plant Mol Biol</i> , 38(6):1071-80.
Thi2.1	Fungal – <i>Fusarium oxysporum f sp. matthioli</i>	Vignutelli et al (1998) <i>Plant J</i> ;14(3):285-95
DB#226	nematodes	Bird and Wilson (1994) <i>Mol. Plant-Microbe Interact.</i> , 7, 419-42

		WO 95.322888
DB#280	nematodes	Bird and Wilson (1994) Mol. Plant-Microbe Interact., 7, 419-42 WO 95.322888
Cat2	nematodes	Niebel et al (1995) Mol Plant Microbe Interact 1995 May-Jun;8(3):371-8
□Tub	nematodes	Aristizabal et al (1996), 8 th International Congress on Plant-Microbe Interaction, Knoxville US B-29
SHSP	nematodes	Fenoll et al (1997) In: Cellular and molecular aspects of plant-nematode interactions. Kluwer Academic, C. Fenoll, F.M.W. Grundler and S.A. Ohi (Eds.),
Tsw12	nematodes	Fenoll et al (1997) In: Cellular and molecular aspects of plant-nematode interactions. Kluwer Academic, C. Fenoll, F.M.W. Grundler and S.A. Ohi (Eds.)
Hs1(pro1)	nematodes	WO 98/122335 - Jung
NsLTP	viral, fungal, bacterial	Molina & Garc'ia-Olmedo (1993) FEBS Lett, 316(2):119-22
RIP	viral, fungal	Turner et al (1997) Proc Natl Acad Sci U S A, 94(8):3866-71

Examples of terminators particularly suitable for use in the gene constructs of the present invention include the *Agrobacterium tumefaciens* nopaline synthase (NOS) gene terminator, the *Agrobacterium tumefaciens* octopine synthase (OCS) gene terminator sequence, the Cauliflower mosaic virus (CaMV) 35S gene terminator sequence, the *Oryza sativa* ADP-glucose pyrophosphorylase terminator sequence (t3'Bt2), the *Zea mays* zein gene terminator sequence, the *rbcs-1A* gene terminator, and the *rbcs-3A* gene terminator sequences, amongst others.

Preferred promoter sequences of the invention include root specific promoters such as but not limited to the ones listed in Table 5 and as outlined in the Examples.

Table 5. Exemplary root specific promoters for use in the performance of the present invention

NAME	ORIGIN	REFERENCE
SbPRP1	Soybean	Suzuki et al., Plant Mol Biol, 21: 109-119, 1993
636 bp fragment of TobRB7	Tobacco	Yamamoto et al., Plant Cell 3:371-382, 1991
GGPS3	Arabidopsis	Okada et al., Plant Physiol 122: 1045-1056, 2000
580 bp fragment of prxEa	Arabidopsis	Wanapu and Shinmyo, Ann N Y Acad Sci 782: 107-114, 1996
lds2 promoter	Barley	Okumura et al., Plant Mol Biol 25: 705-719, 1994
AtPRP3	Arabidopsis	Fowler et al., Plant Physiol 121: 1081-1092, 1999

Those skilled in the art will be aware of additional promoter sequences and terminator sequences which may be suitable for use in performing the invention. Such sequences may readily be used without any undue experimentation.

In the context of the current invention, "ectopic expression" or "ectopic overexpression" of a gene or a protein are conferring to expression patterns and/or expression levels of said gene or protein normally not occurring under natural conditions, more specifically is meant increased expression and/or increased expression levels. Ectopic expression can be achieved in a number of ways including operably linking of a coding sequence encoding said protein to an isolated homologous or heterologous promoter in order to create a chimeric gene and/or operably linking said coding sequence to its own isolated promoter (i.e. the unisolated promoter naturally driving expression of said protein) in order to create a recombinant gene duplication or gene multiplication effect. With "ectopic co-expression" is meant the ectopic expression or ectopic overexpression of two or more genes or proteins. The same or, more preferably, different promoters are used to confer ectopic expression of said genes or proteins.

Preferably, the promoter sequence used in the context of the present invention is operably linked to a coding sequence or open reading frame (ORF) encoding a cytokinin oxidase protein or a homologue, derivative or an immunologically active and/or functional fragment thereof as defined supra.

5 "Downregulation of expression" as used herein means lowering levels of gene expression and/or levels of active gene product and/or levels of gene product activity. Decreases in expression may be accomplished by e.g. the addition of coding sequences or parts thereof in a sense orientation (if resulting in co-suppression) or in an antisense orientation relative to a promoter sequence and furthermore by e.g. insertion
10 mutagenesis (e.g. T-DNA insertion or transposon insertion) or by gene silencing strategies as described by e.g. Angell and Baulcombe (1998 - WO9836083), Lowe et al. (1989 - WO9853083), Lederer et al. (1999 - WO9915682) or Wang et al. (1999 - WO9953050). Genetic constructs aimed at silencing gene expression may have the nucleotide sequence of said gene (or one or more parts thereof) contained therein in a
15 sense and/or antisense orientation relative to the promoter sequence. Another method to downregulate gene expression comprises the use of ribozymes.

Modulating, including lowering, the level of active gene products or of gene product activity can be achieved by administering or exposing cells, tissues, organs or organisms to said gene product, a homologue, derivative and/or immunologically active fragment
20 thereof. Immunomodulation is another example of a technique capable of downregulation levels of active gene product and/or of gene product activity and comprises administration of or exposing to or expressing antibodies to said gene product to or in cells, tissues, organs or organisms wherein levels of said gene product and/or gene product activity are to be modulated. Such antibodies comprise "plantibodies",
25 single chain antibodies, IgG antibodies and heavy chain camel antibodies as well as fragments thereof.

Modulating, including lowering, the level of active gene products or of gene product activity can furthermore be achieved by administering or exposing cells, tissues, organs or organisms to an agonist of said gene product or the activity thereof. Such agonists
30 include proteins (comprising e.g. kinases and proteinases) and chemical compounds identified according to the current invention as described supra.

In the context of the current invention is envisaged the downregulation of the expression of a cytokinin oxidase gene as defined higher. Preferably said cytokinin oxidase gene is

a plant cytokinin oxidase gene, more specifically an *AtCKX*. The invention further comprises downregulation of levels of a cytokinin oxidase protein or of a cytokinin oxidase activity whereby said cytokinin oxidase protein has been defined supra. Preferably said cytokinin oxidase protein is a plant cytokinin oxidase, more specifically an *AtCKX*.

By "modifying cell fate and/or plant development and/or plant morphology and/or biochemistry and/or physiology" is meant that one or more developmental and/or morphological and/or biochemical and/or physiological characteristics of a plant is altered by the performance of one or more steps pertaining to the invention described herein.

"Cell fate" refers to the cell-type or cellular characteristics of a particular cell that are produced during plant development or a cellular process therefor, in particular during the cell cycle or as a consequence of a cell cycle process.

"Plant development" or the term "plant developmental characteristic" or similar term shall, when used herein, be taken to mean any cellular process of a plant that is involved in determining the developmental fate of a plant cell, in particular the specific tissue or organ type into which a progenitor cell will develop. Cellular processes relevant to plant development will be known to those skilled in the art. Such processes include, for example, morphogenesis, photomorphogenesis, shoot development, root development, vegetative development, reproductive development, stem elongation, flowering, and regulatory mechanisms involved in determining cell fate, in particular a process or regulatory process involving the cell cycle.

"Plant morphology" or the term "plant morphological characteristic" or similar term will, when used herein, be understood by those skilled in the art to refer to the external appearance of a plant, including any one or more structural features or combination of structural features thereof. Such structural features include the shape, size, number, position, colour, texture, arrangement, and patternation of any cell, tissue or organ or groups of cells, tissues or organs of a plant, including the root, stem, leaf, shoot, petiole, trichome, flower, petal, stigma, style, stamen, pollen, ovule, seed, embryo, endosperm, seed coat, aleurone, fibre, fruit, cambium, wood, heartwood, parenchyma, aerenchyma, sieve element, phloem or vascular tissue, amongst others.

"Plant biochemistry" or the term "plant biochemical characteristic" or similar term will, when used herein, be understood by those skilled in the art to refer to the metabolic and

catalytic processes of a plant, including primary and secondary metabolism and the products thereof, including any small molecules, macromolecules or chemical compounds, such as but not limited to starches, sugars, proteins, peptides, enzymes, hormones, growth factors, nucleic acid molecules, celluloses, hemicelluloses, calloses, 5 lectins, fibres, pigments such as anthocyanins, vitamins, minerals, micronutrients, or macronutrients, that are produced by plants.

"Plant physiology" or the term "plant physiological characteristic" or similar term will, when used herein, be understood to refer to the functional processes of a plant, including developmental processes such as growth, expansion and differentiation, 10 sexual development, sexual reproduction, seed set, seed development, grain filling, asexual reproduction, cell division, dormancy, germination, light adaptation, photosynthesis, leaf expansion, fibre production, secondary growth or wood production, amongst others; responses of a plant to externally-applied factors such as metals, chemicals, hormones, growth factors, environment and environmental stress factors (eg. 15 anoxia, hypoxia, high temperature, low temperature, dehydration, light, daylength, flooding, salt, heavy metals, amongst others), including adaptive responses of plants to said externally-applied factors.

Means for introducing recombinant DNA into plant tissue or cells include, but are not limited to, transformation using CaCl_2 and variations thereof, in particular the method 20 described by Hanahan (1983), direct DNA uptake into protoplasts (Krens *et al.*, 1982; Paszkowski *et al.*, 1984), PEG-mediated uptake to protoplasts (Armstrong *et al.*, 1990) microparticle bombardment, electroporation (Fromm *et al.*, 1985), microinjection of DNA (Crossway *et al.*, 1986), microparticle bombardment of tissue explants or cells (Christou *et al.*, 1988; Sanford, 1988), vacuum-infiltration of tissue with nucleic acid, or in the case 25 of plants, T-DNA-mediated transfer from *Agrobacterium* to the plant tissue as described essentially by An *et al.* (1985), Dodds *et al.*, (1985), Herrera-Estrella *et al.* (1983a, 1983b, 1985). Methods for transformation of monocotyledonous plants are well known in the art and include *Agrobacterium*-mediated transformation (Cheng *et al.*, 1997 - WO9748814; Hansen 1998 - WO9854961; Hiei *et al.*, 1994 - WO9400977; Hiei *et al.*, 1998 - 30 WO9817813; Rikiishi *et al.*, 1999 - WO9904618; Saito *et al.*, 1995 - WO9506722), microprojectile bombardment (Adams *et al.*, 1999 - US5969213; Bowen *et al.*, 1998 - US5736369; Chang *et al.*, 1994 - WO9413822; Lundquist *et al.*, 1999 - US5874265/US5990390; Vasil and Vasil, 1995 - US5405765. Walker *et al.*, 1999 -

US5955362), DNA uptake (Eyal *et al.*, 1993 - WO9318168), microinjection of *Agrobacterium* cells (von Holt, 1994 - DE4309203) and sonication (Finer *et al.*, 1997 - US5693512).

For microparticle bombardment of cells, a microparticle is propelled into a cell to produce a transformed cell. Any suitable ballistic cell transformation methodology and apparatus can be used in performing the present invention. Exemplary apparatus and procedures are disclosed by Stomp *et al.* (U.S. Patent No. 5,122,466) and Sanford and Wolf (U.S. Patent No. 4,945,050). When using ballistic transformation procedures, the gene construct may incorporate a plasmid capable of replicating in the cell to be transformed.

Examples of microparticles suitable for use in such systems include 1 to 5 μm gold spheres. The DNA construct may be deposited on the microparticle by any suitable technique, such as by precipitation.

A whole plant may be regenerated from the transformed or transfected cell, in accordance with procedures well known in the art. Plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a gene construct of the present invention and a whole plant regenerated therefrom. The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (e.g., apical meristem, axillary buds, and root meristems), and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem).

The term "organogenesis", as used herein, means a process by which shoots and roots are developed sequentially from meristematic centres.

The term "embryogenesis", as used herein, means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether from somatic cells or gametes.

Preferably, the plant is produced according to the inventive method is transfected or transformed with a genetic sequence, or amenable to the introduction of a protein, by any art-recognized means, such as microprojectile bombardment, microinjection, *Agrobacterium*-mediated transformation (including *in planta* transformation), protoplast fusion, or electroporation, amongst others. Most preferably said plant is produced by *Agrobacterium*-mediated transformation.

Agrobacterium-mediated transformation or agrolistic transformation of plants, yeast, moulds or filamentous fungi is based on the transfer of part of the transformation vector sequences, called the T-DNA, to the nucleus and on integration of said T-DNA in the genome of said eukaryote.

5 With "*Agrobacterium*" is meant a member of the *Agrobacteriaceae*, more preferably *Agrobacterium* or *Rhizobacterium* and most preferably *Agrobacterium tumefaciens*.

With "T-DNA", or transferred DNA, is meant that part of the transformation vector flanked by T-DNA borders which is, after activation of the *Agrobacterium vir* genes, nicked at the T-DNA borders and is transferred as a single stranded DNA to the nucleus of an
10 eukaryotic cell.

When used herein, with "T-DNA borders", "T-DNA border region", or "border region" are meant either right T-DNA border (RB) or left T-DNA border (LB). Such a border comprises a core sequence flanked by a border inner region as part of the T-DNA flanking the border and/or a border outer region as part of the vector backbone flanking
15 the border. The core sequences comprise 22 bp in case of octopine-type vectors and 25 bp in case of nopaline-type vectors. The core sequences in the right border region and left border region form imperfect repeats. Border core sequences are indispensable for recognition and processing by the *Agrobacterium* nicking complex consisting of at least VirD1 and VirD2. Core sequences flanking a T-DNA are sufficient to promote transfer of
20 said T-DNA. However, efficiency of transformation using transformation vectors carrying said T-DNA solely flanked by said core sequences is low. Border inner and outer regions are known to modulate efficiency of T-DNA transfer (Wang et al. 1987). One element enhancing T-DNA transfer has been characterized and resides in the right border outer region and is called *overdrive* (Peralta et al. 1986, van Haaren et al. 1987).

25 With "T-DNA transformation vector" or "T-DNA vector" is meant any vector encompassing a T-DNA sequence flanked by a right and left T-DNA border consisting of at least the right and left border core sequences, respectively, and used for transformation of any eukaryotic cell.

With "T-DNA vector backbone sequence" or "T-DNA vector backbone sequences" is
30 meant all DNA of a T-DNA containing vector that lies outside of the T-DNA borders and, more specifically, outside the nicking sites of the border core imperfect repeats.

The current invention includes optimized T-DNA vectors such that vector backbone integration in the genome of a eukaryotic cell is minimized or absent. With "optimized T-

DNA vector" is meant a T-DNA vector designed either to decrease or abolish transfer of vector backbone sequences to the genome of a eukaryotic cell. Such T-DNA vectors are known to the one familiar with the art and include those described by Hanson et al. (1999) and by Stuiver et al. (1999 - WO9901563).

- 5 The current invention clearly considers the inclusion of a DNA sequence encoding a cytokinin oxidase, homologue, derivative or immunologically active and/or functional fragment thereof as defined supra, in any T-DNA vector comprising binary transformation vectors, super-binary transformation vectors, co-integrate transformation vectors, Ri-derived transformation vectors as well as in T-DNA carrying vectors used in
- 10 agrolistic transformation. Preferably, said cytokinin oxidase is a plant cytokinin oxidase, more specifically an *Arabidopsis thaliana* (At)CKX.

With "binary transformation vector" is meant a T-DNA transformation vector comprising:

- (a) a T-DNA region comprising at least one gene of interest and/or at least one selectable marker active in the eukaryotic cell to be transformed; and
- 15 (b) a vector backbone region comprising at least origins of replication active in *E. coli* and *Agrobacterium* and markers for selection in *E. coli* and *Agrobacterium*.

The T-DNA borders of a binary transformation vector can be derived from octopine-type or nopaline-type Ti plasmids or from both. The T-DNA of a binary vector is only transferred to a eukaryotic cell in conjunction with a helper plasmid.

- 20 With "helper plasmid" is meant a plasmid that is stably maintained in *Agrobacterium* and is at least carrying the set of *vir* genes necessary for enabling transfer of the T-DNA. Said set of *vir* genes can be derived from either octopine-type or nopaline-type Ti plasmids or from both.

- 25 With "super-binary transformation vector" is meant a binary transformation vector additionally carrying in the vector backbone region a *vir* region of the Ti plasmid pTiBo542 of the super-virulent *A. tumefaciens* strain A281 (EP0604662, EP0687730). Super-binary transformation vectors are used in conjunction with a helper plasmid.

With "co-integrate transformation vector" is meant a T-DNA vector at least comprising:

- 30 (a) a T-DNA region comprising at least one gene of interest and/or at least one selectable marker active in plants; and
- (b) a vector backbone region comprising at least origins of replication active in *Escherichia coli* and *Agrobacterium*, and markers for selection in *E. coli* and

Agrobacterium, and a set of *vir* genes necessary for enabling transfer of the T-DNA.

The T-DNA borders and said set of *vir* genes of a said T-DNA vector can be derived from either octopine-type or nopaline-type Ti plasmids or from both.

5 With "Ri-derived plant transformation vector" is meant a binary transformation vector in which the T-DNA borders are derived from a Ti plasmid and said binary transformation vector being used in conjunction with a 'helper' Ri-plasmid carrying the necessary set of *vir* genes.

As used herein, the term "selectable marker gene" or "selectable marker" or "marker for selection" includes any gene which confers a phenotype on a cell in which it is expressed to facilitate the identification and/or selection of cells which are transfected or transformed with a gene construct of the invention or a derivative thereof. Suitable selectable marker genes contemplated herein include the ampicillin resistance (Amp^r), tetracycline resistance gene (Tc^r), bacterial kanamycin resistance gene (Kan^r),
10 phosphinothricin resistance gene, neomycin phosphotransferase gene (*npII*), hygromycin resistance gene, β -glucuronidase (GUS) gene, chloramphenicol acetyltransferase (CAT) gene, green fluorescent protein (*gfp*) gene (Haseloff *et al*, 1997), and luciferase gene, amongst others.

With "agrolistics", "agrolistic transformation" or "agrolistic transfer" is meant here a transformation method combining features of *Agrobacterium*-mediated transformation and of biolistic DNA delivery. As such, a T-DNA containing target plasmid is co-delivered with DNA/RNA enabling in planta production of VirD1 and VirD2 with or without VirE2 (Hansen and Chilton 1996; Hansen et al. 1997; Hansen and Chilton 1997 - WO9712046).

With "foreign DNA" is meant any DNA sequence that is introduced in the host's genome
25 by recombinant techniques. Said foreign DNA includes e.g. a T-DNA sequence or a part thereof such as the T-DNA sequence comprising the selectable marker in an expressible format. Foreign DNA furthermore include intervening DNA sequences as defined supra.

With "recombination event" is meant either a site-specific recombination event or a recombination event effected by transposon 'jumping'.

30 With "recombinase" is meant either a site-specific recombinase or a transposase.

With "recombination site" is meant either site-specific recombination sites or transposon border sequences.

With "site specific recombination event" is meant an event catalyzed by a system generally consisting of three elements: a pair of DNA sequences (the site-specific recombination sequences or sites) and a specific enzyme (the site-specific recombinase). The site-specific recombinase catalyzes a recombination reaction only
5 between two site-specific recombination sequences depending on the orientation of the site-specific recombination sequences. Sequences intervening between two site-specific recombination sites will be inverted in the presence of the site-specific recombinase when the site-specific recombination sequences are oriented in opposite directions relative to one another (i.e. inverted repeats). If the site-specific recombination
10 sequences are oriented in the same direction relative to one another (i.e. direct repeats), then any intervening sequences will be deleted upon interaction with the site-specific recombinase. Thus, if the site-specific recombination sequences are present as direct repeats at both ends of a foreign DNA sequence integrated into a eukaryotic genome, such integration of said sequences can subsequently be reversed by interaction of the
15 site-specific recombination sequences with the corresponding site specific recombinase.

A number of different site specific recombinase systems can be used including but not limited to the Cre/lox system of bacteriophage P1, the FLP/FRT system of yeast, the Gin recombinase of phage Mu, the Pin recombinase of *E. coli*, the PinB, PinD and PinF from *Shigella*, and the R/RS system of the pSR1 plasmid. Recombinases generally are
20 integrases, resolvases or flippases. Also dual-specific recombinases can be used in conjunction with direct or indirect repeats of two different site-specific recombination sites corresponding to the dual-specific recombinase (WO99/25840). The two preferred site-specific recombinase systems are the bacteriophage P1 Cre/lox and the yeast FLP/FRT systems. In these systems a recombinase (Cre or FLP) interact specifically
25 with its respective site-specific recombination sequence (lox or FRT respectively) to invert or excise the intervening sequences. The site-specific recombination sequences for each of these two systems are relatively short (34 bp for lox and 47 bp for FRT). Some of these systems have already been used with high efficiency in plants such as tobacco (Dale et al. 1990) and *Arabidopsis* (Osborne et al. 1995). Site-specific
30 recombination systems have many applications in plant molecular biology including methods for control of homologous recombination (e.g. US5527695), for targeted insertion, gene stacking, etc. (WO99/25821) and for resolution of complex T-DNA integration patterns or for excision of a selectable marker (WO99/23202).

Although the site-specific recombination sequences must be linked to the ends of the DNA to be excised or to be inverted, the gene encoding the site specific recombinase may be located elsewhere. For example, the recombinase gene could already be present in the eukaryote's DNA or could be supplied by a later introduced DNA fragment either introduced directly into cells, through crossing or through cross-pollination. Alternatively, a substantially purified recombinase protein could be introduced directly into the eukaryotic cell, e.g. by micro-injection or particle bombardment. Typically, the site-specific recombinase coding region will be operably linked to regulatory sequences enabling expression of the site-specific recombinase in the eukaryotic cell.

With "recombination event effected by transposon jumping" or "transposase-mediated recombination" is meant a recombination event catalyzed by a system consisting of three elements: a pair of DNA sequences (the transposon border sequences) and a specific enzyme (the transposase). The transposase catalyzes a recombination reaction only between two transposon border sequences which are arranged as inverted repeats.

A number of different transposon/transposase systems can be used including but not limited to the Ds/Ac system, the Spm system and the Mu system. These systems originate from corn but it has been shown that at least the Ds/Ac and the Spm system also function in other plants (Fedoroff et al. 1993, Schlappi et al. 1993, Van Sluys et al. 1987). Preferred are the Ds- and the Spm-type transposons which are delineated by 11 bp- and 13 bp- border sequences, respectively.

Although the transposon border sequences must be linked to the ends of the DNA to be excised, the gene encoding the transposase may be located elsewhere. For example, the recombinase gene could already be present in the eukaryote's DNA or could be supplied by a later introduced DNA fragment either introduced directly into cells, through crossing or through cross-pollination. Alternatively, a substantially purified transposase protein could be introduced directly into cells, e.g. by microinjection or by particle bombardment.

As part of the current invention, transposon border sequences are included in a foreign DNA sequence such that they lie outside said DNA sequence and transform said DNA into a transposon-like entity that can move by the action of a transposase.

As transposons often reintegrate at another locus of the host's genome, segregation of the progeny of the hosts in which the transposase was allowed to act might be

necessary to separate transformed hosts containing e.g. only the transposon footprint and transformed hosts still containing the foreign DNA.

In performing the present invention, the genetic element is preferably induced to mobilize, such as, for example, by the expression of a recombinase protein in the cell
5 which contacts the integration site of the genetic element and facilitates a recombination event therein, excising the genetic element completely, or alternatively, leaving a "footprint", generally of about 20 nucleotides in length or greater, at the original integration site. Those hosts and host parts that have been produced according to the inventive method can be identified by standard nucleic acid hybridization and/or
10 amplification techniques to detect the presence of the mobilizable genetic element or a gene construct comprising the same. Alternatively, in the case of transformed host cells, tissues, and hosts wherein the mobilizable genetic element has been excised, it is possible to detect a footprint in the genome of the host which has been left following the excision event, using such techniques. As used herein, the term "footprint" shall be taken
15 to refer to any derivative of a mobilizable genetic element or gene construct comprising the same as described herein which is produced by excision, deletion or other removal of the mobilizable genetic element from the genome of a cell transformed previously with said gene construct. A footprint generally comprises at least a single copy of the recombination loci or transposon used to promote excision. However, a footprint may
20 comprise additional sequences derived from the gene construct, for example nucleotide sequences derived from the left border sequence, right border sequence, origin of replication, recombinase-encoding or transposase-encoding sequence if used, or other vector-derived nucleotide sequences. Accordingly, a footprint is identifiable according to the nucleotide sequence of the recombination locus or transposon of the gene construct
25 used, such as, for example, a sequence of nucleotides corresponding or complementary to a *lox* site or *frt* site.

The term "cell cycle" means the cyclic biochemical and structural events associated with growth and with division of cells, and in particular with the regulation of the replication of DNA and mitosis. Cell cycle includes phases called: G0, Gap1 (G1), DNA synthesis (S),
30 Gap2 (G2), and mitosis (M). Normally these four phases occur sequentially, however, the cell cycle also includes modified cycles wherein one or more phases are absent resulting in modified cell cycle such as endomitosis, acytokinesis, polyploidy, polyteny, and endoreduplication.

The term "cell cycle progression" refers to the process of passing through the different cell cycle phases. The term "cell cycle progression rate" accordingly refers to the speed at which said cell cycle phases are run through or the time spans required to complete said cell cycle phases.

- 5 With "two-hybrid assay" is meant an assay that is based on the observation that many eukaryotic transcription factors comprise two domains, a DNA-binding domain (DB) and an activation domain (AD) which, when physically separated (i.e. disruption of the covalent linkage) do not effectuate target gene expression. Two proteins able to interact physically with one of said proteins fused to DB and the other of said proteins fused to
10 AD will re-unite the DB and AD domains of the transcription factor resulting in target gene expression. The target gene in the yeast two-hybrid assay is usually a reporter gene such as the β -galactosidase gene. Interaction between protein partners in the yeast two-hybrid assay can thus be quantified by measuring the activity of the reporter gene product (Bartel and Fields, 1997). Alternatively, a mammalian two-hybrid system
15 can be used which includes e.g. a chimeric green fluorescent protein encoding reporter gene (Shioda *et al.*, 2000).

- Furthermore, folding simulations and computer redesign of structural motifs of the protein of the invention can be performed using appropriate computer programs (Olszewski, *Proteins* 25 (1996), 286-299; Hoffman, *Comput. Appl. Biosci.* 1 (1995), 675-
20 679). Computer modeling of protein folding can be used for the conformational and energetic analysis of detailed peptide and protein models (Monge, *J. Mol. Biol.* 247 (1995), 995-1012; Renouf, *Adv. Exp. Med. Biol.* 376 (1995), 37-45). In particular, the appropriate programs can be used for the identification of interactive sites of the cytokinin oxidases, its ligands or other interacting proteins by computer assistant
25 searches for complementary peptide sequences (Fassina, *Immunomethods* 5 (1994), 114-120). Further appropriate computer systems for the design of protein and peptides are described in the prior art, for example in Berry, *Biochem. Soc. Trans.* 22 (1994), 1033-1036; Wodak, *Ann. N. Y. Acad. Sci.* 501 (1987), 1-13; Pabo, *Biochemistry* 25 (1986), 5987-5991. The results obtained from the above-described computer analysis
30 can be used for, e.g. the preparation of peptidomimetics of the protein of the invention or fragments thereof. Such pseudopeptide analogues of the natural amino acid sequence of the protein may very efficiently mimic the parent protein (Benkirane, *J. Biol. Chem.* 271 (1996), 33218-33224). For example, incorporation of easily available achiral Ω -

amino acid residues into a protein of the invention or a fragment thereof results in the substitution of amino bonds by polymethylene units of an aliphatic chain, thereby providing a convenient strategy for constructing a peptidomimetic (Banerjee, Biopolymers 39 (1996), 769-777). Superactive peptidomimetic analogues of small peptide hormones in other systems are described in the prior art (Zhang, Biochem. Biophys. Res. Commun. 224 (1996), 327-331). Appropriate peptidomimetics of the protein of the present invention can also be identified by the synthesis of peptidomimetic combinatorial libraries through successive amine alkylation and testing the resulting compounds, e.g., for their binding, kinase inhibitory and/or immunological properties.

Methods for the generation and use of peptidomimetic combinatorial libraries are described in the prior art, for example in Ostresh, Methods in Enzymology 267 (1996), 220-234 and Dörner, Bioorg. Med. Chem. 4 (1996), 709-715.

Furthermore, a three-dimensional and/or crystallographic structure of the protein of the invention can be used for the design of peptidomimetic inhibitors of the biological activity of the protein of the invention (Rose, Biochemistry 35 (1996), 12933-12944; Ruterber, Bioorg. Med. Chem. 4 (1996), 1545-1558).

The compounds to be obtained or identified in the methods of the invention can be compounds that are able to bind to any of the nucleic acids, peptides or proteins of the invention. Other interesting compounds to be identified are compounds that modulate the expression of the genes or the proteins of the invention in such a way that either the expression of said gene or protein is enhanced or decreased by the action of said compound. Alternatively the compound can exert his action by enhancing or decreasing the activity of any of the proteins of the invention. Herein, preferred proteins are novel cytokinin oxidases.

Said compound or plurality of compounds may be comprised in, for example, samples, e.g., cell extracts from, e.g., plants, animals or microorganisms. Furthermore, said compound(s) may be known in the art but hitherto not known to be capable of suppressing or activating cytokinin oxidase interacting proteins. The reaction mixture may be a cell free extract of may comprise a cell or tissue culture. Suitable set ups for the method of the invention are known to the person skilled in the art and are, for example, generally described in Alberts et al., Molecular Biology of the Cell, third edition (1994), in particular Chapter 17. The plurality of compounds may be, e.g., added to the reaction mixture, culture medium or injected into the cell.

If a sample containing a compound or a plurality of compounds is identified in the method of the invention, then it is either possible to isolate the compound from the original sample identified as containing the compound capable of acting as an agonist, or one can further subdivide the original sample, for example, if it consists of a plurality of different compounds, so as to reduce the number of different substances per sample and repeat the method with the subdivisions of the original sample. Depending on the complexity of the samples, the steps described above can be performed several times, preferably until the sample identified according to the method of the invention only comprises a limited number of or only one substance(s). Preferably said sample comprises substances or similar chemical and/or physical properties, and most preferably said substances are identical. Preferably, the compound identified according to the above-described method or its derivative is further formulated in a form suitable for the application in plant breeding or plant cell and tissue culture.

The term "early vigor" refers to the ability of a plant to grow rapidly during early development, and relates to the successful establishment, after germination, of a well-developed root system and a well-developed photosynthetic apparatus.

The term "resistance to lodging" or "standability" refers to the ability of a plant to fix itself to the soil. For plants with an erect or semi-erect growth habit this term also refers to the ability to maintain an upright position under adverse (environmental) conditions. This trait relates to the size, depth and morphology of the root system.

The term 'grafting' as used herein, refers to the joining together of the parts of two different plants so that they bind together and the sap can flow, thus forming a single new plant that can grow and develop. A graft therefore consists of two parts: (i) the lower part is the rootstock as referred to herein and essentially consists of the root system and a portion of the stem, and (ii) the upper part, the scion or graft, which gives rise to the aerial parts of the plant.

As used herein, tblastn refers to an alignment tool that is part of the BLAST (Basic Local Alignment Search Tool) family of programs (<http://www.ncbi.nlm.nih.gov/BLAST/>). BLAST aims to identify regions of optimal local alignment, i.e. the alignment of some portion of two nucleic acid or protein sequences, to detect relationships among sequences which share only isolated regions of similarity (Altschul et al., 1990). In the present invention, tblastn of the BLAST 2.0 suite of programs was used to compare the maize cytokinin oxidase protein sequence against a nucleotide sequence database

dynamically translated in all reading frames (Altschul et al., Nucleic Acids Res. 25: 3389-3402 (1997)).

The following examples and figures are given by means of illustration of the present invention and are in no way limiting. The contents of all references included in

5 this application are incorporated by reference.

BRIEF DESCRIPTION OF THE FIGURES**Figure 1. Schematic representation of plant cytokinin oxidase genes.**

Shown are the structures of different cytokinin oxidase genes isolated from maize
(*ZmCKX1*, accession number AF044603, Biochem. Biophys. Res. Com. 255:328-333,
1999) and Arabidopsis (*AtCKX1* to *AtCKX4*). Exons are denominated with 'E' and
represented by shaded boxes. Introns are represented by white boxes. Further indicated
are the gene sizes (in kb, on top of each structure), the gene accession numbers (under
the names) and a size bar representing 0.5 kb.

Figure 2. Alignment of plant cytokinin oxidase amino acid sequences.

The amino acid sequences from cytokinin oxidases from maize (*ZmCKX1*) and
Arabidopsis (*AtCKX1* to *AtCKX4*) are aligned. Identical amino acid residues are marked
by a black box, similar amino acid residues are in a grey box. Amino acid similarity
groups: (M,I,L,V), (F,W,Y), (G,A), (S,T), (R,K,H), (E,D), (N,Q),

Figure 3. Northern blot analysis of *AtCKX1*-expressing tobacco and *Arabidopsis* plants.

(A) Northern blot analysis of constitutively expressing tobacco plants (lanes 1-8)
compared to wild type SNN tobacco (lane 9)

(B) Comparison of tetracycline-induced gene expression in leaves after 12h of induction
with a constitutively expressing clone. Lanes 2-9, leaves of four different *AtCKX1*-
W38TetR clones (+, -, with or without tetracycline treatment), lane 1, constitutively
expressing 35S:: *AtCKX1* clone.

(C) Northern blot analysis of *Arabidopsis* plants constitutively expressing *AtCKX1* gene.
Lanes 2-4, three different constitutively expressing 35S:: *AtCKX1* clones compared to
wild type Arabidopsis plant (lane 1).

Figure 4: Growth characteristics of 35S::*AtCKX1* transgenic *Arabidopsis* plants.

(A) Two wild type seedlings (left) compared to two 35S:: *AtCKX1* expressing seedlings
(right). Note the increased formation of adventitious roots and increased root branching
in the transgenic seedlings. Pictures were taken 14 days after germination. Plants were
grown *in vitro* on MS medium in petri dishes in a vertical position.

(B) Like A, but roots stained with toluidine blue.

(C) Top view of a petridish with *35S::AtCKX1* transgenic seedlings three weeks after germination.

(D) A *35S::AtCKX1* transgenic plants grown in liquid culture. Roots of wild type seedlings
5 grow poorly under these conditions (not shown).

(E) Transformants (T0) that express the *35S::AtCKX1* gene (three plants on the right), a wild type plant is shown on the left.

(F) Phenotype of T1 plants grown in soil. Wild type plant (left) compared to two *35S::AtCKX1* transgenic plants.

10

Figure 5: Phenotype of *AtCKX2* overexpressing Arabidopsis plants.

T1 generation of *35S::AtCKX2* expressing Arabidopsis plants (two plants on the right) compared to wild type (plant on the left).

15 **Figure 6. Northern blot analysis of *AtCKX2*-expressing tobacco and Arabidopsis plants.**

(A) Northern blot analysis of constitutively expressing tobacco plants (lanes 1-7) compared to wild type SNN tobacco (lane 8)

(B) Northern blot analysis of Arabidopsis plants constitutively expressing *AtCKX2* gene.

20 Lanes 2-8, seven different constitutively expressing *35S::AtCKX2* clones compared to wild type Arabidopsis plant (lane 1).

Figure 7. Shoot phenotype of *AtCKX1* and *AtCKX2* expressing tobacco plants.

(A) Top view of six week old plants.

25 (B) Tobacco plants at the flowering stage.

(C) Kinetics of stem elongation. Arrows mark the onset of flowering. Age of plants (days after germination) and leaf number at that stage are indicated above the arrows. Bars indicate SD; $n = 12$.

(D) Number of leaves ($n = 12$) formed between day 68 and day 100 after germination
30 and final surface area of these leaves (100% of wild type is $3646 \pm 144 \text{ cm}^2$; $n = 3$).

(E) Comparison of leaf size and senescence. Leaves were from nodes number 4, 9, 12, 16 and 20 from the top (from left to right).

Figure 8. Root phenotype of *AtCKX* expressing transgenic tobacco plants.

- (A) Seedlings 17 days after germination.
- (B) Root system of soil grown plants at the flowering stage.
- (C) Root length, number of lateral roots (LR) and adventitious roots (AR) on day 10 after germination.
- (D) Dose-response curve of root growth inhibition by exogenous cytokinin. Bars indicate \pm SD; $n = 30$.

Figure 9: Growth of axillary shoot meristems in *35S::AtCKX1* expressing tobacco plants.**Figure 10: Histology of shoot meristems, leaves and root meristems of *AtCKX1* overexpressing tobacco plants versus wild type (WT) tobacco.**

- (A) Longitudinal median section through the vegetative shoot apical meristem. P, leaf primordia.
- (B) Vascular tissue in second order veins of leaves. X, xylem, PH, a phloem bundle.
- (C) Cross sections of fully developed leaves.
- (D) Scanning electron microscopy of the upper leaf epidermis.
- (E) Root apices stained with DAPI. RM, root meristem.
- (F) Longitudinal median sections of root meristems ten days after germination. RC, root cap; PM, promeristem.
- (G) Transverse root sections 10 mm from the apex. E, epidermis, C1-C4, cortical cell layer, X, xylem, PH, phloem. Bars are 100 μ m.

Figure 11: Northern blot analysis of *AtCKX3* and *AtCKX4*-expressing tobacco plants.

- (A) Northern blot analysis of constitutively expressing *AtCKX3* tobacco plants. Lane designations indicate individual transgenic plant numbers, WT is wild type SNN tobacco. The blot on top was probed with a *AtCKX3* specific probe, the lower blot with a probe specific for the 25S rRNA and serves as a control for RNA loading.

(B) Northern blot analysis of constitutively expressing *AtCKX4* tobacco plants. Lane designations indicate individual transgenic plant numbers, WT is wild type SNN tobacco. The blot on top was probed with an *AtCKX4* specific probe, the lower blot with a probe specific for the 25S rRNA and serves as a control for RNA loading.

5 **Figure 12: Recipocal grafts of *AtCKX2* transgenic tobacco plants and wild type plants.**

(A) Two plants on the left: Control (WT scion grafted on a WT rootstock).

Two plants on the right: WT scion grafted on a *AtCKX2-38* transgenic rootstock.

(B) Left: Control (WT scion grafted on a WT rootstock).

10 Right: Scion of *AtCKX2-38* plant grafted on WT rootstock.

(C) Magnification of root area.

Left: Control (WT scion grafted on a WT rootstock).

Right: WT scion grafted on an *AtCKX2-38* transgenic rootstock.

(D) Formation of adventitious roots.

15 Left: Control (WT scion grafted on an WT rootstock).

Right: WT scion grafted on an *AtCKX2-38* transgenic rootstock.

EXAMPLES**Example 1. Brief description of the sequences of the invention**

Seq ID No	Description
1	<i>AtCKX1</i> genomic
2	<i>AtCKX1</i> protein
3	<i>AtCKX2</i> genomic
4	<i>AtCKX2</i> protein
5	<i>AtCKX3</i> genomic
6	<i>AtCKX3</i> protein
7	<i>AtCKX4</i> genomic
8	<i>AtCKX4</i> protein
9	<i>AtCKX5</i> genomic (short version)
10	<i>AtCKX5</i> protein (short version)
11	<i>AtCKX6</i> genomic
12	<i>AtCKX6</i> protein
13	5'primer <i>AtCKX1</i>
14	3'primer <i>AtCKX1</i>
15	5'primer <i>AtCKX2</i>
16	3'primer <i>AtCKX2</i>
17	5'primer <i>AtCKX3</i>
18	3'primer <i>AtCKX3</i>
19	5'primer <i>AtCKX4</i>
20	3'primer <i>AtCKX4</i>
21	5'primer <i>AtCKX5</i>
22	3'primer <i>AtCKX5</i>
23	5'primer <i>AtCKX6</i>
24	3'primer <i>AtCKX6</i>
25	<i>AtCKX1</i> cDNA
26	<i>AtCKX2</i> cDNA
27	<i>AtCKX3</i> cDNA
28	<i>AtCKX4</i> cDNA
29	<i>AtCKX5</i> cDNA (short version)
30	<i>AtCKX6</i> cDNA
31	<i>AtCKX2</i> cDNA fragment
32	<i>AtCKX2</i> peptide fragment
33	<i>AtCKX5</i> genomic (long version)
34	<i>AtCKX5</i> cDNA (long version)
35	<i>AtCKX5</i> protein (long version)
36	root clavata homolog promoter

Example 2. Identification of candidate cytokinin oxidase encoding genes from *Arabidopsis thaliana*

Six different genes were identified from *Arabidopsis thaliana* that bear sequence similarity to a cytokinin oxidase gene from maize (Morris *et al.*, Biochem Biophys Res Comm 255:328-333, 1999; Houda-Herlin *et al.* Plant J 17:615-626; WO 99/06571). These genes were found by screening 6-frame translations of nucleotide sequences from public genomic databases with the maize protein sequence, employing tblastn program. These sequences were designated as *Arabidopsis thaliana* cytokinin oxidase-like genes or *AtCKX*. They were arbitrarily numbered as *AtCKX1* to *AtCKX6*. The below list summarizes the information on these genes. The predicted ORF borders and protein sequences are indicative, in order to illustrate by approximation the protein sequence divergence between the *Arabidopsis* and maize cytokinin oxidases, as well as amongst the different *Arabidopsis* cytokinin oxidases. The ORF borders and protein sequences shown should not be taken as conclusive evidence for the mode of action of these *AtCKX* genes. For DNA and protein sequence comparisons the program MegAlign from DNASTar was used. This program uses the Clustal method for alignments. For multiple alignments of protein and cDNA sequences the gap penalty and gap length penalty was set at 10 each. For pairwise alignments of proteins the parameters were as follows: Ktuple at 1; Gap penalty at 3; window at 5; diagonals saved at 5. For pairwise alignments of cDNA's the parameters were as follows: Ktuple at 2; Gap penalty at 5; window at 4; diagonals saved at 4. The similarity groups for protein alignments was: (M,I,L,V), (F,W,Y), (G,A), (S,T), (R,K,H), (E,D), (N,Q). The values that are indicated amongst the *Arabidopsis* cDNA and protein sequences represent the lowest and highest values found with all combinations.

A. Gene name: *AtCKX1* (*Arabidopsis thaliana* cytokinin oxidase-like protein 1, SEQ ID NO1)

Location in database (accession number, location on bac): AC002510, *Arabidopsis thaliana* chromosome II section 225 of 255 of the complete sequence. Sequence from clones T32G6.

ORF predicted in the database:

15517..16183, 16415..16542, 16631..16891, 16995..17257, 17344..17752

The *AtCKX1* cDNA sequence is listed as SEQ ID NO 25

Predicted protein sequence: SEQ ID NO 2

5 **Homologies**

% identity with *Z. mays* cDNA:

31,5% (Dnastar/MegAlign - Clustal method)

% similarity with *Z. mays* protein:

32,2% (Dnastar/MegAlign - Clustal method)

10

% identity with other *Arabidopsis* cDNA's (range):

38,2% (*AtCKX2*) - 54,1% (*AtCKX6*) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

15

37,1% (*AtCKX2*) - 58,1% (*AtCKX6*) (Dnastar/MegAlign - Clustal method)

B. Gene name: *AtCKX2* (*Arabidopsis thaliana* cytokinin oxidase-like protein 2, SEQ ID NO3)

20 **Location in database** (accession number, location on bac): AC005917, *Arabidopsis thaliana* chromosome II section 113 of 255 of the complete sequence. Sequence from clones F27F23, F3P11.

ORF predicted in the database:

25 complement, 40721..41012, 41054..41364, 41513..41770, 42535..42662, 43153..43711

Please note: The cDNA sequence identified by the inventor using the gene prediction program NetPlantGene (<http://www.cbs.dtu.dk/services/NetGene2/>) was different than the one annotated in the database. Based on the new cDNA sequence the ORF predicted in the database was revised:

30

complement, 40721..41012, 41095..41364, 41513..41770, 42535..42662, 43153..43711

The protein sequence encoded by this cDNA is listed as SEQ ID NO 4. The cDNA of *AtCKX2* was cloned by RT-PCR from total RNA of *AtCKX2* transgenic plant tissue with the one-step RT-PCR kit (Qiagen, Hilden, Germany) and sequenced using an ABI PRISM Big Dye Terminator cycle sequencing reaction kit (Perkin Elmer Applied Biosystems Division). This confirmed that the cDNA sequence identified and predicted

35

by the inventor was correct. The new *AtCKX2* cDNA sequence is listed as SEQ ID NO 26. An 84-bp fragment corresponding to nucleotides 1171 through 1254 of the *AtCKX2* cDNA is listed as SEQ ID NO 31. The corresponding peptide sequence of this 84-bp cDNA sequence is listed as SEQ ID NO 32.

5

Homologies

% identity with *Z. mays* cDNA:

38,4% (Dnastar/MegAlign - Clustal method)

10

% similarity with *Z. mays* protein:

37,5% (Dnastar/MegAlign - Clustal method)

% identity with other *Arabidopsis* cDNA's (range):

34,9% (*AtCKX6*) – 64,5% (*AtCKX4*) (Dnastar/MegAlign - Clustal method)

15

% similarity with other *Arabidopsis* proteins (range):

36,5% (*AtCKX6*) – 66,1% (*AtCKX4*) (Dnastar/MegAlign - Clustal method)

20

C. Gene name: *AtCKX3* (*Arabidopsis thaliana* cytokinin oxidase-like protein 3, SEQ ID NO 5)

Location in database (accession number, location on bac): AB024035, *Arabidopsis thaliana* genomic DNA, chromosome 5, P1 clone: MHM17, complete sequence.

25

No prediction of the ORF in the database.

The gene was identified by the inventor using several gene prediction programs including GRAIL (<ftp://arthur.epm.ornl.gov/pub/xgrail>), Genscan (<http://CCR-081.mit.edu/GENSCAN.html>) and NetPlantGene (<http://www.cbs.dtu.dk/services/NetGene2/>):

30

complement, 29415..29718, 29813..30081, 30183..30443, 30529..30656, 32107..32716

The new *AtCKX3* cDNA sequence identified by the inventor is listed as SEQ ID NO 27

Predicted protein sequence, based on own ORF prediction: SEQ ID NO 6

35

Homologies

% identity with *Z. mays* cDNA:

38,7% (Dnastar/MegAlign - Clustal method)

5 % similarity with *Z. mays* protein:

39,2% (Dnastar/MegAlign - Clustal method)

% identity with other *Arabidopsis* cDNA's (range):

38,8% (*AtCKX6*) – 51,0% (*AtCKX2*) (Dnastar/MegAlign - Clustal method)

10

% similarity with other *Arabidopsis* proteins (range):

39,9% (*AtCKX6*) – 46,7% (*AtCKX2*) (Dnastar/MegAlign - Clustal method)

D. Gene name: *AtCKX4* (*Arabidopsis thaliana* cytokinin oxidase-like protein 4, SEQ ID

15 NO 7)

Location in database (accession number, location on bac):

1) AL079344, *Arabidopsis thaliana* DNA chromosome 4, BAC clone T16L4 (ESSA project)

20 2) AL161575, *Arabidopsis thaliana* DNA chromosome 4, contig fragment No. 71.

ORF predicted in the database:

1) 76187..76814, 77189..77316, 77823..78080, 78318..78586, 78677..78968

2) 101002..101629, 102004..102131, 102638..102895, 103133..103401,

25 103492..103783

The *AtCKX4* cDNA sequence is listed as SEQ ID NO 28

Predicted protein sequence: SEQ ID NO 8

30 **Homologies**

% identity with *Z. mays* cDNA:

41,0% (Dnastar/MegAlign - Clustal method)

% similarity with *Z. mays* protein:

35 41,0% (Dnastar/MegAlign - Clustal method)

% identity with other *Arabidopsis* cDNA's (range):

35,2% (*AtCKX6*) – 64,5% (*AtCKX2*) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

5 35,1% (*AtCKX6*) – 66,1% (*AtCKX2*) (Dnastar/MegAlign - Clustal method)

E. Gene name: *AtCKX5* (*Arabidopsis thaliana* cytokinin oxidase-like protein 5, SEQ ID NO 9)

10 **Location in database** (accession number, location on bac): AC023754, F1B16, complete sequence, chromosome 1

No prediction of the ORF in the database.

The gene was identified by the inventors using several gene prediction programs
15 including GRAIL (<ftp://arthur.epm.ornl.gov/pub/xgrail>), Genscan (http://CCR-081.mit.edu/GEN_SCAN.html) and NetPlantGene (<http://www.cbs.dtu.dk/services/NetGene2/>).

43756..44347, 44435..44562, 44700..44966, 45493..45755, 46200..46560

20 The new *AtCKX5* cDNA sequence identified and predicted by the inventor is listed as SEQ ID NO 29. The predicted protein sequence for this cDNA is listed as SEQ ID NO 10. A second potential ATG startcodon is present 9 nucleotides more upstream in the genomic sequence. It is unclear which of these 2 startcodons encodes the first amino acid of the protein. Therefore, a second potential *AtCKX5* cDNA starting at this upstream
25 startcodon is also listed in this invention as SEQ ID NO 34. The corresponding genomic sequence is listed as SEQ ID NO 33 and the encoded protein as SEQ ID NO 35.

Homologies

% identity with *Z. mays* cDNA:

30 39,1% (Dnastar/MegAlign - Clustal method)

% similarity with *Z. mays* protein:

36,6% (Dnastar/MegAlign - Clustal method)

35 % identity with other *Arabidopsis* cDNA's (range):

40,1% (AtCKX2) – 44,0% (AtCKX3) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

41,6% (AtCKX4) – 46,4% (AtCKX6) (Dnastar/MegAlign - Clustal method)

5

F. Gene name: *AtCKX6* (*Arabidopsis thaliana* cytokinin oxidase-like protein 6, SEQ ID NO 11)

10 **Location in database** (accession number, location on bac): AL163818, *Arabidopsis thaliana* DNA chromosome 3, P1 clone MAA21 (ESSA project).

ORF predicted in the database:

46630..47215, 47343..47470, 47591..47806, 47899..48161, 48244..48565

15 The *AtCKX6* cDNA sequence is listed as SEQ ID NO 30

Predicted protein sequence: SEQ ID NO 12

Homologies

20 % identity with *Z. mays* cDNA:

37,3% (Dnastar/MegAlign - Clustal method)

% similarity with *Z. mays* protein:

36,1% (Dnastar/MegAlign - Clustal method)

25

% identity with other *Arabidopsis* cDNA's (range):

34,9% (AtCKX2) – 54,1% (AtCKX1) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

30 35,1% (AtCKX4) – 58,1% (AtCKX1) (Dnastar/MegAlign - Clustal method)

Genes *AtCKX3* and *AtCKX5* were not annotated as putative cytokinin oxidases in the database and ORFs for these genes were not given. Furthermore, the ORF (and consequently the protein structures) predicted for *AtCKX2* was different from our own prediction and our prediction was confirmed by sequencing the *AtCKX2* cDNA.

35

A comparison of the gene structure of the *Arabidopsis AtCKX* genes 1 to 4 and the maize *CKX* gene is shown in Fig 1.

The predicted proteins encoded by the *Arabidopsis AtCKX* genes show between 32% and 41% sequence similarity with the maize protein, while they show between 35% and 66% sequence similarity to each other. Because of this reduced sequence conservation, it is not clear *a priori* whether the *Arabidopsis AtCKX* genes encode proteins with cytokinin oxidase activity. An alignment of the *Arabidopsis AtCKX* predicted proteins 1 to 4 and the maize *CKX* gene is shown in Fig 2.

Example 3. Transgenic plants overexpressing *AtCKX1* showed increased cytokinin oxidase activity and altered plant morphology

1. Description of the cloning process

The following primers were used to PCR amplify the *AtCKX1* gene from *Arabidopsis thaliana*, accession Columbia (non-homologous sequences used for cloning are in lower case):

Sequence of 5' primer: cggtcgacATGGGATTGACCTCATCCTTACG (SEQ ID NO:13)

Sequence of 3' primer: gcgtcgacTTATACAGTTCTAGGTTTCGGCAGTAT (SEQ ID NO: 14)

A 2235-bp PCR fragment, amplified by these primers, was inserted in the *Sal* I site of pUC19. The insert was sequenced and confirmed that the PCR amplification product did not contain any mutations. The *Sall*/*Sall* fragment of this vector was subcloned in the *Sall* site downstream of a modified CaMV 35S promoter (carrying three tetracycline operator sequences) in the binary vector pBinHyg-Tx (Gatz *et al.*, 1992). The resulting construct was introduced into tobacco and *Arabidopsis thaliana* through *Agrobacterium*-mediated transformation, using standard transformation protocols.

2. Molecular analysis of the transgenic lines

Several transgenic lines were identified that synthesize the *AtCKX1* transcript at high levels (Fig 3). Transgenic lines expressing *AtCKX1* transcript also showed increased cytokinin oxidase activity as determined by a standard assay for cytokinin oxidase activity based on conversion of [2-³H]iP to adenine as described (Motyka *et al.*, 1996).

This is exemplified for 2 tobacco and 2 *Arabidopsis* lines in Table 6. This result proves that the *AtCKX1* gene encodes a protein with cytokinin oxidase activity.

Table 6. Cytokinin oxidase activity in *AtCKX1* transgenic plant tissues

Leaf sample		
Plant species	Plant line	Cytokinin oxidase activity (nmol Ade/mg protein.h)
<i>Arabidopsis</i>	Col-0 wild-type	0.009
	CKX1-11	0.024
	CKX1-22	0.026
	CKX1-22	0.027
Tobacco	SNN wild-type	0.004
	CKX1-SNN-8	0.016
	CKX1-SNN-28	0.021

3. Phenotypic description of the transgenic lines

3.1 In tobacco:

The plants had a dwarfed phenotype with reduced apical dominance (Figure 7 A, B and C) and increased root production (Figure 8).

Five categories of phenotype:

- 1) strong - 2 clones
- 2) intermediate - 3 clones
- 3) weak - 4 clones
- 4) tall plants (as WT) with large inflorescence - 5 clones
- 5) similar to WT, 9 clones

Height (see Fig. 7 B and C)

- WT: between 100-150 cm
- weak: approximately 75 cm

- intermediate: appr. 40-45 cm (main stem app. 25 cm but overgrown by side branches.
- strong: appr. 10 cm

- 5 The transgenics *AtCKX1-48* and *AtCKX1-50* displayed a strong phenotype. Below are measurements for stem elongation as compared to WT plants:

Line	Wild-type	AtCKX1-48	AtCKX1-50
Days after germination	Height (cm)	Height (cm)	Height (cm)
47	9,5 ± 0,5	1,3 ± 0,3	1,2 ± 0,2
58	22,4 ± 2,3	2,2 ± 0,3	2,3 ± 0,3
68	35,3 ± 2,6	3,1 ± 0,5	2,6 ± 0,5
100	113,3 ± 9,8	7,1 ± 0,8	4,8 ± 0,9
117	138,6 ± 8,1	8,7 ± 0,7	6,6 ± 0,9
131	139,0 ± 9,3	9,3 ± 0,7	8,6 ± 1,0
152	136,6 ± 10,4	10,9 ± 1,1	10,0 ± 1,0
165		11,8 ± 1,9	11,4 ± 1,4
181		16,5 ± 1,7	14,9 ± 1,2
198		19,5 ± 1,5	18,1 ± 1,3

- 10 Experimental: Plants were grown in soil in a greenhouse. Data were collected from at least ten plants per line.

Leaves (see Figure 7 D and E)

- 15 The shape of leaves of *AtCKX1* transgenic expressors was lanceolate (longer and narrow): the width-to-length ratio of mature leaves was reduced from 1:2 in wild type plants to 1:3 in *AtCKX1* transgenics (Figure 7 E). The number of leaves and leaf surface was reduced compared to WT (see Figure 7 D). A prominent difference was also noted for progression of leaf senescence. In WT tobacco, leaf senescence starts in the most basal leaves and leads to a uniform reduction of leaf pigment (Figure 7 E). By contrast, ageing leaves of strongly expressing *AtCKX1* plants stayed green along the leaf veins

and turned yellow in the intercostal regions, indicating altered leaf senescence. The texture of older leaves was more rigid.

Roots

5 In vitro grown plants highly expressing the gene were easily distinguishable from the WT by their ability to form more roots which are thicker (stronger) (Figure 8 A), as well as by forming aerial roots along the stem.

The primary root was longer and the number of lateral and adventitious roots was higher as illustrated in Figure 8 C for *AtCKX1-50* overexpressing seedlings (see also Example 10 9).

The dose-response curve of root growth inhibition by exogenous cytokinin showed that roots of transgenic seedlings are more cytokinin resistant than WT roots (Figure 8 D). The resistance of *AtCKX1* transgenics to IPR was less marked than for *AtCKX2*, which is consistent with the smaller changes in iP-type cytokinins in the latter (see Table 10).

15 A large increase in root biomass was observed for adult plants grown in soil (see Figure 8 B for a plant grown in soil for 4 to 5 months) despite the fact that growth of the aerial plant parts was highly reduced.

Internode distance

- 20
- intermediate phenotype: the 5th internode below inflorescence is about 2.5 cm long and 9th internode was about 0,5 cm long compared to 5 cm and 2 cm for the length of the 5th and 9th internode respectively, in WT plants.
 - strong phenotype: plant *AtCKX1-50* The length of the 20th internode from the bottom measured at day 131 after germination was $1,3 \pm 0,4$ mm compared to 25 $39,2 \pm 3,8$ mm for WT

Apical dominance and branching

More side branches were formed indicating reduced apical dominance compared to WT plants during vegetative growth (see Figure 9). The side branches overgrew the main 30 stem, reaching a height of 40-45 cm for intermediate *AtCKX1* expressors. Even secondary branches appeared. However, the buds were not completely released from apical dominance, i.e. lateral shoots did not really continue to develop. The reduced

apical dominance might be due to reduced auxin production by the smaller shoot apical meristem (see Example 10).

Reproductive development

The onset of flowering in *AtCKX1* transgenics was delayed, the number of flowers and the seed yield per capsule was reduced. The size of flowers was not altered in transgenic plants and the weight of the individual seeds was comparable to the weight of seeds from wild type plants. Data for two representative *AtCKX1* transgenics is summarized below:

A. Onset of flowering

Line	Wild-type	AtCKX1-48	AtCKX1-50
Flowering time (DAG)	106,2 ± 3,3	193,3 ± 4,3	191,8 ± 3,8

Experimental: Data collected for at least ten plants per line. The full elongation of the first flower was defined as onset of flowering. DAG = days after germination.

B. Number of seed capsules per plant

Line	Wild-type	AtCKX1-48	AtCKX1-50
Number of capsules	83,33 ± 5,13	2,00 ± 1,00	2,60 ± 1,67

Experimental: Number of seed capsules was determined at least from 5 different plants. Please note that these plants were grown under greenhouse conditions during winter time. This affects negatively the number of flowers that are formed, in particular in the transgenic clones. However, the general picture that they form a reduced number of flowers is correct. n.d., not determined

C. Seed yield / capsule (mg)

Line	Wild-type	AtCKX1-48	AtCKX1-50
Seed/capsule (mg)	87,41 ± 28,75	23,83 ± 13,36	61,8 ± 40,66

Experimental: Seed yield was determined for at least 12 seed capsules. The size of seed capsules was very variable, hence the large standard deviations. n.d., not determined

D. Weight of 100 seeds (mg)

Line	Wild-type	AtCKX1-48	AtCKX1-50
Seeds weight (mg)	9,73 ± 0,44	10,70 ± 1,60	9,54 ± 0,94

Experimental: The seed biomass was determined as the weight of 100 seed from at least 5 different seed capsules. n.d., not determined

3.2 In Arabidopsis

- onset of germination was same as for WT
- the total root system was enlarged and the number of side roots and adventitious roots was enhanced (see Figure 4 A through D)
- the growth of aerial organs was reduced resulting in a dwarfed phenotype (see Figure 4 E and F) and the leaf biomass was reduced. Leaf and flower formation is delayed.
- the life cycle was longer compared to WT and the seed yield was lower compared to WT

The following morphometric data illustrate these phenotypes:

Root development

A. Total length of the root system

Line	Wild-type	AtCKX1-11	AtCKX1-15
Length (mm)	32,5	76,5	68,4

5

B. Primary root length

Line	Wild-type	AtCKX1-11	AtCKX1-15
Length (mm)	32,3 ± 3,8	52,3 ± 4,8	39,9 ± 4,2

C. Lateral roots (LR) length

10

Line	Wild-type	AtCKX1-11	AtCKX1-15
Length (mm)	0,2 ± 0,4	15,6 ± 11,0	10,4 ± 7,6

D. Adventitious roots length

Line	Wild-type	AtCKX1-11	AtCKX1-15
Length (mm)	0,03 ± 0,18	8,6 ± 8,5	19,1 ± 11,0

15

E. Number of lateral roots (LR)

Line	Wild-type	AtCKX1-11	AtCKX1-15
Number of LR	0,3 ± 0,5	10,4 ± 5,4	2,6 ± 1,1

F. Number of adventitious roots (AR)

Line	Wild-type	AtCKX1-11	AtCKX1-15
Number of AR	0,03 ± 0,18	1,6 ± 1,1	2,6 ± 1,1

20

Experimental: Measurements were carried out on plants 8 days after germination in vitro on MS medium. At least 17 plants per line were scored.

Shoot development

A. Leaf surface

Line	Wild-type	AtCKX1-11-7 T3 homozygous plants	AtCKX1-11-12 T3 homozygous plants	AtCKX1-15-1 T3 homozygous plants
Leaf surface (cm ²)	21,16 ± 1,73	2,28 ± 0,58	2,62 ± 0,28	1,66 ± 0,22

Experimental: Leaf surface area of main rosette leaves formed after 30 days after germination was measured. 3 plants per clone were analysed.

Reproductive development

Onset of flowering

Line	Wild-type	AtCKX1-11 T3 heterozygous plants	AtCKX2-2 T2 heterozygous plants	AtCKX2-5 T2 heterozygous plants
Flowering time (DAG)	43,6 ± 5,8	69,7 ± 9,4	51,2 ± 4,1	45,1 ± 6,9

Experimental: Plants were grown under greenhouse condition. At least 13 plants per clone were analysed. DAG = days after germination

Conclusion: The analysis of *AtCKX1* transgenic *Arabidopsis* plants confirmed largely the results obtained from tobacco and indicates the general nature of the consequences of a reduced cytokinin content. The total root system was enlarged (the total root length was increased app. 110-140% in *AtCKX1* transgenics), the shoot developed more slowly (retarded flowering) and the leaf biomass was reduced. The seed yield was lower in the transgenics as well (data not shown).

Example 4. Transgenic plants overexpressing AtCKX2 showed increased cytokinin oxidase activity and altered plant morphology

1. Description of the cloning process

- 5 The following primers were used to PCR amplify the AtCKX2 gene from *Arabidopsis thaliana*, accession Columbia (non-homologous sequences used for cloning are in lower case):

Sequence of 5' primer: gcggtaccAGAGAGAGAAACATAAACAATGGC (SEQ ID NO:15)

- 10 Sequence of 3' primer: gcggtaccCAATTTTACTTCCACCAAATGC (SEQ ID NO:16)

A 3104-bp PCR fragment, amplified by these primers, was inserted in the KpnI site of pUC19. The insert was sequenced to check that no differences to the published sequence were introduced by the PCR procedure. The KpnI/KpnI fragment of this vector
15 was subcloned in the KpnI site downstream of a modified CaMV 35S promoter (carrying three tetracycline operator sequences) in the binary vector pBinHyg-Tx (Gatz *et al.*, 1992). The resulting construct was introduced into tobacco and *Arabidopsis thaliana* through *Agrobacterium*-mediated transformation, using standard transformation protocols.

20

2. Molecular analysis of the transgenic lines

Several transgenic lines were identified that synthesize the AtCKX2 transcript at high levels (Fig 6). Transgenic lines expressing AtCKX2 transcript also showed increased cytokinin oxidase activity. This is exemplified for 2 tobacco and 3 *Arabidopsis* lines in
25 Table 7. This result proves that the AtCKX2 gene encodes a protein with cytokinin oxidase activity.

30

Table 7. Cytokinin oxidase activity in *AtCKX2* transgenic plant tissues

Sample		
Plant species and tissue	Plant line	Cytokinin oxidase activity (nmol Ade/mg protein.h)
<i>Arabidopsis</i> callus	Col-0 wild-type	0.037
	CKX2-15	0.351
	CKX2-17	0.380
	CKX2-55	0.265
Tobacco leaves	SNN wild-type	0.009
	CKX2-SNN-18	0.091
	CKX2-SNN-19	0.091

3. Phenotypic description of the transgenic lines

3.1 In tobacco (see Fig 7 to 10):

Three categories of phenotype:

- 1) strong - 15 clones (similar to intermediate phenotype of *AtCKX1*)
- 2) weak - 6 clones
- 3) others - similar to WT plants, 7 clones

Aerial plant parts

The observations concerning plant height, internode distance, branching, leaf form and yellowing were similar as for *AtCKX1* transgenics with some generally minor quantitative differences in that the dwarfing characteristics were more severe in *AtCKX1* transgenics than in *AtCKX2* transgenics (compare *AtCKX1* plants with *AtCKX2* plants in Figure 7 A and B). This is illustrated below for stem elongation and internode distance measurements of clones with a strong phenotype *AtCKX2-38* and *AtCKX2-40*:

Stem elongation

Line	Wild-type	AtCKX2-38	AtCKX2-40
Days after germination	Height (cm)	Height (cm)	Height (cm)
47	9,5 ± 0,5	2,4 ± 0,1	2,6 ± 0,2
58	22,4 ± 2,3	5,5 ± 0,7	5,3 ± 0,5
68	35,3 ± 2,6	7,1 ± 0,8	7,0 ± 0,7
100	113,3 ± 9,8	15,5 ± 2,5	20,3 ± 6,4
117	138,6 ± 8,1	19,8 ± 3,8	29,5 ± 6,0
131	139,0 ± 9,3	26,5 ± 7,0	33,4 ± 5,8
152	136,6 ± 10,4	33,7 ± 6,3	33,9 ± 6,4
165		36,2 ± 4,3	

- 5 Experimental: Plants were grown in soil in a green house. Data were collected from at least ten plants per line.

Internode distance

Line	Wild-type	AtCKX2-38
Internode distance (mm)	39,2 ± 3,8	7,2 ± 1,6

- 10 Experimental: The length of the 20th internode from the bottom was measured at day 131 after germination.

Roots

- 15 In vitro grown plants highly expressing the gene were easily distinguishable from WT plants by their ability to form more roots which are thicker (stronger) as well as by forming aerial roots along the stem.

The primary root was longer and the number of lateral and adventitious roots was higher as illustrated in Figure 8 C for *AtCKX2-38* overexpressing seedlings (see also Example 9).

- 20 The dose-response curve of root growth inhibition by exogenous cytokinin showed that roots of transgenic seedlings were more cytokinin resistant than WT roots (Figure 8 D). The resistance of *AtCKX1-28* transgenics to iPR was less marked than for *AtCKX2-38*, which is consistent with the smaller changes in iP-type cytokinins in the latter (see Table 10).

An increase in fresh and dry weight of the root biomass of T0 lines of AtCKX2 transgenic plants compared to WT was observed for plant grown in soil, as illustrated in the following table:

Line	Wild-type	AtCKX2 (T0)
Fresh weight (g)	45,2 ± 15,4	77,1 ± 21,3
Dry weight (g)	6,3 ± 1,9	8,6 ± 2,2

Experimental: Six WT plants and six independent T0 lines of 35S::AtCKX2 clone were grown on soil. After flowering the root system was washed with water, the soil was removed as far as possible and the fresh weight and dry weight was measured.

An increase in fresh and dry weight of the root biomass was also observed for F1 progeny of AtCKX2 transgenics grown in hydroponics as compared to WT, as illustrated in the following table:

Line	Wild-type	AtCKX2-38	AtCKX2-40
Fresh weight ROOT (g)	19,76 ± 6,79	33,38 ± 7,76	50,04 ± 15,59
Dry weight ROOT (g)	2,36 ± 0,43	2,61 ± 0,39	3,52 ± 1,06
Fresh weight SHOOT (g)	159,8 ± 44,53	33,66 ± 2,67	48,84 ± 11,83
Fresh weight SHOOT/ROOT ratio	8,24 ± 0,63	1,04 ± 0,18	1,08 ± 0,51

Experimental: Soil grown plants were transferred 60 days after germination to a hydroponic system (Hoagland's solution) and grown for additional 60 days. The hydroponic solution was aerated continuously and replaced by fresh solution every third day.

In summary, transgenic plants grown in hydroponic solution formed approximately 65-150% more root biomass (fresh weight) than wild type plants. The increase in dry weight was 10-50%. This difference is possibly in part due to the larger cell volume of the transgenics. This reduces the relative portion of cell walls, which forms the bulk of dry matter material. The shoot biomass was reduced to 20%-70% of wild type shoots. The

difference in fresh weight leads to a shift in the shoot/root ratio, which was approximately 8 in wild type but approximately 1 in the transgenic clones.

Conclusion:

- 5 An increase in root growth and biomass was observed for *AtCKX2* transgenic seedlings and adult plants grown under different conditions compared to WT controls despite the fact that growth of the aerial plant parts is reduced. Quantitative differences were observed between different transgenic plants: higher increases in root biomass were observed for the strongest expressing clones.

10

Reproductive development

- The onset of flowering in *AtCKX2* transgenics was delayed, the number of flowers and the seed yield per capsule was reduced. These effects were very similar to those observed in the *AtCKX1* transgenic plants but they were less prominent in the *AtCKX2* transgenics, as indicated in the tables below. The size of flowers was not altered in transgenic plants and the weight of the individual seeds was comparable to the weight of seeds from wild type plants.

20

A. Onset of flowering

Line	Wild-type	AtCKX1-48	AtCKX1-50	AtCKX2-38	AtCKX2-40
Flowering time (DAG)	106,2 ± 3,3	193,3 ± 4,3	191,8 ± 3,8	140,6 ± 6,5	121,9 ± 9,8

Experimental: Data collected for at least ten plants per line. The full elongation of the first flower was defined as onset of flowering. DAG = days after germination.

25

B. Number of seed capsules per plant

Line	Wild-type	AtCKX1-48	AtCKX1-50	AtCKX2-38	AtCKX2-40
Number of capsules	83,33 ± 5,13	2,00 ± 1,00	2,60 ± 1,67	4,30 ± 2,58	n.d.

30

Experimental: Number of seed capsules was determined at least from 5 different plants. Please note that these plants were grown under green house conditions during winter time. This affects negatively the number of flowers that are formed,

in particular in the transgenic clones. However, the general picture that they form a reduced number of flowers is correct. n.d., not determined

C. Seed yield / capsule (mg)

Line	Wild-type	AtCKX1-48	AtCKX1-50	AtCKX2-38	AtCKX2-40
Seed/capsule (mg)	87,41 ± 28,75	23,83 ± 13,36	61,8 ± 40,66	46,98 ± 29,30	n.d.

Experimental: Seed yield was determined for at least 12 seed capsules. The size of seed capsules was very variable, hence the large standard deviations. n.d., not determined

D. Weight of 100 seeds (mg)

Line	Wild-type	AtCKX1-48	AtCKX1-50	AtCKX2-38	AtCKX2-40
Seeds weight (mg)	9,73 ± 0,44	10,70 ± 1,60	9,54 ± 0,94	10,16 ± 0,47	n.d.

Experimental: The seed biomass was determined as the weight of 100 seed from at least 5 different seed capsules. n.d., not determined

3.2 In Arabidopsis:

The following morphometric data were obtained for *AtCKX2* transgenics:

Root development

A. Total length of the root system

Line	Wild-type	AtCKX2-2	AtCKX2-5
Length (mm)	32,5	50,6	48,5

B. Primary root length

Line	Wild-type	AtCKX2-2	AtCKX2-5
Length (mm)	32,3 ± 3,8	30,7 ± 4,8	31,6 ± 6,8

C. Lateral roots length

Line	Wild-type	AtCKX2-2	AtCKX2-5
Length (mm)	0,2 ± 0,4	5,5 ± 9,0	1,9 ± 2,5

D. Adventitious roots length

Line	Wild-type	AtCKX2-2	AtCKX2-5
Length (mm)	0,03 ± 0,18	14,4 ± 10,2	14,9 ± 9,1

E. Number of lateral roots (LR)

Line	Wild-type	AtCKX2-2	AtCKX2-5
Number of LR	0,3 ± 0,5	2,9 ± 2,3	1,9 ± 1,0

F. Number of adventitious roots (AR)

Line	Wild-type	AtCKX2-2	AtCKX2-5
Number of AR	0,03 ± 0,18	1,8 ± 0,9	1,8 ± 1,0

Experimental: Measurements were carried out on plants 8 d.a.g. in vitro on MS medium. At least 17 plants per line were scored.

Shoot developmentLeaf surface

Line	Wild-type	AtCKX2-2 T2 heterozygous plants	AtCKX2-5 T2 heterozygous plants	AtCKX2-9 T2 heterozygous plants
Leaf surface (cm ²)	21,16 ± 1,73	8,20 ± 2,35	8,22 ± 0,55	7,72 ± 0,85

Experimental: Leaf surface area of main rosette leaves formed after 30 days after germination was measured. 3 plants per clone were analysed.

Reproductive developmentOnset of flowering

Line	Wild-type	AtCKX1-11 T3 heterozygous plants	AtCKX2-2 T2 heterozygous plants	AtCKX2-5 T2 heterozygous plants
Flowering time (DAG)	43,6 ± 5,8	69,7 ± 9,4	51,2 ± 4,1	45,1 ± 6,9

Experimental: Plants were grown under greenhouse condition. At least 13 plants per clone were analysed. DAG = days after germination.

Conclusion: Arabidopsis *AtCKX2* transgenics had reduced leaf biomass and a dwarfing phenotype similar to *AtCKX1* transgenics (compare Figure 5 with Figure 4 F). The total root system was also enlarged in *AtCKX2* transgenic Arabidopsis. The total root length is increased approximately 50% in *AtCKX2* transgenics. The *AtCKX1* transgenics have longer primary roots, more side roots and form more adventitious roots. *AtCKX2* transgenics lack the enhanced growth of the primary root but form more side roots and lateral roots than WT.

10 **Summary:**

The phenotypes observed for *AtCKX2* transgenics were very similar but not identical to the *AtCKX1* transgenics, which in turn were very similar but not identical to the results obtained for the tobacco transgenics. This confirms the general nature of the consequences of a reduced cytokinin content in these two plant species and therefore, similar phenotypes can be expected in other plant species as well. The main difference between tobacco and Arabidopsis is the lack of enhanced primary root growth in *AtCKX2* overexpressing plants (data not shown).

20 **Example 5. Transgenic plants overexpressing *AtCKX3* showed increased cytokinin oxidase activity and altered plant morphology**

1. Description of the cloning process

The following primers were used to PCR amplify the *AtCKX3* gene from *Arabidopsis thaliana*, accession Columbia (non-homologous sequences used for cloning are in lower case):

Sequence of 5' primer: gcggtaccTTCATTGATAAGAATCAAGCTATTCA (SEQ ID NO:17)

Sequence of 3' primer: gcggtaccCAAAGTGGTGAGAACGACTAACA (SEQ ID NO:18)

30 A 3397-bp PCR fragment, produced by this PCR amplification, was inserted in the KpnI site of pBluescript. The insert was sequenced to confirm that the PCR product has no sequence changes as compared to the gene. The KpnI/KpnI fragment of this vector was subcloned in the KpnI site downstream of a modified CaMV 35S promoter (carrying three

tetracycline operator sequences) in the binary vector pBinHyg-Tx (Gatz *et al.*, 1992). The resulting construct was introduced into tobacco and *Arabidopsis thaliana* through *Agrobacterium*-mediated transformation, using standard transformation protocols.

5 2. Molecular analysis of the transgenic lines

Several transgenic tobacco lines were identified that synthesize the *AtCKX3* transcript at high levels (Fig 11 A.). Transgenic tobacco lines expressing *AtCKX3* transcript also showed increased cytokinin oxidase activity. This is exemplified for three plants in Table 8. This proves that the *AtCKX3* gene encodes a protein with cytokinin oxidase activity.

10

Table 8. Cytokinin oxidase activity in *AtCKX4* transgenic plant tissues

Sample		
Plant species and tissue	Plant line	Cytokinin oxidase activity (nmol Ade/mg protein.h)
tobacco leaves	SNN wild-type	0.011
	CKX3-SNN-3	0.049
	CKX3-SNN-6	0.053
	CKX3-SNN-21	0.05

15 3. Plant phenotypic analysis

The phenotypes generated by overexpression of the *AtCKX3* gene in tobacco and *Arabidopsis* were basically similar as those of *AtCKX1* and *AtCKX2* expressing plants, i.e. enhanced rooting and dwarfing. However, overexpression of the *AtCKX3* gene in tobacco resulted in a stronger phenotype compared to *AtCKX2*. In this sense *AtCKX3* overexpression was more similar to *AtCKX1* overexpression.

20

Example 6. Transgenic plants overexpressing AtCKX4 showed increased cytokinin oxidase activity and altered plant morphology

1. Description of the cloning process

- 5 The following primers were used to PCR amplify the AtCKX4 gene from *Arabidopsis thaliana*, accession Columbia (non-homologous sequences used for cloning are in lower case):

Sequence of 5' primer: gcggtaccCCCATTAACCTACCCGTTTG (SEQ ID NO:19)

Sequence of 3' primer: gcggtaccAGACGATGAACGTACTTGTCTGTA (SEQ ID NO:20)

10

- A 2890-bp PCR fragment, produced by this PCR amplification, was inserted in the KpnI site of pBluescript. The insert was sequenced to confirm that the PCR product has no sequence changes as compared to the gene. The KpnI/KpnI fragment of this vector was subcloned in the KpnI site downstream of a modified CaMV 35S promoter (carrying three
15 tetracycline operator sequences) in the binary vector pBinHyg-Tx (Gatz *et al.*, 1992). The resulting construct was introduced into tobacco and *Arabidopsis thaliana* through *Agrobacterium*-mediated transformation, using standard transformation protocols.

2. Molecular analysis of the transgenic lines

- 20 Several transgenic tobacco lines synthesized the AtCKX4 transcript at high levels (Fig 11 B.). Transgenic lines expressing AtCKX4 transcript also showed increased cytokinin oxidase activity. This is exemplified for 3 *Arabidopsis* and 3 tobacco lines in Table 9. This result proves that the AtCKX4 gene encodes a protein with cytokinin oxidase activity.

Table 9. Cytokinin oxidase activity in *AtCKX4* transgenic plant tissues

Sample		
Plant species and tissue	Plant line	Cytokinin oxidase activity (nmol Ade/mg protein.h)
<i>Arabidopsis</i> callus	Col-0 wild-type	0.037
	CKX4-37	0.244
	CKX4-40	0.258
	CKX4-41	0.320
tobacco leaves	SNN wild-type	0.011
	CKX4-SNN-3	0.089
	CKX4-SNN-18	0.085
	CKX4-SNN-27	0.096

5 Overall, the data showed that the apparent K_m values for the four cytokinin oxidases were in the range of 0.2 to 9.5 μM with iP as substrate, which further demonstrates that the proteins encoded by *AtCKX1* through 4 are indeed cytokinin oxidase enzymes as disclosed herein.

10 3. Plant phenotypic analysis

The phenotypes generated by overexpression of the *AtCKX4* gene in tobacco and *Arabidopsis* were basically similar as those of *AtCKX1* and *AtCKX2* expressing plants, i.e. enhanced rooting, reduced apical dominance, dwarfing and yellowing of intercostal regions in older leaves of tobacco. An additional phenotype in tobacco was lanceolate
 15 leaves (altered length-to-width ratio).

General observations of *AtCKX* overexpressing tobacco plants

Overall, the phenotypic analysis demonstrated that *AtCKX* gene overexpression caused drastic developmental alterations in the plant shoot and root system in tobacco, including
 20 enhanced development of the root system and dwarfing of the aerial plant part. Other effects such as altered leaf senescence, formation of adventitious root on stems, and others were also observed as disclosed herein. The alterations were very similar, but not

identical, for the different genes. In tobacco, *AtCKX1* and *AtCKX3* overexpressors were alike as were *AtCKX2* and *AtCKX4*. Generally, the two former showed higher expression of the traits, particularly in the shoot. Therefore, a particular cytokinin oxidase gene may be preferred for achieving the phenotypes that are described in the
5 embodiments of this invention.

Example 7. Cloning of the *AtCKX5* gene

The following primers were used to PCR amplify the *AtCKX5* gene from *Arabidopsis thaliana*, accession Columbia (non-homologous sequences used for cloning are in lower
10 case):

Sequence of 5' primer: ggggtaccTTGATGAATCGTGAAATGAC (SEQ ID NO:21)

Sequence of 3' primer: ggggtaccCTTTCCTCTTGGTTTGTCTGT (SEQ ID NO:22)

The sequence of the 5' primer includes the two potential startcodons of the *AtCKX5*
15 protein, the most 5' startcodon is underlined and a second ATG is indicated in italics. A 2843-bp PCR fragment, produced by this PCR amplification, was inserted as a blunt-end product in pCR-Blunt II-TOPO cloning vector (Invitrogen).

Example 8. Cloning of the *AtCKX6* gene

The following primers were used to PCR amplify the *AtCKX6* gene from *Arabidopsis thaliana*, accession Columbia (non-homologous sequences used for cloning are in lower
20 case):

Sequence of 5' primer: gctctagaTCAGGAAAAGAACCATGCTTATAG (SEQ ID NO:23)

Sequence of 3' primer: gctctagaTCATGAGTATGAGACTGCCTTTTG (SEQ ID NO:24)
25

A 1949-bp PCR fragment, produced by this PCR amplification, was inserted as a blunt-end product in pCR-Blunt II-TOPO cloning vector (Invitrogen).

Example 9. Tobacco seedling growth test demonstrated early vigor of *AtCKX* transgenics 30

Seeds of *AtCKX1-50* and *AtCKX2-38* overexpressing transgenics and WT tobacco were sown *in vitro* on MS medium, brought to culture room 4 days after cold treatment and

germinated after 6 days. Observations on seedling growth were made 10 days after germination (see also Figure 8C) and are summarized below. At least 20 individuals were scored per clone. Similar data have been obtained in two other experiments.

A. Total length of the root system

Line	Wild-type	AtCKX1-50	AtCKX2-38
Length (mm)	61,1	122,0	106,5

B. Primary root length

Line	Wild-type	AtCKX1-50	AtCKX2-38
Length (mm)	32,3 \pm 2,6	50,8 \pm 4,5	52,4 \pm 4,8

C. Lateral roots length

Line	Wild-type	AtCKX1-50	AtCKX2-38
Length (mm)	9,8 \pm 5,5	18,0 \pm 8,1	13,0 \pm 6,0

D. Adventitious roots length

Line	Wild-type	AtCKX1-50	AtCKX2-38
Length (mm)	19,0 \pm 5,0	53,0 \pm 12,0	42,0 \pm 9,8

E. Number of lateral roots (LR)

Line	Wild-type	AtCKX1-50	AtCKX2-38
Number of LR	1,9 \pm 0,9	6,5 \pm 2,2	5,6 \pm 2,0

F. Number of adventitious roots (AR)

Line	Wild-type	AtCKX1-50	AtCKX2-38
Number of AR	2,2 \pm 0,6	3,5 \pm 0,9	3,6 \pm 1,3

AtCKX1 and AtCKX2 plants, general observations:

Seedlings of *AtCKX1* and *AtCKX2* overexpressing tobacco plants had 60% more adventitious roots and three times more lateral roots than untransformed control plants 10 days after germination. The length of the primary root was increased by about 70%. This – together with more and longer side roots and secondary roots – resulted in a 70-

100% increase in total root length. These results showed that overexpression of cytokinin oxidase enhances the growth and development of both the main root and the adventitious roots, resulting in early vigor.

5 **Example 10. Histological analysis of altered plant morphology in *AtCKX1* overexpressing tobacco plants**

Microscopic analysis of different tissues revealed that the morphological changes in *AtCKX* transgenics are reflected by distinct changes in cell number and rate of cell formation (see Figure 10). The shoot apical meristem (SAM) of *AtCKX1* transgenics was smaller than in wild type and fewer cells occupy the space between the central zone and the peripheral zone of lateral organ formation, but the cells were of the same size (Figure 10 A). The reduced cell number and size of the SAM as a consequence of a reduced cytokinin content indicates that cytokinins have a role in the control of SAM proliferation.

15 No obvious changes in the differentiation pattern occurred, suggesting that the spatial organization of the differentiation zones in the SAM is largely independent from cell number and from the local cytokinin concentration. The overall tissue pattern of leaves in cytokinin oxidase overexpressers was unchanged. However, the size of the phloem and xylem was significantly reduced (Figure 10 B). By contrast, the average cell size of leaf parenchyma and epidermal cells was increased four- to fivefold (Figure 10 C, D). New cells of *AtCKX1* transgenics are formed at 3-4% of the rate of wild type leaves and final leaf cell number was estimated to be in the range of 5-6% of wild type. This indicates an absolute requirement for cytokinins in leaves to maintain the cell division cycle. Neither cell size nor cell form of floral organs was altered and seed yield per capsule was similar

25 in wild type and *AtCKX* transgenic plants. The cell population of root meristems of *AtCKX1* transgenic plants was enlarged approximately 4-fold and the cell numbers in both the central and lateral columnella were enhanced (Figure 10 E, F). The final root diameter was increased by 60% due to an increased diameter of all types of root cells. The radial root patterns was identical in wild type and transgenics, with the exception that frequently a fourth layer of cortex cells was noted in transgenic roots (Figure 10 G).

30 The increased cell number and the slightly reduced cell length indicates that the enhanced root growth is due to an increased number of cycling cells rather than increased cell growth. In the presence of lowered cytokinin content, root meristem cells

must undergo additional rounds of mitosis before they leave the meristem and start to elongate. The exit from the meristem is therefore regulated by a mechanism that is sensitive to cytokinins. Apparently, cytokinins have a negative regulatory role in the root meristem and wild type cytokinin concentrations are inhibitory to the development of a maximal root system. Therefore, reducing the level of active cytokinins by overexpressing cytokinin oxidases stimulates root development, which results in an increase in the size of the root with more lateral and adventitious roots as compared to WT plants.

Example 11. *AtCKX1* and *AtCKX2*-overexpressing tobacco plants had a reduced cytokinin content.

Among the 16 different cytokinin metabolites that were measured, the greatest change occurred in the iP-type cytokinins in *AtCKX2* overexpressers (Table 10): the overall decrease in the content of iP-type cytokinins is more pronounced in *AtCKX2* expressing plants than in *AtCKX1* transgenics. *AtCKX1* transgenics showed a stronger phenotype in the shoot. It is not known which cytokinin metabolite is relevant for the different traits that were analyzed. It may be that different cytokinin forms play different roles in the various development processes. Smaller alterations were noted for Z-type cytokinins, which could be due to a different accessibility of the substrate or a lower substrate specificity of the protein. The total content of iP and Z metabolites in individual transgenic clones was between 31% and 63% of wild type. The cytokinin reserve pool of O-glucosides was also lowered in the transgenics (Table 10). The concentration of N-glucosides and DHZ-type cytokinins was very low and was not or only marginally, altered in transgenic seedlings (data not shown).

Table 10. Cytokinin content of *AtCKX* transgenic plants. Cytokinin extraction, immunopurification, HPLC separation and quantification by ELISA methods was carried out as described by Faiss et al., 1997. Three independently pooled samples of approximately 100 two week old seedlings (2.5 g per sample) were analyzed for each clone. Concentrations are in pmol x g fresh weight⁻¹. Abbreviations: iP, N⁶-(Δ^2 isopentenyl)adenine; iPR, N⁶-(Δ^2 isopentenyl)adenine riboside; iPRP, N⁶-(Δ^2 isopentenyl)adenine riboside 5'-monophosphate; Z, *trans*-zeatin; ZR, zeatin riboside; ZRP, zeatin riboside 5'-monophosphate; ZOG, zeatin O-glucoside; ZROG, zeatin riboside O-glucoside.

Line	WT	AtCKX1-2		AtCKX1-28		AtCKX2-38		AtCKX2-40	
Cytokinin metabolite	Concentration	Concentration	% of WT	Concentration	% of WT	Concentration	% of WT	Concentration	% of WT
iP	5,90 ± 1,80	4,76 ± 0,82	81	4,94 ± 2,62	84	1,82 ± 0,44	31	2,85 ± 0,62	48
iPR	2,36 ± 0,74	1,53 ± 0,14	65	0,75 ± 0,27	32	0,55 ± 0,39	23	0,89 ± 0,07	38
iPRP	3,32 ± 0,73	0,87 ± 0,26	26	1,12 ± 0,13	34	0,80 ± 0,48	24	1,68 ± 0,45	51
Z	0,24 ± 0,06	0,17 ± 0,02	71	0,22 ± 0,03	92	0,21 ± 0,06	88	0,22 ± 0,02	92
ZR	0,60 ± 0,13	0,32 ± 0,12	53	0,34 ± 0,03	57	0,34 ± 0,15	57	0,32 ± 0,05	53
ZRP	0,39 ± 0,17	0,42 ± 0,11	107	0,28 ± 0,15	72	0,06 ± 0,01	15	0,17 ± 0,06	44
ZOG	0,46 ± 0,20	0,32 ± 0,09	70	0,26 ± 0,13	57	0,20 ± 0,07	43	0,12 ± 0,02	26
ZROG	0,48 ± 0,17	0,30 ± 0,06	63	0,47 ± 0,02	98	0,23 ± 0,05	48	0,30 ± 0,13	63
Total	13,75	8,69	63	8,38	61	4,21	31	6,55	48

Example 12. Grafting experiments showed that dwarfing and enhanced root development due to *AtCKX* overexpression is confined to transgenic tissues

To investigate which phenotypic effects of cytokinin oxidase overexpression are restricted to expressing tissues, i.e. are cell- or organ-autonomous traits, grafting experiments were performed. Reciprocal grafts were made between an *AtCKX2* transgenic tobacco plant and a WT tobacco. The transgenic plant used in this experiment was *AtCKX2*-38, which displayed a strong phenotype characterized by enhanced root growth and reduced development of the aerial plant parts. As described in Example 3 through 6, these were two important phenotypes that resulted from cytokinin oxidase overexpression in tobacco and arabidopsis.

Plants were about 15 cm tall when grafted and the graft junction was about 10 cm above the soil. Figure 12 shows plants 15 weeks after grafting. The main results were that : (i) the aerial phenotype of a WT scion grafted on a transgenic rootstock was similar to the WT control graft (= WT scion on WT rootstock). Importantly, this showed that overexpression of the *AtCKX2* transgene in the rootstock did not induce dwarfing of the non-transgenic aerial parts of the plant (see Figure 12 A). Improved root growth of the transgenic rootstock was maintained, indicating that improved root growth of *AtCKX* transgenics is autonomous and does not depend on an *AtCKX* transgenic shoot (Figure 12 C). Interestingly, the WT scions grafted on the transgenic rootstocks looked healthier and were better developed. Notably, senescence of the basal leaves was retarded in these plants (see Figure 12 A); (ii) the transgenic scion grafted on the WT rootstock looked similar to the aerial part of the transgenic plant from which it was derived, i.e. the shoot dwarfing phenotype is also autonomous and not dependent on the improved root growth (see Figure 12 B).

In addition to the above-mentioned better appearance of WT shoots grafted on a transgenic rootstock, the formation of adventitious roots on the basal part of WT shoots was noted (Figure 12 D, right plant). Formation of adventitious roots also occurred on the stem of *AtCKX* transgenics but not on stems of WT control grafts (Figure 12 D, left plant) and therefore seems to be a non-autonomous trait.

In summary, it is disclosed in this invention that enhanced root formation and dwarfing of the shoot in *AtCKX* overexpressing tobacco are autonomous traits and can be uncoupled by grafting procedures. Surprisingly, grafting of a WT scion on an *AtCKX*

transgenic rootstock resulted in more vigorously growing plants and retardation of leaf senescence.

As an alternative to grafting, tissue-specific promoters could be used for uncoupling the autonomous phenotypic effects of cytokinin overexpression. Therefore, it is disclosed in this invention that cytokinin oxidase overexpression in a tissue specific manner can be used to alter the morphology of a plant such as the shoot or root system.

Example 13.. Expression of an *AtCKX* gene under a root-specific promoter in transgenic plants leads to increased root production

An *AtCKX* gene (see example 4) is cloned under control of the root *clavata* homolog promoter of *Arabidopsis* (SEQ ID NO 36) , which is a promoter that drives root-specific expression. Other root-specific promoters may also be used for the purpose of this invention. See Table 5 for exemplary root-specific promoters.

Transgenic plants expressing the *AtCKX* gene specifically in the roots show increased root production without negatively affecting growth and development of the aerial parts of the plant. Positive effects on leaf senescence and growth of aerial plant parts are observed.

Example 14. Suppression of an *AtCKX* gene under a senescence-induced promoter in transgenic plants leads to delayed leaf senescence and enhanced seed yield.

A chimeric gene construct derived from an *AtCKX* gene and designed to suppress expression of endogenous cytokinin oxidase gene(s) is cloned under control of a senescence-induced promoter. For example, promoters derived from senescence-associated genes (SAG) such as the SAG12 promoter can be used (Quirino et al., 2000). Transgenic plants suppressing endogenous cytokinin oxidase gene(s) specifically in senescing leaves show delayed leaf senescence and higher seed yield without negatively affecting the morphology and growth and development of the plant.

Example 15. Overexpression of an *AtCKX* gene in the female reproductive organs leads to parthenocarpic fruit development

- 5 The open reading frame of an *AtCKX* gene is cloned under control of a promoter that confers overexpression in the female reproductive organs such as for example the DefH9 promoter from *Antirrhinum majus* or one of its homologues, which have high expression specificity in the placenta and ovules. Transgenic plants with enhanced cytokinin oxidase activity in these tissues show parthenocarpic fruit development.

REFERENCES

- WO0105985. Method to modulate the expression of genes inducing the parthenocarpic trait in plants.
- 5 Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., and Watson, J. D. (1994). "Molecular Biology of the Cell." Garland Publishing Inc.
- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D.J. (1997) "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs." *Nucl. Acids Res.* **25**, 3389-3402.
- 10 An, G., Watson, B. D., Stachel, S., Gordon, M. P., and Nester, E. W. (1985). New cloning vehicles for transformation of higher plants. *EMBO J.* **4**, 277-284.
- Armstrong, C. L., Petersen, W. P., Buchholz, W. G., Bowen, B. A., and Sulc, S. L. (1990). Factors affecting PEG-mediated stable transformation of maize protoplasts. *Plant Cell Reports* **9**, 335-339.
- 15 Banerjee, A., Pramanik, A., Bhattacharjya, S., and Balaram, P. (1996). Omega amino acids in peptide design: incorporation into helices. *Biopolymers* **39**, 769-777.
- Baron, M. H. and Baltimore, D. (1982). Antibodies against the chemically synthesized genome-linked protein of poliovirus react with native virus-specific proteins. *Cell* **28**, 395-404.
- 20 Bartel, P. L. and Fields, S. (1997). "The Yeast Two-Hybrid System." Oxford University Press.
- Benkirane, N., Guichard, G., Briand, J. P., and Muller, S. (1996). Exploration of requirements for peptidomimetic immune recognition. Antigenic and immunogenic properties of reduced peptide bond pseudopeptide analogues of a histone hexapeptide. *J.Biol Chem.* **271**, 33218-33224.
- 25 Berry, A. and Brenner, S. E. (1994). A prototype computer system for de novo protein design. *Biochem.Soc.Trans.* **22**, 1033-1036.
- Christou, P., McCabe, D. E., and Swain, W. F. (1988). Stable transformation of soybean callus by DNA-coated gold particles. *Plant Physiol.* **87**, 671-674.

Crossway, A., Oakes, J. V., Irvine, J. M., Ward, B., Knauf, V. C., and Shewmaker, C. K. (1986). Integration of foreign DNA following microinjection of tobacco mesophyll protoplasts. *Mol.Gen.Genet.* **202**, 179-185.

5 Dale, E. C. and Ow, D. W. (1990). Intra- and intermolecular site-specific recombination in plant cells mediated by bacteriophage P1 recombinase. *Gene* **91**, 79-85.

Dodds, J. H. (1985). "Plant genetic engineering." Cambridge University Press.

Doerner, P., Jorgensen, J. E., You, R., Steppuhn, J., and Lamb, C. (1996). Control of root growth and development by cyclin expression. *Nature* **380**, 520-523.

10 Dorner, B., Husar, G. M., Ostresh, J. M., and Houghten, R. A. (1996). The synthesis of peptidomimetic combinatorial libraries through successive amide alkylations. *Bioorg.Med.Chem.* **4**, 709-715.

15 Ellis, J. G., Llewellyn, D. J., Dennis, E. S., and Peacock, W. J. (1987). Maize Adh-1 promoter sequences control anaerobic regulation: addition of upstream promoter elements from constitutive genes is necessary for expression in tobacco. *EMBO J.* **6**, 11-16.

Faiss, M., Zalubilová, J., Strnad, M., Schmölling, T. (1997). Conditional transgenic expression of the ipt gene indicates a function for cytokinins in paracrine signaling in whole tobacco plants. *Plant J.* **12**, 401-415.

20 Fassina, G. and Melli, M. (1994). Identification of interactive sites of proteins and protein receptors by computer-assisted searches for complementary peptide sequences. *Immunomethods.* **5**, 114-120.

Fedoroff, N. V. and Smith, D. L. (1993). A versatile system for detecting transposition in Arabidopsis. *Plant J.* **3**, 273-289.

25 Hanahan, D. (1983). Studies on transformation of Escherichia coli with plasmids. *J.Mol.Biol* **166**, 557-580.

Hansen, G. and Chilton, M. D. (1996). "Agrolistic" transformation of plant cells: integration of T-strands generated in planta. *Proc.Natl.Acad.Sci.U.S.A* **93**, 14978-14983.

- Hansen, G., Shillito, R. D., and Chilton, M. D. (1997). T-strand integration in maize protoplasts after codelivery of a T-DNA substrate and virulence genes. *Proc.Natl.Acad.Sci.U.S.A* **94**, 11726-11730.
- Hanson, B., Engler, D., Moy, Y., Newman, B., Ralston, E., and Gutterson, N. (1999). A
5 simple method to enrich an Agrobacterium-transformed population for plants containing only T-DNA sequences. *Plant J.* **19**, 727-734.
- Harlow, E. and Lane, D. (1988). "Antibodies: A Laboratory Manual." Cold Spring Harbor Laboratory Press.
- Herrera-Estrella, L., De Block, M., Messens, E. H. J. P., Van Montagu, M., and Schell, J.
10 (1983). Chimeric genes as dominant selectable markers in plant cells. *EMBO J.* **2**, 987-995.
- Hoffman, D. L., Laiter, S., Singh, R. K., Vaisman, I. I., and Tropsha, A. (1995). Rapid protein structure classification using one-dimensional structure profiles on the bioSCAN parallel computer. *Comput.Appl.Biosci.* **11**, 675-679.
- 15 Krens, F. A., Molendijk, L., Wullems, G. J., and Schilperoort, R. A. (1982). *In vitro* transformation of plant protoplasts with Ti-plasmid DNA. *Nature* **296**, 72-74.
- Lerner, R. A. (1982). Tapping the immunological repertoire to produce antibodies of predetermined specificity. *Nature* **299**, 593-596.
- Lerner, R. A., Green, N., Alexander, H., Liu, F. T., Sutcliffe, J. G., and Shinnick, T. M.
20 (1981). Chemically synthesized peptides predicted from the nucleotide sequence of the hepatitis B virus genome elicit antibodies reactive with the native envelope protein of Dane particles. *Proc.Natl.Acad.Sci.U.S.A* **78**, 3403-3407.
- Liddle, J. E. and Cryer, A. (1991). "A Practical Guide to Monoclonal Antibodies." Wiley New York.
- 25 Loffler, J., Langui, D., Probst, A., and Huber, G. (1994). Accumulation of a 50 kDa N-terminal fragment of beta-APP695 in Alzheimer's disease hippocampus and neocortex. *Neurochem.Int.* **24**, 281-288.
- Monge, A., Lathrop, E. J., Gunn, J. R., Shenkin, P. S., and Friesner, R. A. (1995). Computer modeling of protein folding: conformational and energetic analysis of reduced
30 and detailed protein models. *J.Mol.Biol* **247**, 995-1012.

Morris, R.O. et al. (1999). Isolation of a gene encoding a glycosylated cytokinin oxidase from maize. *Biochem. Biophys. Res. Commun.* **255**, 328-333

Motyka, V., Faiss, M., Strnad, M., Kaminek, M. and Schmuelling, T. (1996). Changes in cytokinin content and cytokinin oxidase activity in response to derepression of *ipt* gene transcription in transgenic tobacco calli and plants. *Plant Physiol.* **112**, 1035-1043.

Murakami, T., Simonds, W. F., and Spiegel, A. M. (1992). Site-specific antibodies directed against G protein beta and gamma subunits: effects on alpha and beta gamma subunit interaction. *Biochemistry* **31**, 2905-2911.

Olszewski, K. A., Kolinski, A., and Skolnick, J. (1996). Folding simulations and computer redesign of protein A three-helix bundle motifs. *Proteins* **25**, 286-299.

Osborne, B. I., Wirtz, U., and Baker, B. (1995). A system for insertional mutagenesis and chromosomal rearrangement using the Ds transposon and Cre-lox. *Plant J.* **7**, 687-701.

Ostresh, J. M., Blondelle, S. E., Dorner, B., and Houghten, R. A. (1996). Generation and use of nonsupport-bound peptide and peptidomimetic combinatorial libraries. *Methods Enzymol.* **267**, 220-234.

Pabo, C. O. and Suchanek, E. G. (1986). Computer-aided model-building strategies for protein design. *Biochemistry* **25**, 5987-5991.

Paszkowski, J., Shillito, R. D., Saul, M., Mandak, V., and Hohn, T. H. B. P. I. (1984). Direct gene transfer to plants. *EMBO J.* **3**, 2717-2722.

Peralta, E. G., Hellmiss, R., and Ream, W. (1986). Overdrive, a T-DNA transmission enhancer on the *A. tumefaciens* tumour-inducing plasmid. *EMBO J.* **5**, 1137-1142.

Quirino, B.F., Noh, Y.-S., Himelbau, E., and Amasino, R.M. (2000). Molecular aspects of leaf senescence. *Trends in Plant Science* **5**, 278-282.

Renouf, D. V. and Hounsell, E. F. (1995). Molecular modelling of glycoproteins by homology with non-glycosylated protein domains, computer simulated glycosylation and molecular dynamics. *Adv.Exp.Med.Biol* **376**, 37-45.

Rinaldi, A.C. and Comandini, O. (1999). Cytokinin oxidase strikes again. *Trends in Plant Sc.* **4**, 300.

- Rose, R. B., Craik, C. S., Douglas, N. L., and Stroud, R. M. (1996). Three-dimensional structures of HIV-1 and SIV protease product complexes. *Biochemistry* **35**, 12933-12944.
- 5 Rutenber, E. E., McPhee, F., Kaplan, A. P., Gallion, S. L., Hogan, J. C., Jr., Craik, C. S., and Stroud, R. M. (1996). A new class of HIV-1 protease inhibitor: the crystallographic structure, inhibition and chemical synthesis of an aminimide peptide isostere. *Bioorg.Med.Chem.* **4**, 1545-1558.
- Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989). "Molecular Cloning: A Laboratory Manual." Cold Spring Harbor Laboratory Press.
- 10 Schlappi, M., Smith, D., and Fedoroff, N. (1993). TnpA trans-activates methylated maize Suppressor-mutator transposable elements in transgenic tobacco. *Genetics* **133**, 1009-1021.
- Shioda, T., Andriole, S., Yahata, T., and Isselbacher, K. J. (2000). A green fluorescent protein-reporter mammalian two-hybrid system with extrachromosomal maintenance of a prey expression plasmid: Application to interaction screening. *Proc.Natl.Acad.Sci.U.S.A* **97**, 5220-5224.
- 15 Tamura, R. N., Cooper, H. M., Collo, G., and Quaranta, V. (1991). Cell type-specific integrin variants with alternative alpha chain cytoplasmic domains. *Proc.Natl.Acad.Sci.U.S.A* **88**, 10183-10187.
- 20 Van Haaren, M. J., Sedee, N. J., Schilperoort, R. A., and Hooykaas, P. J. (1987). Overdrive is a T-region transfer enhancer which stimulates T-strand production in *Agrobacterium tumefaciens*. *Nucleic Acids Res.* **15**, 8983-8997.
- Van Sluys, M. A., Tempe, J., and Fedoroff, N. (1987). Studies on the introduction and mobility of the maize Activator element in *Arabidopsis thaliana* and *Daucus carota*.
- 25 *EMBO J.* **6**, 3881-3889.
- Wang, K., Genetello, C., Van Montagu, M., and Zambryski, P. C. (1987). Sequence context of the T-DNA border repeat element determines its relative activity during T-DNA transfer to plant cells. *Mol.Gen.Genet.* **210**, 338-346.
- Woulfe, J., Lafortune, L., de Nadai, F., Kitabgi, P., and Beaudet, A. (1994). Post-
- 30 translational processing of the neurotensin/neuromedin N precursor in the central

nervous system of the rat—II. Immunohistochemical localization of maturation products. *Neuroscience* **60**, 167-181.

- Zhang, Y. L., Dawe, A. L., Jiang, Y., Becker, J. M., and Naider, F. (1996). A superactive peptidomimetic analog of a farnesylated dodecapeptide yeast pheromone.
- 5 *Biochem.Biophys.Res.Commun.* **224**, 327-331.

CLAIMS

1. Use of a nucleic acid encoding a plant cytokinin oxidase or encoding a protein that reduces the level of active cytokinins in plants or plant parts for stimulating root growth or
5 for enhancing the formation of lateral or adventitious roots or for altering root geotropism.
2. A method for stimulating root growth or for enhancing the formation of lateral or adventitious roots or for altering root geotropism comprising expression of a nucleic acid encoding a plant cytokinin oxidase selected from the group consisting of:
 - 10 (a) nucleic acids comprising a DNA sequence as given in any of SEQ ID NOs 27, 1, 3, 5, 7, 9, 11, 25, 26, 28 to 31, 33 or 34, or the complement thereof,
 - (b) nucleic acids comprising the RNA sequences corresponding to any of SEQ ID NOs 27, 1, 3, 5, 7, 9, 11, 25, 26, 28 to 31, 33 or 34, or the complement thereof,
 - (c) nucleic acids specifically hybridizing to any of SEQ ID NOs 27, 1, 3, 5, 7, 9, 11,
15 25, 26, 28 to 31, 33 or 34, or to the complement thereof,
 - (d) nucleic acids encoding a protein comprising the amino acid sequence as given in any of SEQ ID NOs 2, 4, 6, 8, 10, 12, 32 or 35, or the complement thereof,
 - (e) nucleic acids as defined in any of (a) to (d) characterized in that said nucleic acid is DNA, genomic DNA, cDNA, synthetic DNA or RNA wherein T is replaced by U,
 - 20 (f) a nucleic acid which is degenerated to a nucleic acid as given in any of SEQ ID NOs 27, 1, 3, 5, 7, 9, 11, 25, 26, 28 to 31, 33 or 34, or which is degenerated to a nucleic acid as defined in any of (a) to (e) as a result of the genetic code,
 - (g) nucleic acids which are diverging from a nucleic acid encoding a protein as given in any of SEQ ID NOs 2, 4, 6, 8, 10, 12 or 35 or which is diverging from a
25 nucleic acid as defined in any of (a) to (e), due to the differences in codon usage between the organisms,
 - (h) nucleic acids encoding a protein as given in SEQ ID NOs 2, 4, 6, 8, 10, 12 or 35 or nucleic acids as defined in (a) to (e) which are diverging due to the differences between alleles,
 - 30 (i) nucleic acids encoding a protein as given in any of SEQ ID NOs 2, 4, 6, 8, 10, 12 or 35,
 - (j) functional fragments of nucleic acids as defined in any of (a) to (i) having the biological activity of a cytokinin oxidase, and

(k) nucleic acids encoding a plant cytokinin oxidase,
or comprising expression, preferably in roots, of a nucleic acid encoding a protein that
reduces the level of active cytokinins in plants or plant parts.

- 5 3. An isolated nucleic acid encoding a novel plant protein having cytokinin oxidase
activity selected from the group consisting of:
- (a) a nucleic acid comprising a DNA sequence as given in any of SEQ ID NOs 29,
3, 5, 9, 26, 27, 31, 33 or 34, or the complement thereof,
 - (b) a nucleic acid comprising the RNA sequences corresponding to any of SEQ ID
10 NOs 29, 3, 5, 9, 26, 27, 31, 33 or 34, or the complement thereof,
 - (c) a nucleic acid specifically hybridizing to a nucleic acid as given in any of SEQ ID
NOs 29, 3, 5, 9, 26, 27, 31, 33 or 34, or the complement thereof,
 - (d) a nucleic acid encoding a protein with an amino acid sequence comprising the
15 polypeptide as given in SEQ ID NO 32 and which is at least 70% similar to the
amino acid sequence as given in SEQ ID NO 4,
 - (e) a nucleic acid encoding a protein with an amino acid sequence which is at least
47% similar to the amino acid sequence as given in SEQ ID NO 6,
 - (f) a nucleic acid encoding a protein with an amino acid sequence which is at least
47% similar to the amino acid sequence as given in SEQ ID NO 10 or 35,
 - 20 (g) a nucleic acid encoding a protein comprising the amino acid sequence as given
in any of SEQ ID NOs 4, 6, 10, 32 or 35,
 - (h) a nucleic acid which is degenerated to a nucleic acid as given in any of SEQ ID
NOs 29, 3, 5, 9, 26, 27, 33 or 34 or which is degenerated to a nucleic acid as
defined in any of (a) to (g) as a result of the genetic code,
 - 25 (i) a nucleic acid which is diverging from a nucleic acid encoding a protein as given
in any of SEQ ID NOs 4, 6, 10 or 35 or which is diverging from a nucleic acid as
defined in any of (a) to (g) due to the differences in codon usage between the
organisms,
 - (j) a nucleic acid encoding a protein as given in SEQ ID NOs 4, 6, 10 or 35, or a
30 nucleic acid as defined in (a) to (g) which is diverging due to the differences
between alleles,
 - (k) a nucleic acid encoding an immunologically active fragment of a cytokinin
oxidase encoded by a nucleic acid as given in any of SEQ ID NOs 29, 3, 5, 9, 26,

27, 31, 33 or 34, or an immunologically active fragment of a nucleic acid as defined in any of (a) to (j),

(l) a nucleic acid encoding a functional fragment of a cytokinin oxidase encoded by a nucleic acid as given in any of SEQ ID NOs 29, 3, 5, 9, 26, 27, 31, 33 or 34, or
5 a functional fragment of a nucleic acid as defined in any of (a) to (j), wherein said fragment has the biological activity of a cytokinin oxidase, and

(m) a nucleic acid encoding a protein as defined in SEQ ID NO 4, 6, 10 or 35,

provided that said nucleic acid is not the nucleic acid as deposited under any of the following Genbank accession numbers: AC005917, AB024035, and AC023754.

- 10 4. An isolated nucleic acid according to claim 3 which is DNA, cDNA, genomic DNA or synthetic DNA, or RNA wherein T is replaced by U.
- 5 5. A nucleic acid molecule of at least 15 nucleotides in length hybridizing specifically with a nucleic acid of claim 3 or 4.
- 15 6. A nucleic acid molecule of at least 15 nucleotides in length specifically amplifying a nucleic acid of claim 3 or 4.
7. A vector comprising a nucleic acid of claim 3 or 4.
8. A vector according to claim 7 which is an expression vector wherein the nucleic acid is operably linked to one or more control sequences allowing the expression of said nucleic acid in prokaryotic and/or eukaryotic host cells.
- 20 9. A host cell containing a nucleic acid according to claim 3 or 4 or a vector according to claim 7 or 8.
10. The host cell of claim 9, wherein the host cell is a bacterial, insect, fungal, plant or animal cell.
11. An isolated polypeptide encodable by a nucleic acid of claim 3 or 4, or a homologue
25 or a derivative thereof, or an immunologically active or a functional fragment thereof.
12. The polypeptide of claim 11 which has an amino acid sequence as given in any of SEQ ID NOs 4, 6, 10 or 35, or a homologue or a derivative thereof, or an immunologically active or a functional fragment thereof.

13. A method for producing a polypeptide according to claim 11 or 12 comprising culturing a host cell of claim 9 or 10 under conditions allowing the expression of the polypeptide and recovering the produced polypeptide from the culture.
14. An antibody specifically recognizing a polypeptide of claim 11 or 12 or a specific epitope thereof.
15. A method for the production of transgenic plants, plant cells or plant tissues comprising the introduction of a nucleic acid of claim 3 or 4 in an expressible format or a vector of claim 7 or 8 in said plant, plant cell or plant tissue.
16. A method for the production of altered plants, plant cells or plant tissues comprising the introduction of a polypeptide of claim 11 or 12 directly into a cell, a tissue or an organ of said plant.
17. A method for effecting the expression of a polypeptide of claim 11 or 12 comprising the introduction of a nucleic acid of claim 3 or 4 operably linked to one or more control sequences or a vector of claim 7 or 8 stably into the genome of a plant cell.
18. The method of claim 16 or 17 further comprising regenerating a plant from said plant cell.
19. A transgenic plant cell comprising a nucleic acid of claim 3 or 4 which is operably linked to regulatory elements allowing transcription and/or expression of said nucleic acid in plant cells or a transgenic plant cell obtainable by a method of claim 16 or 17.
20. The transgenic plant cell of claim 18 wherein said nucleic acid of claim 3 or 4 is stably integrated into the genome of said plant cell.
21. A transgenic plant or plant tissue comprising plant cells of claim 19 or 20.
22. A harvestable part of a plant of claim 21.
23. The harvestable part of a plant of claim 22 which is selected from the group consisting of seeds, leaves, fruits, stem cultures, rhizomes, roots, tubers and bulbs.
24. The progeny derived from any of the plants or plant parts of any of claims 21 to 23.
25. A method for stimulating root growth comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

26. A method for enhancing the formation of lateral or adventitious roots comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.
- 5 27. A method for altering root geotropism comprising altering the expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.
28. A method of any of claims 25 to 27, said method leading to an increase in yield.
29. The method of any of claims 25 to 28 wherein said expression of said nucleic acid
10 occurs under the control of a strong constitutive promoter.
30. The method of any of claims 25 to 28 wherein said expression of said nucleic acid occurs under the control of a promoter that is preferentially expressed in roots.
31. A method for identifying and obtaining proteins interacting with a polypeptide of claim 11 or 12 comprising a screening assay wherein a polypeptide of claim 11 or 12 is
15 used.
32. The method of claim 31 comprising a two-hybrid screening assay wherein a polypeptide of claim 11 or 12 as a bait and a cDNA library as prey are used.
33. A method for modulating the interaction between a polypeptide of claim 11 or 12 and interacting protein partners obtainable by a method according to claim 31 or 32.
- 20 34. A method for identifying and obtaining compounds interacting with a polypeptide of claim 11 or 12 comprising the steps of:
- a) providing a two-hybrid system wherein a polypeptide of claim 11 or 12 and an interacting protein partner obtainable by a method according to claim 31 or 32 are expressed,
 - 25 b) interacting said compound with the complex formed by the expressed polypeptides as defined in (a), and,
 - c) performing measurement of interaction of said compound with said polypeptide or the complex formed by the expressed polypeptides as defined in (a).
35. A method for identifying compounds or mixtures of compounds which specifically
30 bind to a polypeptide of claim 11 or 12, comprising:

a) combining a polypeptide of claim 11 or 12 with said compound or mixtures of compounds under conditions suitable to allow complex formation, and,

b) detecting complex formation, wherein the presence of a complex identifies a compound or mixture which specifically binds said polypeptide.

5 36. A method of any of claims 31 to 35 wherein said compound or mixture inhibits the activity of said polypeptide of claim 11 or 12 and can be used for the rational design of chemicals.

37. Use of a compound or mixture identified by means of a method of any of claims 31 to 35 as a plant growth regulator or herbicide.

10 38. A method for production of a plant growth regulator or herbicide composition comprising the steps of the method of any of claims 31 to 35 and formulating the compounds obtained from said steps in a suitable form for the application in agriculture or plant cell or tissue culture.

39. A method for the design of or screening for growth-promoting chemicals or
15 herbicides comprising the use of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or a vector of claim 7 or 8.

40. Use of a nucleic acid molecule of claim 3 or 4 or a nucleic acid as defined in claim 2, the vector of claim 7 or 8, a polypeptide of claim 11 or 12 for increasing yield.

41. Use of a nucleic acid molecule of claim 3 or 4 or a nucleic acid as defined in claim 2,
20 the vector of claim 7 or 8, a polypeptide of claim 11 or 12 for stimulating root growth.

42. Use of a nucleic acid molecule of claim claim 3 or 4 or a nucleic acid as defined in claim 2, the vector of claim 7 or 8, a polypeptide of claim 11 or 12 for enhancing the formation of lateral or adventitious roots.

43. Use of a nucleic acid molecule of claim claim 3 or 4 or a nucleic acid as defined in
25 claim 2, the vector of claim 7 or 8, a polypeptide of claim 11 or 12 for altering root geotropism.

44. Diagnostic composition comprising at least a nucleic acid molecule of any of claims 3 to 6, the vector of claim 7 or 8, a polypeptide of claim 11 or 12 or an antibody of claim 14.

45. A method for increasing the size of the root meristem comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression of a nucleic acid encoding a protein that reduces the level of active cytokinins in plants or plant parts, preferably in roots.
- 5 46. A method for increasing the root size comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression of another nucleic acid encoding a protein that reduces the level of active cytokinins in plants or plant parts, preferably in roots.
- 10 47. A method for increasing the size of the shoot meristem comprising downregulation of expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2, preferably in shoots.
48. A method for delaying leaf senescence comprising downregulation of expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 in leaves, preferably in senescing leaves.
- 15 49. A method for altering leaf senescence comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 in senescing leaves.
50. A method for increasing leaf thickness comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression of a nucleic acid encoding a protein that reduces the level of active cytokinins in plants or
20 plant parts.
51. A method for reducing the vessel size comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression of a nucleic acid encoding a protein that reduces the level of active cytokinins in plants or plant parts.
- 25 52. A method for increasing the vessel size comprising downregulation of expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 in plants or plant parts.
53. A method for inducing parthenocarpy comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression of a

nucleic acid encoding a protein that reduces the level of active cytokinins in plants or plant parts, preferably in the placenta, ovules and tissues derived therefrom.

54. A method for improving standability of seedlings comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 or comprising expression
5 of a nucleic acid encoding a protein that reduces the level of active cytokinins in seedlings, preferably in the roots of seedlings.
55. A method for increasing branching comprising expression of a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2 in plants or plant parts.
56. A method for improving lodging resistance comprising expression of a nucleic acid of
10 claim 3 or 4 or a nucleic acid as defined in claim 2 in plants or plant parts, preferably in stems or axillary buds.
57. Use of a transgenic rootstock in grafting procedures with a scion for improving the root-related characteristics of the resulting plant or tree characterized in an enhanced root growth due to expression of a plant cytokinin oxidase in said rootstock.
- 15 58. Use according to claim 57 wherein said plant cytokinin oxidase is encoded by a nucleic acid of claim 3 or 4 or a nucleic acid as defined in claim 2.
59. A transgenic plant comprising a transgenic rootstock expressing a plant cytokinin oxidase according to claim 57 or 58 and further comprising a scion.
60. A harvestable part of a plant of claim 59.
- 20 61. A method for stimulating root growth and development comprising expression of a nucleic acid encoding a plant cytokinin oxidase in a transgenic plant cell or tissue culture.
62. A method according to claim 61 wherein said nucleic acid is at least one of the nucleic acids of claim 3 or as defined in claim 2.




FIGURE 1

2/12

ZmCKX1	1	1	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
AtCKX1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
AtCKX2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
AtCKX3	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
AtCKX4	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
ZmCKX1	82	101	110	120	130	140	150	160	170	180	190	200										
AtCKX1	101	110	120	130	140	150	160	170	180	190	200											
AtCKX2	70	80	90	100	110	120	130	140	150	160	170	180	190	200								
AtCKX3	83	93	103	113	123	133	143	153	163	173	183	193	203									
AtCKX4	77	87	97	107	117	127	137	147	157	167	177	187	197	207								
ZmCKX1	173	201	210	220	230	240	250	260	270	280	290	300										
AtCKX1	190	210	220	230	240	250	260	270	280	290	300											
AtCKX2	154	164	174	184	194	204	214	224	234	244	254	264	274	284	294	304						
AtCKX3	171	181	191	201	211	221	231	241	251	261	271	281	291	301								
AtCKX4	177	187	197	207	217	227	237	247	257	267	277	287	297	307								
ZmCKX1	273	301	310	320	330	340	350	360	370	380	390	400										
AtCKX1	290	310	320	330	340	350	360	370	380	390	400											
AtCKX2	254	264	274	284	294	304	314	324	334	344	354	364	374	384	394	404						
AtCKX3	271	281	291	301	311	321	331	341	351	361	371	381	391	401								
AtCKX4	277	287	297	307	317	327	337	347	357	367	377	387	397	407								
ZmCKX1	373	401	410	420	430	440	450	460	470	480	490	500										
AtCKX1	377	407	417	427	437	447	457	467	477	487	497	507										
AtCKX2	340	350	360	370	380	390	400	410	420	430	440	450	460	470	480	490	500					
AtCKX3	358	368	378	388	398	408	418	428	438	448	458	468	478	488	498	508						
AtCKX4	363	373	383	393	403	413	423	433	443	453	463	473	483	493	503							
ZmCKX1	466	501	510	520	530	540	550	560	570	580	590	600										
AtCKX1	473	503	513	523	533	543	553	563	573	583	593	603										
AtCKX2	435	445	455	465	475	485	495	505	515	525	535	545	555	565	575	585	595	605				
AtCKX3	451	461	471	481	491	501	511	521	531	541	551	561	571	581	591	601						
AtCKX4	457	467	477	487	497	507	517	527	537	547	557	567	577	587	597	607						

FIGURE 2

3/12




Figure 3

4/12







Figure 4

5/12




Figure 5

6/12




Figure 6

7/12




Figure 7


8/12




A



B



C



D

Figure 8

9/12



Figure 9

10/12




Figure 10

11/12




Figure 11


12/12



A



B



C



D

Figure 12

SEQUENCE LISTING

<110> Schmülling, Thomas
Werner, Tomás

<120> Method for modifying plant morphology, biochemistry and
physiology

<130> CROP-005-PCT

<140>

<141>

<150> EP 00870132.8

<151> 2000-06-16

<150> US 60/258,415

<151> 2000-12-27

<150> EP 01870053.4

<151> 2001-03-16

<160> 36

<170> PatentIn Ver. 2.1

<210> 1

<211> 2236

<212> DNA

<213> Arabidopsis thaliana

<400> 1

```
atgggattga cctcaccctt acgggtccat agacaaaaca acaagacttt cctcggaatc 60
ttcatgatct tagttctaag ctgtatacca ggtagaacca atctttgttc caatcattct 120
35 gttagtaccc caaaagaatt accttcttca aatccttcag atattcggtc ctcattagtt 180
tcactagatt tggagggtta tataagcttc gacgatgtcc acaatgtggc caaggacttt 240
ggcaacagat accagttacc acctttggca attctacatc caaggtcagt ttttgatatt 300
tcacgatga tgaagcatat agtacatctg ggctccacct caaatcttac agtagcagct 360
agaggccatg gtcactcgct tcaaggacaa gctctagctc atcaagggtg tgtcatcaaa 420
40 atggagtcac ttcgaagtcc tgatatcagg atttataagg ggaagcaacc atatgttgat 480
gtctcagggtg gtgaaatatg gataaacatt ctacgcgaga ctctaaaata cggctcttca 540
ccaaagtcct ggacagacta ccttcatttg accggtggag gtacactatc taatgctgga 600
atcagcggtc aagcattcaa gcatggaccc caaatcaaca acgtctacca gctagagatt 660
gttacaggta tttcattcat gctttatctc tgcggtagtc tcaaaaaaat atgcacctgt 720
45 aaagaatatc catctcttca tgagcaaaaa cactgacgac tttaaataat ttttgactat 780
aaaacaagag tgcataaggca caaatgtgaa atatgcaaca cacaattgta acttgcacca 840
agaaaaaagt tataaaaaa aacaactgat aagcaatata tttccaatat ttaatcaggg 900
aaaggagaag tcgtaacctg ttctgagaag cgggaattctg aacttttctt cagtgttctt 960
ggcgggcttg gacagtttgg cataatcacc cgggcacgga tctctcttga accagcaccg 1020
50 catatggtaa agttctatct tgaacaaagt tcaacaata tacgctatga ttctaagaac 1080
cactttcctg acacagtcaa ataactttta atagggttaa tggatcaggg tactctactc 1140
tgacttttct gcattttcaa gggaccaaga atatctgatt tcgaaggaga aaacttttga 1200
ttacgttgaa ggatttgtga taatcaatag aacagacctt ctcaataatt ggcgatcgct 1260
attcagtcce aacgattcca cacaggcaag cagattcaag tcagatggga aaactcttta 1320
55 ttgcctagaa gtggtcaaat atttcaacct agaagaagct agctctatgg atcaggtaag 1380
```

atgtgaaagc aatatataac tagacttagt ttccacagag agctccaaat caaccgttgg 1440
 ctactagcct actaacataa tgaatggtg ccgtgcagga aactggcaag ttactttcag 1500
 agttaaatta tattccatcc actttgtttt catctgaagt gccatatatc gagtttctgg 1560
 atcgcgtagc tatcgagag agaaaactaa gagcaaagg tttatgggag gttccacatc 1620
 5 cctggctgaa tctcctgatt cctaagagca gcatatacca atttgctaca gaagttttca 1680
 acaacattct cacaagcaac aacaacggtc ctatccttat ttatccagtc aatcaatcca 1740
 agtaagttag caaatgccca aaagcaaatg cgtccagtga ttctgaaaca taaattacta 1800
 accatatcca acattttgtg gtttcagggtg gaagaaacat acatccttga taactccaaa 1860
 tgaagatata ttctatctcg tagcctttct cccctctgca gtgccaaatt cctcagggaa 1920
 10 aaacgatcta gagtaccttt tgaacaaaaa ccaaagagtt atgaacttct gcgcagcagc 1980
 aaacctcaac gtgaagcagt atttgcccca ttatgaaact caaaaagagt ggaaatcaca 2040
 ctttggtcaaa agatgggaaa catttgcaaa gaggaacaa gcctacgacc ctctagcgat 2100
 tctagcacct ggccaaagaa tattccaaaa gacaacagga aaattatctc ccatccaact 2160
 cgcaaagtca aaggcaacag gaagtctca aaggtaccat tacgcatcaa tactgccgaa 2220
 15 acctagaact gtataa 2236

<210> 2

<211> 575

20 <212> PRT

<213> Arabidopsis thaliana

<400> 2

Met Gly Leu Thr Ser Ser Leu Arg Phe His Arg Gln Asn Asn Lys Thr
 25 1 5 10 15
 Phe Leu Gly Ile Phe Met Ile Leu Val Leu Ser Cys Ile Pro Gly Arg
 20 25 30
 30 Thr Asn Leu Cys Ser Asn His Ser Val Ser Thr Pro Lys Glu Leu Pro
 35 35 40 45
 Ser Ser Asn Pro Ser Asp Ile Arg Ser Ser Leu Val Ser Leu Asp Leu
 50 55 60
 35 Glu Gly Tyr Ile Ser Phe Asp Asp Val His Asn Val Ala Lys Asp Phe
 65 70 75 80
 40 Gly Asn Arg Tyr Gln Leu Pro Pro Leu Ala Ile Leu His Pro Arg Ser
 85 90 95
 Val Phe Asp Ile Ser Ser Met Met Lys His Ile Val His Leu Gly Ser
 100 105 110
 45 Thr Ser Asn Leu Thr Val Ala Ala Arg Gly His Gly His Ser Leu Gln
 115 120 125
 Gly Gln Ala Leu Ala His Gln Gly Val Val Ile Lys Met Glu Ser Leu
 130 135 140
 50 Arg Ser Pro Asp Ile Arg Ile Tyr Lys Gly Lys Gln Pro Tyr Val Asp
 145 150 155 160
 55 Val Ser Gly Gly Glu Ile Trp Ile Asn Ile Leu Arg Glu Thr Leu Lys
 165 170 175

Tyr Gly Leu Ser Pro Lys Ser Trp Thr Asp Tyr Leu His Leu Thr Val
 180 185 190
 5 Gly Gly Thr Leu Ser Asn Ala Gly Ile Ser Gly Gln Ala Phe Lys His
 195 200 205
 Gly Pro Gln Ile Asn Asn Val Tyr Gln Leu Glu Ile Val Thr Gly Lys
 210 215 220
 10 Gly Glu Val Val Thr Cys Ser Glu Lys Arg Asn Ser Glu Leu Phe Phe
 225 230 235 240
 Ser Val Leu Gly Gly Leu Gly Gln Phe Gly Ile Ile Thr Arg Ala Arg
 245 250 255
 15 Ile Ser Leu Glu Pro Ala Pro His Met Val Lys Trp Ile Arg Val Leu
 260 265 270
 Tyr Ser Asp Phe Ser Ala Phe Ser Arg Asp Gln Glu Tyr Leu Ile Ser
 275 280 285
 Lys Glu Lys Thr Phe Asp Tyr Val Glu Gly Phe Val Ile Ile Asn Arg
 290 295 300
 25 Thr Asp Leu Leu Asn Asn Trp Arg Ser Ser Phe Ser Pro Asn Asp Ser
 305 310 315 320
 Thr Gln Ala Ser Arg Phe Lys Ser Asp Gly Lys Thr Leu Tyr Cys Leu
 325 330 335
 30 Glu Val Val Lys Tyr Phe Asn Pro Glu Glu Ala Ser Ser Met Asp Gln
 340 345 350
 Glu Thr Gly Lys Leu Leu Ser Glu Leu Asn Tyr Ile Pro Ser Thr Leu
 355 360 365
 35 Phe Ser Ser Glu Val Pro Tyr Ile Glu Phe Leu Asp Arg Val His Ile
 370 375 380
 Ala Glu Arg Lys Leu Arg Ala Lys Gly Leu Trp Glu Val Pro His Pro
 385 390 395 400
 40 Trp Leu Asn Leu Leu Ile Pro Lys Ser Ser Ile Tyr Gln Phe Ala Thr
 405 410 415
 Glu Val Phe Asn Asn Ile Leu Thr Ser Asn Asn Asn Gly Pro Ile Leu
 420 425 430
 45 Ile Tyr Pro Val Asn Gln Ser Lys Trp Lys Lys His Thr Ser Leu Ile
 435 440 445
 Thr Pro Asn Glu Asp Ile Phe Tyr Leu Val Ala Phe Leu Pro Ser Ala
 450 455 460
 55

	Val	Pro	Asn	Ser	Ser	Gly	Lys	Asn	Asp	Leu	Glu	Tyr	Leu	Leu	Lys	Gln	
	465					470					475					480	
5	Asn	Gln	Arg	Val	Met	Asn	Phe	Cys	Ala	Ala	Ala	Asn	Leu	Asn	Val	Lys	
				485						490						495	
	Gln	Tyr	Leu	Pro	His	Tyr	Glu	Thr	Gln	Lys	Glu	Trp	Lys	Ser	His	Phe	
				500					505					510			
10	Gly	Lys	Arg	Trp	Glu	Thr	Phe	Ala	Gln	Arg	Lys	Gln	Ala	Tyr	Asp	Pro	
			515					520					525				
	Leu	Ala	Ile	Leu	Ala	Pro	Gly	Gln	Arg	Ile	Phe	Gln	Lys	Thr	Thr	Gly	
		530					535					540					
15	Lys	Leu	Ser	Pro	Ile	Gln	Leu	Ala	Lys	Ser	Lys	Ala	Thr	Gly	Ser	Pro	
	545					550					555					560	
	Gln	Arg	Tyr	His	Tyr	Ala	Ser	Ile	Leu	Pro	Lys	Pro	Arg	Thr	Val		
20				565						570					575		
	<210> 3																
25	<211> 2991																
	<212> DNA																
	<213> Arabidopsis thaliana																
	<400> 3																
30	atgggcta	aatc	ttcg	ttta	aat	gatc	acttta	atcac	gggttt	taat	gatcac	caaat	catca	60			
	aacgg	tatta	aaatt	gattt	accta	aatcc	ctta	acctca	ccct	ctctac	cgatc	cttcc	120				
	atcat	ctccg	cagcc	ctctca	tgact	tcgga	aacata	aacca	ccgt	gacccc	cggcg	gcgta	180				
	atctg	ccccct	cctcc	accgc	tgat	atctct	cgtct	cctcc	aata	cgcgc	aaacg	gaaaa	240				
	agta	cattcc	aagta	gcggc	tcgt	ggccaa	ggcc	actcct	taaa	cggcca	agcct	cggtc	300				
35	tccgg	cgagg	taat	cgtaa	catga	cggtg	atcact	gacg	tggt	gggttc	aaaag	acaag	360				
	aagta	cgctg	acgt	ggcggc	cggg	acgtta	tggt	gggatg	tgct	taagaa	gacgg	cgagg	420				
	aaagg	gggtg	cgcc	gggttc	ttgg	acggat	tatt	tgata	taacc	gcgcg	agga	acgttg	480				
	tcga	atgggtg	gaatt	gggtg	tcaag	tgttt	cga	aacggtc	ctct	tgtag	taac	gtcctt	540				
	gaatt	ggacg	ttatt	actgg	tacgc	atctt	ctaa	actttg	atgt	acatac	aaca	acaaaa	600				
40	actg	ttttt	tttt	atagta	tttt	tcattt	tttg	taccat	aggt	tttatg	tttt	atagtt	660				
	gtg	ctaaact	tctt	gcacca	cacg	taagtc	ttcg	aaacac	aaa	atgcgta	acgc	atctat	720				
	atgt	ttttt	tac	atattga	atgt	tgttca	tgag	aaataa	agta	aattaca	tata	cacaca	780				
	ttt	attgtcg	tac	atatata	aata	aattaaa	gaca	aatttt	caca	attggg	agcg	tgtaa	840				
	ttt	gggattt	ttg	taatgta	catg	catgac	gc	atgcata	ggag	cttttc	gg	ttttctta	900				
45	gatt	tgtgta	gtatt	ttcaaa	tatat	cathtt	at	tttctttc	gaata	aaagag	gtgg	tatat	960				
	ttta	aaatag	caac	atttca	gaatt	ttttct	ttga	atttac	act	tttttaa	ttgt	tattgt	1020				
	taata	tggat	tttga	ataaaa	taatt	tcagg	gaa	aggtgaa	atgt	tgacat	gctc	gcgaca	1080				
	gcta	aacc	gaatt	gttct	atgg	agtggt	agg	aggttg	ggt	caatttg	gaatt	tataac	1140				
	gag	agccaga	attg	ttttg	accat	gcacc	taaa	cgggta	cgt	atcatca	tatt	tttacc	1200				
50	ttt	gttttag	tcag	cattca	tttt	tcatta	gta	attccgt	ttca	atttct	aaatt	ttttt	1260				
	agt	caataga	aaat	gattct	tatg	tcagag	ctt	gattatt	tagt	gatttt	tatt	gagata	1320				
	aaata	aaaata	taac	cta	acg	gaa	ataatta	tttt	actaat	cgg	ataatgt	ctg	attaaaa	1380			
	cat	tttatga	tatt	acacta	agag	agttag	agac	gatg	atca	caaaac	atga	agcttt	1440				
	cttag	atgg	atccta	aaaac	taa	agttag	taca	agtttg	gaatt	taggt	caaat	gctta	1500				
55	agtt	gcatta	attt	gaacaa	aat	ctatgca	ttga	ataaaa	aaa	agatatg	gatt	atttta	1560				


```

    taaagtatag tccttgtaat cctaggactt gttgtctaatt cttgtcttat gcgtgcaaat 1620
    ctttttgatg tcaatatata atccttggtt attagagtca agctctttca ttagtcaact 1680
    actcaaatat actccaaagt ttagaatata gtcttctgac taattagaat cttacaaccg 1740
    ataaacggtta caatttggtt atcatttttaaaaacagatt tggtcataat atacgatgac 1800
5   gttctgtttt agtttcatct attcacaat tttatataat ttttttcaag aaaatattga 1860
    aatactatac tgtaatatgg tttctttata tatgtgtgta taaattaaat gggattggtt 1920
    tctctaaatg aaattgtgta ggccaaatgg tttcggatgc tctacagtga tttcacaact 1980
    tttcacaagg accaagaacg tttgatatca atggcaaacg atattggagt cgactattta 2040
    gaagggtcaaa tttttctatc aaacggtgtc gttgacacct cttttttccc accttcagat 2100
10  caatctaaag tcgctgatct agtcaagcaa cacggtatca tctatgttct tgaagtagcc 2160
    aagtattatg atgatcccaa tctccccatc atcagcaagg tactacacat ttacattttc 2220
    atcatcgttt ttatcatacc ataagatatt taaatgattc atcattgcac cacattaaga 2280
    tattcatcat catcatcggt acattttttt ttgcatctta tgcttctcat aatctactat 2340
    tgtgtagggtt attgacacat taacgaaaac attaagttac ttgcccgggt tcatatcaat 2400
15  gcacgacgtg gcctacttcg atttcttgaa ccgtgtacat gtcgaagaaa ataaactcag 2460
    atctttggga ttatgggaac ttctcatcc ttggcttaac ctctacgttc ctaaactctc 2520
    gattctcgat tttcataacg gtgtgtgcaa agacattctt ctttaagcaaa aatcagcttc 2580
    gggactcgct cttctctatc caacaaaccg gaataagtac atacttctct tcattcatat 2640
    ttatcttcaa gaaccaaatg aaataaattt ctatgaactg attatgctgt tattgttaga 2700
20  tgggacaatc gtatgtcggc gatgatacca gagatcgatg aagatgttat atatattatc 2760
    ggactactac aatccgctac cccaaaggat cttccagaag tggagagcgt taacgagaag 2820
    ataattaggt tttgcaagga ttcagggtatt aagattaagc aatatctaata gcattatact 2880
    agtaaagaag attggattga gcatttttga tcaaaatggg atgatttttc gaagaggaaa 2940
    gatctatttg atcccaagaa actgttatct ccagggcaag acatcttttg a 2991
25

```

<210> 4

<211> 501

<212> PRT

30 <213> Arabidopsis thaliana

<400> 4

```

Met Ala Asn Leu Arg Leu Met Ile Thr Leu Ile Thr Val Leu Met Ile
  1           5           10           15
35 Thr Lys Ser Ser Asn Gly Ile Lys Ile Asp Leu Pro Lys Ser Leu Asn
    20           25           30
40 Leu Thr Leu Ser Thr Asp Pro Ser Ile Ile Ser Ala Ala Ser His Asp
    35           40           45
    Phe Gly Asn Ile Thr Thr Val Thr Pro Gly Gly Val Ile Cys Pro Ser
    50           55           60
45 Ser Thr Ala Asp Ile Ser Arg Leu Leu Gln Tyr Ala Ala Asn Gly Lys
    65           70           75           80
    Ser Thr Phe Gln Val Ala Ala Arg Gly Gln Gly His Ser Leu Asn Gly
    85           90           95
50 Gln Ala Ser Val Ser Gly Gly Val Ile Val Asn Met Thr Cys Ile Thr
    100          105          110
55 Asp Val Val Val Ser Lys Asp Lys Lys Tyr Ala Asp Val Ala Ala Gly
    115          120          125

```

Thr Leu Trp Val Asp Val Leu Lys Lys Thr Ala Glu Lys Gly Val Ser
 130 135 140
 5 Pro Val Ser Trp Thr Asp Tyr Leu His Ile Thr Val Gly Gly Thr Leu
 145 150 155 160
 Ser Asn Gly Gly Ile Gly Gly Gln Val Phe Arg Asn Gly Pro Leu Val
 165 170 175
 10 Ser Asn Val Leu Glu Leu Asp Val Ile Thr Gly Lys Gly Glu Met Leu
 180 185 190
 15 Thr Cys Ser Arg Gln Leu Asn Pro Glu Leu Phe Tyr Gly Val Leu Gly
 195 200 205
 Gly Leu Gly Gln Phe Gly Ile Ile Thr Arg Ala Arg Ile Val Leu Asp
 210 215 220
 20 His Ala Pro Lys Arg Ala Lys Trp Phe Arg Met Leu Tyr Ser Asp Phe
 225 230 235 240
 Thr Thr Phe Thr Lys Asp Gln Glu Arg Leu Ile Ser Met Ala Asn Asp
 245 250 255
 25 Ile Gly Val Asp Tyr Leu Glu Gly Gln Ile Phe Leu Ser Asn Gly Val
 260 265 270
 30 Val Asp Thr Ser Phe Phe Pro Pro Ser Asp Gln Ser Lys Val Ala Asp
 275 280 285
 Leu Val Lys Gln His Gly Ile Ile Tyr Val Leu Glu Val Ala Lys Tyr
 290 295 300
 35 Tyr Asp Asp Pro Asn Leu Pro Ile Ile Ser Lys Val Ile Asp Thr Leu
 305 310 315 320
 Thr Lys Thr Leu Ser Tyr Leu Pro Gly Phe Ile Ser Met His Asp Val
 325 330 335
 40 Ala Tyr Phe Asp Phe Leu Asn Arg Val His Val Glu Glu Asn Lys Leu
 340 345 350
 45 Arg Ser Leu Gly Leu Trp Glu Leu Pro His Pro Trp Leu Asn Leu Tyr
 355 360 365
 Val Pro Lys Ser Arg Ile Leu Asp Phe His Asn Gly Val Val Lys Asp
 370 375 380
 50 Ile Leu Leu Lys Gln Lys Ser Ala Ser Gly Leu Ala Leu Leu Tyr Pro
 385 390 395 400
 Thr Asn Arg Asn Lys Trp Asp Asn Arg Met Ser Ala Met Ile Pro Glu
 405 410 415
 55

Ile Asp Glu Asp Val Ile Tyr Ile Ile Gly Leu Leu Gln Ser Ala Thr
420 425 430

Pro Lys Asp Leu Pro Glu Val Glu Ser Val Asn Glu Lys Ile Ile Arg
5 435 440 445

Phe Cys Lys Asp Ser Gly Ile Lys Ile Lys Gln Tyr Leu Met His Tyr
450 455 460

10 Thr Ser Lys Glu Asp Trp Ile Glu His Phe Gly Ser Lys Trp Asp Asp
465 470 475 480

Phe Ser Lys Arg Lys Asp Leu Phe Asp Pro Lys Lys Leu Leu Ser Pro
15 485 490 495

15 Gly Gln Asp Ile Phe
500

20

<210> 5
<211> 3302
<212> DNA
<213> Arabidopsis thaliana

25

<400> 5
atggcgagtt ataactcttcg ttcacaagtt cgtcttatag caataacaat agtaatcatc 60
attactctct caactccgat cacaaccaac acatcaccac aaccatggaa taccctttca 120
cacaacgaat tcgccggaaa actcacctcc tcctcctcct ccgtcgaatc agccgccaca 180
30 gatttcggcc acgtcaccaa aatcttccct tccgccgtct taatcccttc ctccgttgaa 240
gacatcacag atctcataaa actctctttt gactctcaac tgtcttttcc tttagccgct 300
cgtgggtcacg gacacagcca ccgtggccaa gcctcggcta aagacggagt tgtgggtcaac 360
atgcggtcca tggtaaaccg ggatcgaggt atcaagggtg ctaggacctg tttatatgtt 420
gacgtggacg ctgctgtggct atggattgag gtgttgaata aaactttgga gttagggtta 480
35 acgccggttt cttggacgga ttatttgtat ttaacagtcg gtgggacgtt atcaaacggc 540
ggaattagtg gacaaacgtt tcggtacggt ccacagatca ctaatgttct agagatggat 600
gttattactg gtacgtacca cgatcttttt cacacagaga ttaaaaaaaaa cagtaatagt 660
gattttaact tcgtacgttt ctgatagaca acaagaact tcgtacgttt ttcgaagttt 720
tttcgtcttt ttcatttttag atctgcgcgg ccatttttgg ttatgctatt gtttgtttgt 780
40 attgtttgtc tctgtttatt tatttctcga acttgttgat agcttttctt cttttcacac 840
atcaatctaa tcaccttttt tggctttaag attagaaaga agatacggac taggtaaaaa 900
taggtgggtg taaacgtaga cgcattaaaa aaatattggt ttttttattt tttgataagc 960
aaaattggtg gttggtctaa gattataaac ttgatattaa tgcaaaggtc gatctagcaa 1020
tagaagatta atcaatatc ttgtgtttt aacaacagat tatttcatca ttaaaatcgt 1080
45 gaaacaaaga aattttggta gtatacatc cgtgtagttt tgttagttta ttaaaaaaaa 1140
tagtatatag ttttgttaaa acgcgattta tttagtaaca cattagtata ttacacgttt 1200
aaccaactaa actttttttt ttgaataatt atgttctata tttcttactc aaattatgca 1260
aatttcgtgg attcgaagtc aaatttctgc gaaatttaca tgggtcatata ttataaaact 1320
gttcatataa cccggtgaac aaacagacaa ttaagggttt gaatgggtac ggcggttggg 1380
50 gcggacacaa ccgtcaatag atcagaccgt tttttattta ccattcatca attatattcc 1440
gcagtgggtt ggggtaaaaa aaatagaaga aaaccgcagc ggaccaattc cataccgttt 1500
ttacatacaa ataaacatgg tgcgaacgg tttattgtcc gcctcaaaaa tgaaatggac 1560
taaaccgcag ataaattaga ccgctttgtc cgtgcctcc attcatagac taaaaaaaaa 1620
caacaaaaaa aaaaatggtc ccacgccccat gattttacac gaggtttctt gtggcgtaag 1680
55 gacaaaactc aaaagttcac aacgtttggt cctaaccagg tgtaatggat taagtaacag 1740

tcaatcttct tattataget gtatccatta tgtccacata tgcattccata tacattacac 1800
 tgttgggtctc aagtgttagtt agattacgaa gactttcaag ttccattttt tggttaggag 1860
 ataaacataa tttaatgata ccgacttttag cactctaggc tcaaaacaag tacagaagag 1920
 aatagtttta tttcaaaactc gttgcattgt tgtatcaatt aattgtgtta gtctttgtat 1980
 5 attcttacat aacgggtccaa gtttgttgaa atagtttact tactaaactt ttcctaattg 2040
 ggtcaaatctt tatttttatag gaaaaggaga gattgcaact tgttccaagg acatgaactc 2100
 ggatcttttc ttccgcggtgt taggagggtt ggttcaattc ggcattataa caagagccag 2160
 aattaaactt gaagtagctc cgaaaagggt atgttaaatt tgtaaattat gcaactacag 2220
 aaaattctat gaaatttatg aatgaacata tatgcatttt tggatttttg taggccaagt 2280
 10 ggttaagggt tctatacata gatttctccg aattcacaag agatcaagaa cgagtgatat 2340
 cgaaaacgga cggtgtagat ttcttagaag gttccattat ggtgggacct ggcccaccgg 2400
 ataactggag atccacgtat tatccaccgt ccgatcactt gaggatcgcc tcaatggtca 2460
 aacgacatcg tgtcatctac tgccttgaag tcgtcaagta ttacgacgaa acttctcaat 2520
 acacagtcaa cgagggtccgt acatacatat aatcataaat catacatgta taattgggag 2580
 15 atctttatgc attattcaat tatattaatt tacttttagtt atttaactta tgcaggaaat 2640
 ggaggagtta agcgatagtt taaaccatgt aagaggggtt atgtacgaga aagatgtgac 2700
 gtatatggat ttcttaaaacc gagttcgaac cggagagcta aacctgaaat ccaaaggcca 2760
 atgggatggt ccacatccat ggcttaattc ctctgtacca aaaactcaaa tctccaaatt 2820
 tgatgatggt gtttttaagg gtattatcct aagaaataac atcactagcg gtcctgttct 2880
 20 tgtttatcct atgaatcgca acaagtaagt ttaactcgat attgcaaaat ttactatcta 2940
 catcttcggt ttggaatccg aaatattctt acaagctaatt tttatgcggc gtttttaggt 3000
 ggaatgatcg gatgtctgcc gctatacccg aggaagatgt attttatgag gtaggggttt 3060
 taagatccgc ggggttttgac aattgggagg cttttgatca agaaaacatg gaaatactga 3120
 agttttgtga ggatgctaatt atgggggtta tacaatatct tccttatcat tcatcacaag 3180
 25 aaggatgggt tagacatttt ggtccgaggt ggaatatatt cgtagagaga aaatataaat 3240
 atgatcccaa aatgatatta tcaccgggac aaaatatatt tcaaaaaata aactcgagtt 3300
 ag 3302

30 <210> 6
 <211> 523
 <212> PRT
 <213> Arabidopsis thaliana

35 <400> 6
 Met Ala Ser Tyr Asn Leu Arg Ser Gln Val Arg Leu Ile Ala Ile Thr
 1 5 10 15
 Ile Val Ile Ile Ile Thr Leu Ser Thr Pro Ile Thr Thr Asn Thr Ser
 40 20 25 30
 Pro Gln Pro Trp Asn Ile Leu Ser His Asn Glu Phe Ala Gly Lys Leu
 35 40 45
 45 Thr Ser Ser Ser Ser Val Glu Ser Ala Ala Thr Asp Phe Gly His
 50 55 60
 Val Thr Lys Ile Phe Pro Ser Ala Val Leu Ile Pro Ser Ser Val Glu
 65 70 75 80
 50 Asp Ile Thr Asp Leu Ile Lys Leu Ser Phe Asp Ser Gln Leu Ser Phe
 85 90 95
 55 Pro Leu Ala Ala Arg Gly His Gly His Ser His Arg Gly Gln Ala Ser
 100 105 110

Ala Lys Asp Gly Val Val Val Asn Met Arg Ser Met Val Asn Arg Asp
 115 120 125
 5 Arg Gly Ile Lys Val Ser Arg Thr Cys Leu Tyr Val Asp Val Asp Ala
 130 135 140
 Ala Trp Leu Trp Ile Glu Val Leu Asn Lys Thr Leu Glu Leu Gly Leu
 145 150 155 160
 10 Thr Pro Val Ser Trp Thr Asp Tyr Leu Tyr Leu Thr Val Gly Gly Thr
 165 170 175
 15 Leu Ser Asn Gly Gly Ile Ser Gly Gln Thr Phe Arg Tyr Gly Pro Gln
 180 185 190
 Ile Thr Asn Val Leu Glu Met Asp Val Ile Thr Gly Lys Gly Glu Ile
 195 200 205
 20 Ala Thr Cys Ser Lys Asp Met Asn Ser Asp Leu Phe Phe Ala Val Leu
 210 215 220
 Gly Gly Leu Gly Gln Phe Gly Ile Ile Thr Arg Ala Arg Ile Lys Leu
 225 230 235 240
 25 Glu Val Ala Pro Lys Arg Ala Lys Trp Leu Arg Phe Leu Tyr Ile Asp
 245 250 255
 30 Phe Ser Glu Phe Thr Arg Asp Gln Glu Arg Val Ile Ser Lys Thr Asp
 260 265 270
 Gly Val Asp Phe Leu Glu Gly Ser Ile Met Val Asp His Gly Pro Pro
 275 280 285
 35 Asp Asn Trp Arg Ser Thr Tyr Tyr Pro Pro Ser Asp His Leu Arg Ile
 290 295 300
 Ala Ser Met Val Lys Arg His Arg Val Ile Tyr Cys Leu Glu Val Val
 305 310 315 320
 40 Lys Tyr Tyr Asp Glu Thr Ser Gln Tyr Thr Val Asn Glu Glu Met Glu
 325 330 335
 45 Glu Leu Ser Asp Ser Leu Asn His Val Arg Gly Phe Met Tyr Glu Lys
 340 345 350
 Asp Val Thr Tyr Met Asp Phe Leu Asn Arg Val Arg Thr Gly Glu Leu
 355 360 365
 50 Asn Leu Lys Ser Lys Gly Gln Trp Asp Val Pro His Pro Trp Leu Asn
 370 375 380
 Leu Phe Val Pro Lys Thr Gln Ile Ser Lys Phe Asp Asp Gly Val Phe
 385 390 395 400
 55

Lys Gly Ile Ile Leu Arg Asn Asn Ile Thr Ser Gly Pro Val Leu Val
 405 410 415
 Tyr Pro Met Asn Arg Asn Lys Trp Asn Asp Arg Met Ser Ala Ala Ile
 5 420 425 430
 Pro Glu Glu Asp Val Phe Tyr Ala Val Gly Phe Leu Arg Ser Ala Gly
 435 440 445
 10 Phe Asp Asn Trp Glu Ala Phe Asp Gln Glu Asn Met Glu Ile Leu Lys
 450 455 460
 Phe Cys Glu Asp Ala Asn Met Gly Val Ile Gln Tyr Leu Pro Tyr His
 15 465 470 475 480
 Ser Ser Gln Glu Gly Trp Val Arg His Phe Gly Pro Arg Trp Asn Ile
 485 490 495
 Phe Val Glu Arg Lys Tyr Lys Tyr Asp Pro Lys Met Ile Leu Ser Pro
 20 500 505 510
 Gly Gln Asn Ile Phe Gln Lys Ile Asn Ser Ser
 515 520
 25
 <210> 7
 <211> 2782
 <212> DNA
 30 <213> Arabidopsis thaliana
 <400> 7
 atgactaata ctctctgttt aagcctcatt accctaataa cgctttttat aagtttaacc 60
 ccaaccttaa tcaaatcaga tgagggcatt gatgttttct taccatattc actcaacctt 120
 35 acgggtcctaa ccgatccctt ctccatctct gccgcttctc acgacttcgg taacataacc 180
 gacgaaaatc ccggcgccgt cctctgccct tcctccacca cggagggtggc tcgtctcctc 240
 cgtttcgtca acggaggatt ctcttacaat aaaggetcaa ccagccccgc gtctactttc 300
 aaagtggctg ctcgaggcca agccactcc ctccgtggcc aagcctctgc acccggagggt 360
 gtcgtcgtga acatgacgtg tctcgccatg gcggctaaac cagcggcggt tgttatctcg 420
 40 gcagacggga cttacgtctga cgtggctgcc gggacgatgt ggggtggatgt tctgaaggcg 480
 gcgggtggata gaggcgtctc gccggttaca tggacggatt atttgtatct cagcgtcggc 540
 gggacgttgt cgaacgctgg aatcgggtgg cagacgttta gacacggccc tcagattagt 600
 aacgttcatg agcttgacgt tattaccggg acgtaaatac caaaacttca ctaatctcgt 660
 tacaattttt taattttttg gtaatatata ttttgtacgg ctcaactctt aattaagaat 720
 45 gaaacagtat ctatgatctt ctagatgctc tttttttgtc tgcaagcttt aattgtagta 780
 acatcagcga tatatatatc acatgcatgt gtattattga tgataatata taatgtttta 840
 gttacaaatt tgattctcaa ggtaaaactc acacgccata accagtataa aactccaaaa 900
 atcacgtttt ggtcagaaat acatattcctt cattaacagt agttatgcta taatttgtga 960
 ttataaataa ctccggaggt tgttcacaat actaaatttc aggaaaagggt gaaatgatga 1020
 50 cttgctctcc aaagttaaac cctgaattgt tctatggagt tttaggagggt ttgggtcaat 1080
 tcgggtattat aacgagggcc aggattgcgt tggatcatgc acccacaagg gtatgtatca 1140
 tgcattcata gtgtaataca ttataattt taatgtagtg gtcctaaatc caaaatttga 1200
 tttgatttgg ttggaacgta cgtatatata ataagtcaaa aggctgattt tgaagacgaa 1260
 tttatatact tttgttgaat taaatctgat tttgcttacg ttttattaga ttctgcgtaa 1320
 55 taaatcctag gacttgctcg agtgtaatct tgtcttatgc ttgcaaactc tgttgatgtc 1380

```

aatatccta atctttttt atatttccct acgtaagttt tagatatagt tatttttaaac 1440
tgctataaat tgtgtacgta tagacttttag ataaaaagtt gtgggtcgctt gcacctattt 1500
gtttatcgct atagtgatc aaaggtctat atatgattct tgggttttct ttttgaaaaa 1560
aatagaccat acaatccaag gaagatgatc ttaaatggac taatttatgg atataaattg 1620
5 atatacaaat ctgcagggtga aatgggtctcg cactactctac agtgacttct cggcttttaa 1680
aagagaccaa gagcgtttaa tatcaatgac caatgatctc ggagttgact ttttggaagg 1740
tcaacttatg atgtcaaag gcttcgtaga cactctcttc tccctactct ccgatcaaac 1800
aagagtcgca tctcttgtga atgaccaccg gatcatctat gttctcgaag tagccaagta 1860
ttatgacaga accacccttc ccattattga ccagggtacta aaatccatta ttcgatgatga 1920
10 ttatcttcac acaatcagta tcatcaccaa ttaccatcat cacttgatcat atatgatcca 1980
aagtaaatat atcacatgat ataaataaat cgttcaaate ttttttttta aagaataaaa 2040
gaatcatttt caagcattac tcatacacat ctacgaatca ccgtgaccat atataaccat 2100
acgcttatta aataatcatt tttgtttgta ggtgattgac acgttaagta gaactctagg 2160
tttcgcteca ggggtttatgt tcgtacaaga tggtccgtat ttcgatttct tgaaccgtgt 2220
15 ccgaaacgaa gaagataaac tcagatcttt aggactatgg gaagttcctc atccatggct 2280
taacatcttt gtcccggggg ctgcaatcca agattttcat gatgggtgta ttaatggcct 2340
tcttctaaac caaacctcaa cttctggtgt tactctcttc tatcccacaa accgaaacaa 2400
gtaaataattt actttttgat tttgttttat ttgaaagtat atcccaataa tgtatgttaa 2460
attgttaaca agaattttatt ttattaatag atggaacaac cgcagtgtcaa cgatgacacc 2520
20 ggacgaagat gttttttatg tgatcggatt actgcaatca gctgggtggat ctcaaaattg 2580
gcaagaactt gaaaatctca acgacaaggt tattcagttt tgtgaaaact cgggaattaa 2640
gattaaggaa tatttgatgc actatacaag aaaagaagat tgggttaaac attttggacc 2700
aaaatgggat gattttttta gaaagaaaat tatgtttgat cccaaaagac tattgtctcc 2760
aggacaagac atatttaatt aa 2782

```

<210> 8

<211> 524

<212> PRT

30 <213> Arabidopsis thaliana

<400> 8

```

Met Thr Asn Thr Leu Cys Leu Ser Leu Ile Thr Leu Ile Thr Leu Phe
  1             5             10             15

```

```

35 Ile Ser Leu Thr Pro Thr Leu Ile Lys Ser Asp Glu Gly Ile Asp Val
      20             25             30

```

```

40 Phe Leu Pro Ile Ser Leu Asn Leu Thr Val Leu Thr Asp Pro Phe Ser
      35             40             45

```

```

Ile Ser Ala Ala Ser His Asp Phe Gly Asn Ile Thr Asp Glu Asn Pro
  50             55             60

```

```

45 Gly Ala Val Leu Cys Pro Ser Ser Thr Thr Glu Val Ala Arg Leu Leu
  65             70             75             80

```

```

Arg Phe Ala Asn Gly Gly Phe Ser Tyr Asn Lys Gly Ser Thr Ser Pro
      85             90             95

```

```

50 Ala Ser Thr Phe Lys Val Ala Ala Arg Gly Gln Gly His Ser Leu Arg
      100            105            110

```

```

55 Gly Gln Ala Ser Ala Pro Gly Gly Val Val Val Asn Met Thr Cys Leu
      115            120            125

```

	Ala Met Ala Ala Lys Pro Ala Ala Val Val Ile Ser Ala Asp Gly Thr	
	130	135 140
5	Tyr Ala Asp Val Ala Ala Gly Thr Met Trp Val Asp Val Leu Lys Ala	
	145	150 155 160
	Ala Val Asp Arg Gly Val Ser Pro Val Thr Trp Thr Asp Tyr Leu Tyr	
		165 170 175
10	Leu Ser Val Gly Gly Thr Leu Ser Asn Ala Gly Ile Gly Gly Gln Thr	
		180 185 190
	Phe Arg His Gly Pro Gln Ile Ser Asn Val His Glu Leu Asp Val Ile	
15		195 200 205
	Thr Gly Lys Gly Glu Met Met Thr Cys Ser Pro Lys Leu Asn Pro Glu	
		210 215 220
20	Leu Phe Tyr Gly Val Leu Gly Gly Leu Gly Gln Phe Gly Ile Ile Thr	
		225 230 235 240
	Arg Ala Arg Ile Ala Leu Asp His Ala Pro Thr Arg Val Lys Trp Ser	
		245 250 255
25	Arg Ile Leu Tyr Ser Asp Phe Ser Ala Phe Lys Arg Asp Gln Glu Arg	
		260 265 270
	Leu Ile Ser Met Thr Asn Asp Leu Gly Val Asp Phe Leu Glu Gly Gln	
30		275 280 285
	Leu Met Met Ser Asn Gly Phe Val Asp Thr Ser Phe Phe Pro Leu Ser	
		290 295 300
35	Asp Gln Thr Arg Val Ala Ser Leu Val Asn Asp His Arg Ile Ile Tyr	
		305 310 315 320
	Val Leu Glu Val Ala Lys Tyr Tyr Asp Arg Thr Thr Leu Pro Ile Ile	
		325 330 335
40	Asp Gln Val Ile Asp Thr Leu Ser Arg Thr Leu Gly Phe Ala Pro Gly	
		340 345 350
	Phe Met Phe Val Gln Asp Val Pro Tyr Phe Asp Phe Leu Asn Arg Val	
45		355 360 365
	Arg Asn Glu Glu Asp Lys Leu Arg Ser Leu Gly Leu Trp Glu Val Pro	
		370 375 380
50	His Pro Trp Leu Asn Ile Phe Val Pro Gly Ser Arg Ile Gln Asp Phe	
		385 390 395 400
	His Asp Gly Val Ile Asn Gly Leu Leu Leu Asn Gln Thr Ser Thr Ser	
		405 410 415
55		

	Gly Val Thr	Leu Phe Tyr	Pro Thr	Asn Arg Asn	Lys Trp	Asn Asn Arg	
	420			425		430	
5	Met Ser Thr	Met Thr Pro	Asp Glu	Asp Val Phe	Tyr Val	Ile Gly Leu	
	435		440		445		
	Leu Gln Ser	Ala Gly Gly	Ser Gln	Asn Trp	Gln Glu	Leu Glu Asn	Leu
	450		455		460		
10	Asn Asp Lys	Val Ile Gln	Phe Cys	Glu Asn	Ser Gly	Ile Lys	Ile Lys
	465		470		475		480
	Glu Tyr Leu	Met His Tyr	Thr Arg	Lys Glu	Asp Trp	Val Lys	His Phe
		485		490			495
15	Gly Pro Lys	Trp Asp Asp	Phe Leu	Arg Lys	Lys Ile	Met Phe	Asp Pro
		500		505		510	
20	Lys Arg Leu	Leu Ser Pro	Gly Gln	Asp Ile	Phe Asn		
	515		520				
	<210> 9						
25	<211> 2805						
	<212> DNA						
	<213> Arabidopsis thaliana						
	<400> 9						
30	atgacgtcaa	gctttcttct	cctgacgttc	gccatatgta	aactgatcat	agccgtgggt	60
	ctaaacgtgg	gcccagtgga	gctcctccgc	atcggagcca	tagatgtcga	cggccacttc	120
	accgtccacc	cttccgactt	agcctccgtc	tcctcagact	tcggtatgct	gaagtcacct	180
	gaagagccat	tggccgtgct	tcatccatca	tcggccgaag	acgtggcacg	actcgtcaga	240
	acagcttacg	gttcagccac	ggcgtttccg	gtctcagccc	gaggccacgg	ccattccata	300
35	aacggacaag	ccgcggcggg	gaggaacggt	gtgggtgggt	aaatgaacca	cggcgtaacc	360
	gggacgcccc	agccactcgt	ccgaccggat	gaaatgtatg	tggatgtatg	gggtggagag	420
	ttatgggtcg	atgtgttgaa	gaaaacgttg	gagcatggct	tagcaccaaa	atcatggacg	480
	gattacttgt	atctaaccgt	tggagggtaca	ctctccaatg	caggaatcag	tgggtcaagct	540
	tttcaccatg	gtcctcaa	tagtaacgtc	cttgagctcg	acgttgtaac	tgggttagtat	600
40	taaaacattc	aagttcatat	attttaaatg	cttttgtctg	aagttttact	aataacaaga	660
	aattgatata	aaaaagtagg	gaaaggagag	gtgatgagat	gctcagaaga	agagaacaca	720
	aggctattcc	atggagtctt	tgggtggatta	ggtcaatttg	ggatcatcac	tcgagcacga	780
	atctctctcg	aaccagctcc	ccaaagggtg	atattttttt	aatgactagc	tatcaaaaat	840
	ccctggcggg	tccatacgtt	gtaatctttt	tagtttttac	tgttgatggg	attttttata	900
45	tattttggat	aataaaaccc	taaaatggta	tattgtgatg	acagggtgaga	tggatacggg	960
	tattgtattc	gagcttcaaa	gtgtttacgg	aggaccaaga	gtacttaatc	tcaatgcatg	1020
	gtcaattaaa	gtttgattac	gtggaagggt	ttgtgattgt	ggacgaagga	ctcgtcaaca	1080
	attggagatc	ttctttcttc	tctccacgta	acccgcgtaa	gatctcctct	gttagttcca	1140
	acggctctgt	tttgtattgc	cttgagatca	ccaagaacta	ccacgactcc	gactccgaaa	1200
50	tcgttgatca	ggtcactttc	attattcact	tagaaaaaag	cgatattttc	attttttata	1260
	ttgatgaata	tctggaagga	tttaacgcta	tgcgactatt	gggaaatcat	tatgaaaaaa	1320
	tatttagttt	atatgattga	aagtgtctc	catagtattt	ttgttggtgc	gactttatta	1380
	taacttaaat	ttggaagagg	acatgaagaa	gaagccagag	aggatctaca	gagatctagc	1440
	ttttccacct	gaacttaata	atgcacattt	atataattat	ttttcttctt	ctaaagttta	1500
55	gtttatcact	agcgaattaa	tcatggttac	taattaaagta	qtggacaggg	tcatggacca	1560

```

ctcactcacc aaataatgat tectcttttac tcttaagttt aattttaata aaaccaactc 1620
tactggaatc ttaacttata cttgggttttg gtaggctttt atagcaacac ggttttttta 1680
attttcctat tccagatttt gtatatataa tgtecgatttt ttttcttttt gtttcaggaa 1740
gttgagattc tgatgaagaa attgaatttc ataccgacat cggctctttac aacggattta 1800
5 caatatgtgg actttctcga ccgggtacac aaggccgaat tgaagctccg gtccaagaat 1860
ttatgggagg ttccacaccc atgggtcaac ctcttcgtgc caaaatcaag aatctctgac 1920
ttcgataaag gcgttttcaa gggcattttg ggaaataaaa caagtggccc tattcttata 1980
taccatga acaaagacaa gtaagtcttg acattaccat tgattactac ttctaaattt 2040
cttctctaga aaaaagaata aaacgagttt tgcattgcat gcatgcaaag ttacacttgt 2100
10 ggggattaat tagtggtcca agaaaaaag tttgtcaaaa ttgaaaaaaa ctagacacgt 2160
ggtacatggg attgtccgaa aaacggtgtc cacatgtgca tcgaaccagc taagattgac 2220
aacaacactt cgtcggctcg tatttctctt tttgttttgt gaccaaacc gatggtccag 2280
attgggttta ttgttttta agttcctaga actcatggtg ggtgggtccc aatcagattc 2340
tctgagacca aaccgatctc aacgaaccct ccgcacatca ttgattatta cattaatata 2400
15 gatattgtcg ttgctgacgt gtcgtaattt gatgttattg tcagatggga cgagaggagc 2460
tcagccgtga cgccggatga ggaagttttc tatctggtgg ctctattgag atcagcttta 2520
acggacggtg aagagacaca gaagctagag tatctgaaag atcagaaccg tcggatcttg 2580
gagttctgtg aacaagccaa gatcaatgtg aagcagtatc ttcctcacca cgcaacacag 2640
gaagagtggg tggctcattt tggggacaag tgggatcggt tcagaagctt aaaggctgag 2700
20 ttgatccgc gacacatact cgctactggt cagagaatct ttcaaaacc atctttgtct 2760
ttgtttctc cgctcgtcgc ttcttcgtca gcggcttcat ggtga 2805

```

<210> 10

25 <211> 536

<212> PRT

<213> Arabidopsis thaliana

<400> 10

```

30 Met Thr Ser Ser Phe Leu Leu Leu Thr Phe Ala Ile Cys Lys Leu Ile
    1             5             10             15

```

```

Ile Ala Val Gly Leu Asn Val Gly Pro Ser Glu Leu Leu Arg Ile Gly
    20             25             30

```

```

35 Ala Ile Asp Val Asp Gly His Phe Thr Val His Pro Ser Asp Leu Ala
    35             40             45

```

```

40 Ser Val Ser Ser Asp Phe Gly Met Leu Lys Ser Pro Glu Glu Pro Leu
    50             55             60

```

```

Ala Val Leu His Pro Ser Ser Ala Glu Asp Val Ala Arg Leu Val Arg
    65             70             75             80

```

```

45 Thr Ala Tyr Gly Ser Ala Thr Ala Phe Pro Val Ser Ala Arg Gly His
    85             90             95

```

```

Gly His Ser Ile Asn Gly Gln Ala Ala Ala Gly Arg Asn Gly Val Val
    100            105            110

```

```

50 Val Glu Met Asn His Gly Val Thr Gly Thr Pro Lys Pro Leu Val Arg
    115            120            125

```

```

55 Pro Asp Glu Met Tyr Val Asp Val Trp Gly Gly Glu Leu Trp Val Asp
    130            135            140

```

	Val	Leu	Lys	Lys	Thr	Leu	Glu	His	Gly	Leu	Ala	Pro	Lys	Ser	Trp	Thr	145	150	155	160
5	Asp	Tyr	Leu	Tyr	Leu	Thr	Val	Gly	Gly	Thr	Leu	Ser	Asn	Ala	Gly	Ile	165	170	175	
	Ser	Gly	Gln	Ala	Phe	His	His	Gly	Pro	Gln	Ile	Ser	Asn	Val	Leu	Glu	180	185	190	
10	Leu	Asp	Val	Val	Thr	Gly	Lys	Gly	Glu	Val	Met	Arg	Cys	Ser	Glu	Glu	195	200	205	
15	Glu	Asn	Thr	Arg	Leu	Phe	His	Gly	Val	Leu	Gly	Gly	Leu	Gly	Gln	Phe	210	215	220	
	Gly	Ile	Ile	Thr	Arg	Ala	Arg	Ile	Ser	Leu	Glu	Pro	Ala	Pro	Gln	Arg	225	230	235	240
20	Val	Arg	Trp	Ile	Arg	Val	Leu	Tyr	Ser	Ser	Phe	Lys	Val	Phe	Thr	Glu	245	250	255	
	Asp	Gln	Glu	Tyr	Leu	Ile	Ser	Met	His	Gly	Gln	Leu	Lys	Phe	Asp	Tyr	260	265	270	
25	Val	Glu	Gly	Phe	Val	Ile	Val	Asp	Glu	Gly	Leu	Val	Asn	Asn	Trp	Arg	275	280	285	
30	Ser	Ser	Phe	Phe	Ser	Pro	Arg	Asn	Pro	Val	Lys	Ile	Ser	Ser	Val	Ser	290	295	300	
	Ser	Asn	Gly	Ser	Val	Leu	Tyr	Cys	Leu	Glu	Ile	Thr	Lys	Asn	Tyr	His	305	310	315	320
35	Asp	Ser	Asp	Ser	Glu	Ile	Val	Asp	Gln	Glu	Val	Glu	Ile	Leu	Met	Lys	325	330	335	
	Lys	Leu	Asn	Phe	Ile	Pro	Thr	Ser	Val	Phe	Thr	Thr	Asp	Leu	Gln	Tyr	340	345	350	
40	Val	Asp	Phe	Leu	Asp	Arg	Val	His	Lys	Ala	Glu	Leu	Lys	Leu	Arg	Ser	355	360	365	
	Lys	Asn	Leu	Trp	Glu	Val	Pro	His	Pro	Trp	Leu	Asn	Leu	Phe	Val	Pro	370	375	380	
45	Lys	Ser	Arg	Ile	Ser	Asp	Phe	Asp	Lys	Gly	Val	Phe	Lys	Gly	Ile	Leu	385	390	395	400
50	Gly	Asn	Lys	Thr	Ser	Gly	Pro	Ile	Leu	Ile	Tyr	Pro	Met	Asn	Lys	Asp	405	410	415	
	Lys	Trp	Asp	Glu	Arg	Ser	Ser	Ala	Val	Thr	Pro	Asp	Glu	Glu	Val	Phe	420	425	430	
55																				

Tyr Leu Val Ala Leu Leu Arg Ser Ala Leu Thr Asp Gly Glu Glu Thr
 435 440 445
 5 Gln Lys Leu Glu Tyr Leu Lys Asp Gln Asn Arg Arg Ile Leu Glu Phe
 450 455 460
 Cys Glu Gln Ala Lys Ile Asn Val Lys Gln Tyr Leu Pro His His Ala
 465 470 475 480
 10 Thr Gln Glu Glu Trp Val Ala His Phe Gly Asp Lys Trp Asp Arg Phe
 485 490 495
 Arg Ser Leu Lys Ala Glu Phe Asp Pro Arg His Ile Leu Ala Thr Gly
 500 505 510
 15 Gln Arg Ile Phe Gln Asn Pro Ser Leu Ser Leu Phe Pro Pro Ser Ser
 515 520 525
 Ser Ser Ser Ser Ala Ala Ser Trp
 20 530 535
 <210> 11
 25 <211> 1936
 <212> DNA
 <213> Arabidopsis thaliana
 <400> 11
 30 atgcttatag taagaagttt caccatcttg cttctcagct gcatagcctt taagttggct 60
 tgctgcttct ctagcagcat ttcttctttg aaggcgcttc ccctagtagg ccatttggag 120
 tttgaacatg tccatcacgc ctccaaagat tttggaaatc gataccagtt gatccctttg 180
 gcggtcttac atcccaaate ggtaagcgac atcgccctca cgatacgaca catctggatg 240
 atgggcactc attcacagct tacagtggca gcgagaggctc gtggacattc actccaaggc 300
 35 caagctcaaa caagacatgg aattgttata cacatggaat cactccatcc ccagaagctg 360
 caggtctaca gtgtggattc ccctgctcca tatgttgatg tgtctgggtg tgagctgtgg 420
 ataaacattt tgcattgagac cctcaagtac gggcttgac caaatcatg gacggattac 480
 ctgcatttaa ctgtagggtg tactctgtcc aatgctggaa taagcggcca ggcatccga 540
 catggaccac agatcagcaa tgttcatcaa ctggagattg tcacagggtta gttcagagtt 600
 40 gcagtattcg tgttttgaag gcatagactc tatatgggtg gtgactatta acaacatgaa 660
 gagattcccg agaatagcta cccactaatg tcatgcctat ttattgactg caggaaaagg 720
 cgagatccta aactgtacaa agaggcagaa cagcgactta tttaatggtg ttcttgggtg 780
 tttagggtcag tttggcatca taacgcgggc aagaatagca ttggaaccag caccaaccat 840
 ggtaaacat aaataaataa aaaacttaaa aactgaacac gcgtgtgtcc tcctaactct 900
 45 gtataatgga caggtaaaat ggataagagt gttatacctg gattttgcag cttttgccaa 960
 ggaccaagag caactaatat ctgcccaggg ccacaaattc gattacatag aagggtttgt 1020
 gataataaac aggacaggcc tcctgaacag ctggagggtg tctttcaccg cagaagagcc 1080
 tttagaagca agccaattca agtttgatg aaggactctg tattgtctgg agctagccaa 1140
 gtatttgaag caagataaca aagacgtaat caaccagggtg agaaaacaga gtagaagcaa 1200
 50 tcggtagaat cttcttttgt agatgacatt cattggaact gaaaatatat atatatattgt 1260
 ccaatccagg aagtgaagaa aacattatca gagctaagct acgtgacgtc gacactgttt 1320
 acaacggagg tagcatatga agcattcttg gacagggtac atgtgtctga ggtaaaactc 1380
 cgatcgaaag ggcagtggga ggtgccacat ccatggctga acctcctggg accaagaagc 1440
 aaaatcaatg aatttgcaag aggtgtattt ggaaacatac taacggatac aagcaacggc 1500
 55 ccagtcacgt tctaccaggt gaacaaatca aagtaagaaa gaaagaaaga aagagctagt 1560

catgattttg tttcttttca cttgttgaca aaacaaaagc atgttggtga gcagggtggga 1620
 caatcaaaca tcagcagtaa caccggagga agaggtattc tacctggtgg cgatcctaac 1680
 atcggcatct ccagggtcgg caggaaagga tggagtagaa gagatcttga ggcggaacag 1740
 aagaatactg gaattcagtg aagaagcagg gatagggttg aagcagtatc tgccacatta 1800
 5 cagcacaaga gaagagtgga gatcccattt cggggacaag tggggagaat ttgtgaggag 1860
 gaaatccaga tatgatccat tggcaattct tgcgcctggc caccgaattt ttcaaaaggc 1920
 agtctcatatc tcatga 1936

10 <210> 12
 <211> 504
 <212> PRT
 <213> Arabidopsis thaliana

15 <400> 12
 Met Leu Ile Val Arg Ser Phe Thr Ile Leu Leu Leu Ser Cys Ile Ala
 1 5 10 15
 Phe Lys Leu Ala Cys Cys Phe Ser Ser Ser Ile Ser Ser Leu Lys Ala
 20 20 25 30
 Leu Pro Leu Val Gly His Leu Glu Phe Glu His Val His His Ala Ser
 35 40 45
 25 Lys Asp Phe Gly Asn Arg Tyr Gln Leu Ile Pro Leu Ala Val Leu His
 50 55 60
 Pro Lys Ser Val Ser Asp Ile Ala Ser Thr Ile Arg His Ile Trp Met
 65 70 75 80
 30 Met Gly Thr His Ser Gln Leu Thr Val Ala Ala Arg Gly Arg Gly His
 85 90 95
 Ser Leu Gln Gly Gln Ala Gln Thr Arg His Gly Ile Val Ile His Met
 35 100 105 110
 Glu Ser Leu His Pro Gln Lys Leu Gln Val Tyr Ser Val Asp Ser Pro
 115 120 125
 40 Ala Pro Tyr Val Asp Val Ser Gly Gly Glu Leu Trp Ile Asn Ile Leu
 130 135 140
 His Glu Thr Leu Lys Tyr Gly Leu Ala Pro Lys Ser Trp Thr Asp Tyr
 145 150 155 160
 45 Leu His Leu Thr Val Gly Gly Thr Leu Ser Asn Ala Gly Ile Ser Gly
 165 170 175
 Gln Ala Phe Arg His Gly Pro Gln Ile Ser Asn Val His Gln Leu Glu
 50 180 185 190
 Ile Val Thr Gly Lys Gly Glu Ile Leu Asn Cys Thr Lys Arg Gln Asn
 195 200 205
 55 Ser Asp Leu Phe Asn Gly Val Leu Gly Gly Leu Gly Gln Phe Gly Ile

	210	215	220
	Ile Thr Arg Ala Arg	Ile Ala Leu Glu Pro	Ala Pro Thr Met Asp Gln
	225	230	235 240
5	Glu Gln Leu Ile Ser	Ala Gln Gly His Lys Phe Asp Tyr Ile Glu Gly	
	245	250	255
10	Phe Val Ile Ile Asn Arg Thr Gly Leu Leu Asn Ser Trp Arg Leu Ser		
	260	265	270
	Phe Thr Ala Glu Glu Pro Leu Glu Ala Ser Gln Phe Lys Phe Asp Gly		
	275	280	285
15	Arg Thr Leu Tyr Cys Leu Glu Leu Ala Lys Tyr Leu Lys Gln Asp Asn		
	290	295	300
	Lys Asp Val Ile Asn Gln Glu Val Lys Glu Thr Leu Ser Glu Leu Ser		
	305	310	315 320
20	Tyr Val Thr Ser Thr Leu Phe Thr Thr Glu Val Ala Tyr Glu Ala Phe		
	325	330	335
	Leu Asp Arg Val His Val Ser Glu Val Lys Leu Arg Ser Lys Gly Gln		
25	340	345	350
	Trp Glu Val Pro His Pro Trp Leu Asn Leu Leu Val Pro Arg Ser Lys		
	355	360	365
30	Ile Asn Glu Phe Ala Arg Gly Val Phe Gly Asn Ile Leu Thr Asp Thr		
	370	375	380
	Ser Asn Gly Pro Val Ile Val Tyr Pro Val Asn Lys Ser Lys Trp Asp		
	385	390	395 400
35	Asn Gln Thr Ser Ala Val Thr Pro Glu Glu Glu Val Phe Tyr Leu Val		
	405	410	415
	Ala Ile Leu Thr Ser Ala Ser Pro Gly Ser Ala Gly Lys Asp Gly Val		
40	420	425	430
	Glu Glu Ile Leu Arg Arg Asn Arg Arg Ile Leu Glu Phe Ser Glu Glu		
	435	440	445
45	Ala Gly Ile Gly Leu Lys Gln Tyr Leu Pro His Tyr Thr Thr Arg Glu		
	450	455	460
	Glu Trp Arg Ser His Phe Gly Asp Lys Trp Gly Glu Phe Val Arg Arg		
	465	470	475 480
50	Lys Ser Arg Tyr Asp Pro Leu Ala Ile Leu Ala Pro Gly His Arg Ile		
	485	490	495
55	Phe Gln Lys Ala Val Ser Tyr Ser		
	500		

```

5    <210> 13
    <211> 31
    <212> DNA
    <213> Artificial Sequence

    <220>
10   <223> Description of Artificial Sequence:oligonucleotide
    : primer or probe

    <400> 13
    cggtcgacat gggattgacc tcatccttac g
15                                     31

    <210> 14
    <211> 35
    <212> DNA
20   <213> Artificial Sequence

    <220>
    <223> Description of Artificial Sequence:oligonucleotide
    : primer or probe
25

    <400> 14
    gcgtcgactt atacagttct aggtttcggc agtat
30                                     35

    <210> 15
    <211> 33
    <212> DNA
    <213> Artificial Sequence

35   <220>
    <223> Description of Artificial Sequence:oligonucleotide
    : primer or probe

    <400> 15
40   gcggtaccag agagagaaac ataaacaaat ggc
33

    <210> 16
    <211> 31
45   <212> DNA
    <213> Artificial Sequence

    <220>
50   <223> Description of Artificial Sequence:oligonucleotide
    : primer or probe

    <400> 16
    gcggtacca attttacttc caccaaaatg c
55                                     31

```

<210> 17
<211> 34
<212> DNA
<213> Artificial Sequence
5
<220>
<223> Description of Artificial Sequence:oligonucleotide
: primer or probe
10 <400> 17
gcggtacctt cattgataag aatcaagcta ttca 34

<210> 18
15 <211> 31
<212> DNA
<213> Artificial Sequence

<220>
20 <223> Description of Artificial Sequence:oligonucleotide
: primer or probe

<400> 18
25 gcggtaccca aagtggtag aacgactaac a 31

<210> 19
<211> 28
<212> DNA
30 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
: primer or probe
35
<400> 19
gcggtacccc cattaaccta cccgtttg 28

<210> 20
40 <211> 32
<212> DNA
<213> Artificial Sequence

<220>
45 <223> Description of Artificial Sequence:oligonucleotide
: primer or probe

<400> 20
50 gcggtaccag acgatgaacg tacttgtctg ta 32

<210> 21
<211> 28
55 <212> DNA

<213> Artificial Sequence

<220>

5 <223> Description of Artificial Sequence:oligonucleotide
: primer or probe

<400> 21
ggggtacctt gatgaatcgt gaaatgac 28

10 <210> 22
<211> 31
<212> DNA
<213> Artificial Sequence

15 <220>
<223> Description of Artificial Sequence:oligonucleotide
: primer or probe

20 <400> 22
ggggtaccct ttctctcttg ttttgcctg t 31

25 <210> 23
<211> 32
<212> DNA
<213> Artificial Sequence

30 <220>
<223> Description of Artificial Sequence:oligonucleotide
: primer or probe

35 <400> 23
gctctagatc aggaaaagaa ccatgcttat ag 32

40 <210> 24
<211> 32
<212> DNA
<213> Artificial Sequence

45 <220>
<223> Description of Artificial Sequence:oligonucleotide
: primer or probe

50 <400> 24
gctctagatc atgagtatga gactgccttt tg 32

55 <210> 25
<211> 1728
<212> DNA
<213> Arabidopsis thaliana

55 <400> 25

	atgggattga	cctcatcctt	acggttccat	agacaaaaca	acaagacttt	cctcggaaac	60
	ttcatgatct	tagttctaag	ctgtatacca	ggtagaacca	atctttgttc	caatcattct	120
	gttagtacct	caaaagaatt	accttcttca	aatccttcag	atattcggtc	ctcattagtt	180
	tcactagatt	tgagggggta	tataagcttc	gacgatgtcc	acaatgtggc	caaggacttt	240
5	ggcaacagat	accagttacc	acctttggca	attctacatc	caaggtcagt	ttttgatatt	300
	tcatcgatga	tgaagcatat	agtacatctg	ggctccacct	caaactctac	agtagcagct	360
	agaggccatg	gtcactcgct	tcaaggacaa	gctctagctc	atcaagggtg	tgtcatcaaa	420
	atggagtcac	ttcgaagtcc	tgatatcagg	atctataagg	ggaagcaacc	atatgttgat	480
	gtctcagggtg	gtgaaatatg	gataaacatt	ctacgcgaga	ctctaaaata	cggctcttca	540
10	ccaaagtcct	ggacagacta	ccttcatttg	accgttggag	gtacactatc	taatgctgga	600
	atcagcgggtc	aagcattcaa	gcatggaccc	caaactcaaca	acgtctacca	gctagagatt	660
	gttagcaggga	aaggagaagt	cgtaacctgt	tctgagaagc	ggaattctga	acttttcttc	720
	agtgttcttg	gcgggcttgg	acagtttggc	ataatcacc	gggcacggat	ctctcttgaa	780
	ccagcacgcg	atatgggttaa	atggatcagg	gtactctact	ctgacttttc	tgcattttca	840
15	agggaccaag	aatatctgat	ttcgaaggag	aaaacttttg	attacgttga	aggattttgtg	900
	ataatcaata	gaacagacct	tctcaataat	tggcgatcgt	cattcagtc	caacgattcc	960
	acacaggcaa	gcagattcaa	gtcagatggg	aaaactcttt	attgcctaga	agtgggtcaaa	1020
	tattttcaacc	cagaagaagc	tagctctatg	gatcaggaaa	ctggcaagtt	acttttcagag	1080
	ttaaattata	ttccattcc	tttgttttca	tctgaagtgc	catatatcga	gtttctggat	1140
20	cgctgcata	tcgcagagag	aaaactaaga	gcaaagggtt	tatgggagg	tccacatccc	1200
	tggttgaatc	tcctgattcc	taagagcagc	atataccaat	ttgctacaga	agttttcaac	1260
	aacattctca	caagcaacaa	caacggctct	atccttattt	atccagtc	tcaatccaag	1320
	tggagaagaa	atacatcttt	gataactcca	aatgaagata	tattctatct	cgtagccttt	1380
	ctccccctctg	cagtgcctaa	ttctcagggt	aaaaacgata	tagagtacct	tttgaaacaa	1440
25	aaccaagag	ttatgaactt	ctgcgcagca	gcaaacctca	acgtgaagca	gtattttgcc	1500
	cattatgaaa	ctcaaaaaga	gtggaaatca	cactttggca	aaagatggga	aacatttgca	1560
	cagaggaaac	aagcctacga	ccctctagcg	attctagcac	ctggccaaag	aatattccaa	1620
	aagacaacag	gaaaattatc	tcccatccaa	ctcgcaaggt	caaaggcaac	aggaagtcct	1680
	caaagggtacc	attacgcata	aatactgccc	aaacctagaa	ctgtataa		1728
30							
	<210> 26						
	<211> 1506						
	<212> DNA						
35	<213> Arabidopsis thaliana						
	<400> 26						
	atggctaata	ttcgtttaat	gatcacttta	atcacgggtt	taatgatcac	caaactatca	60
	aacgggtatta	aaattgattt	acctaatacc	cttaacctca	ccctctctac	cgatccttcc	120
40	atcatctccg	cagcctctca	tgacttcgga	aacataacca	ccgtgacccc	cggcggcgta	180
	atctgcccct	cctccaccgc	tgatatctct	cgtctcctcc	aatacgccgc	aaacggaaaa	240
	agtacattcc	aagtagcggc	tcgtggccaa	ggccactcct	taaacggcca	agcctcgggtc	300
	tccggcggag	taatcgta	catgacgtgt	atcactgacg	tggtgggttc	aaaagacaag	360
	aagtacgctg	acgtggcggc	cgggacgtta	tgggtggatg	tgcttaagaa	gacggcggag	420
45	aaaggggtgt	cgccgggttc	ttggacggat	tatttgcata	taaccgtcgg	aggaacgttg	480
	tcgaattggtg	gaattggtgg	tcaagtgttt	cgaaacgggtc	ctcttggttag	taacgtcctt	540
	gaattggacg	ttattactgg	gaaagggtgaa	atgttgacat	gctcgcgaca	gctaaaccca	600
	gaattgttct	atggagtgtt	aggagggtttg	ggtcaatttg	gaattataac	gagagccaga	660
	attgttttgg	accatgcacc	taaacggggc	aaatgggttc	ggatgctcta	cagtgtattc	720
50	acaactttta	caaaggacca	agaacgtttg	atatcaatgg	caaacgatat	tggagtcgac	780
	tatttagaag	gtcaaatatt	tctatcaaac	ggtgtcgttg	acacctcttt	tttccacct	840
	tcagatcaat	ctaaagtgcg	tgatctagtc	aagcaacacg	gtatcatcta	tgttcttgaa	900
	gtagccaagt	attatgatga	tcccaatctc	cccatcatca	gcaagggttat	tgacacatta	960
	acgaaaacat	taagttactt	gccgggttcc	atatcaatgc	acgacgtggc	ctacttcgat	1020
55	ttcttgaacc	gtgtacatgt	cgaagaaaat	aaactcagat	ctttgggatt	atgggaactt	1080

cctcatcctt ggcttaacct ctacgttcct aaatctcgga ttctcgattt tcataacggt 1140
 gttgtcaaag acattcttct taagcaaaaa tcagcttcgg gactcgtctt tctctatcca 1200
 acaaaccgga ataaatggga caatcgtatg tcggcgatga taccagagat cgatgaagat 1260
 gttatatata ttatcggact actacaatcc gctaccccaa aggatcttcc agaagtggag 1320
 5 agcggttaacg agaagataat taggttttgc aaggattcag gtattaagat taagcaatat 1380
 ctaatgcatt atactagtaa agaagattgg attgagcatt ttggatcaaa atgggatgat 1440
 ttttcgaaga ggaaagatct atttgatccc aagaaactgt tatctccagg gcaagacatc 1500
 ttttga 1506

10 <210> 27
 <211> 1572
 <212> DNA
 <213> Arabidopsis thaliana

15 <400> 27
 atggcgagtt ataatcttctg ttacacaagtt cgtcttatag caataacaat agtaatcatc 60
 attactctct caactccgat cacaaccaac acatcaccac aaccatggaa tatcctttca 120
 cacaacgaat tcgcgggaaa actcacctcc tctctctctt cgcgcgaatc agccgccaca 180
 20 gatttcggcc acgtcaccaa aatcttccct tccgcgtctt taatcccttc ctccgttgaa 240
 gacatcacag atctcataaa actctctttt gactctcaac tgtcttttcc tttagccgct 300
 cgtggtcacg gacacagcca ccgtggccaa gcctcggcta aagacggagt tgtggtcaac 360
 atgcggtcca tggtaaaccg ggatcgaggt atcaagggtg ctaggacctg tttatatgtt 420
 gacgttgacg ctgctgggct atggattgag gtgttgaata aaactttgga gttagggtta 480
 25 acgcgggttt cttggacgga ttatttgtat ttaacagtcg gtgggacgtt atcaaacggc 540
 ggaattagtg gacaaacgtt tcggtacggt ccacagatca ctaatgttct agagatggat 600
 gttattactg gaaaaggaga gattgcaact tgttccaagg acatgaactc ggatcttttc 660
 ttgcggtgt taggaggttt gggatcaattc ggcattataa caagagccag aattaaactt 720
 gaagtagctc cgaaaagggc caagtgttta aggtttctat acatagattt ctccgaattc 780
 30 acaagagatc aagaacgagt gatatcgaaa acggacggtg tagatttctt agaaggttcc 840
 attatggtyg accatggccc accggataac tggagatcca cgtattatcc accgtccgat 900
 cacttgagga tcgcctcaat ggtcaaacga catcgtgtca tctactgcct tgaagtcgtc 960
 aagtattacg acgaaacttc tcaatacaca gtcaacgagg aaatggagga gtttaagcgt 1020
 agtttaaac atgtaagagg gtttatgtac gagaaagatg tgacgtatat ggatttccta 1080
 35 aaccgagttc gaaccggaga gctaaacctg aaatccaaag gccaatggga tgttccacat 1140
 ccatggctta atctcttcgt accaaaaact caaatctcca aatttgatga tgggtgtttt 1200
 aagggtatta tcctaagaaa taacatcact agcggctctg ttcttgttta tcctatgaat 1260
 cgcaacaagt ggaatgatcg gatgtctgcc gctatacccg aggaagatgt attttatgcy 1320
 gtagggtttt taagatccgc gggttttgac aattgggagg cttttgatca agaaaacatg 1380
 40 gaaatactga agttttgtga ggatgcta atgggggtta tacaatatct tccttatcat 1440
 tcatcacaag aaggatgggt tagacatttt ggtccgaggt ggaatatatt cgtagagaga 1500
 aaatataaat atgatcccaa aatgatatta tcaccgggac aaaatatatt tcaaaaaata 1560
 aactcgagtt ag 1572

45 <210> 28
 <211> 1575
 <212> DNA
 <213> Arabidopsis thaliana

50 <400> 28
 atgactaata ctctctgttt aagcctcatc accctaataa cgctttttat aagtttaacc 60
 ccaaccttaa tcaaatcaga tgagggcatt gatgttttct taccatatac actcaacctt 120
 acggctctaa ccgatccctt ctccatctct gccgcttctc acgacttcgg taacataacc 180
 55 gacgaaaatc ccggcgccgt cctctgcctt tctccacca cggaggtggc tcgtctcctc 240

	cgtttcgcta	acggaggatt	ctcttacaat	aaaggctcaa	ccagccccgc	gtctactttc	300
	aaagtggctg	ctcgaggcca	aggccactcc	ctccgtggcc	aagcctctgc	acccggaggt	360
	gtcgtcgtga	acatgacgtg	tctcgccatg	gcggctaaac	cagcggcggg	tggtatctcg	420
	gcagacggga	cttacgctga	cgtggctgcc	gggacgatgt	gggtggatgt	tctgaaggcg	480
5	gcggtggata	gaggcgtctc	gccggttaca	tggacggatt	atttgtatct	cagcgtcggc	540
	gggacgttgt	cgaacgctgg	aatcgggtgt	cagacgttta	gacacggccc	tcagattagt	600
	aacgttcctg	agcttgacgt	tattaccgga	aaaggtgaaa	tgatgacttg	ctctccaaag	660
	ttaaaccctg	aattgttcta	tggagtttta	ggaggtttgg	gtcaattcgg	tattataacg	720
	agggccagga	ttgcgttggg	tcatgcaccc	acaagggtga	aatggctctg	catactctac	780
10	agtgaacttct	cggtctttaa	aagagaccaa	gagcgtttaa	tatcaatgac	caatgatctc	840
	ggagttgact	ttttggaagg	tcaacttatg	atgtcaaatg	gcttcgtaga	cacctctttc	900
	ttcccactct	ccgatcaaac	aagagtcgca	tctcttgtga	atgaccaccg	gaticatctat	960
	gttctcgaag	tagccaagta	ttatgacaga	accacccttc	ccattattga	ccaggtgatt	1020
	gacacgttaa	gtagaactct	aggtttcgtc	ccagggttta	tgttcgtaca	agatgttccg	1080
15	tatttcgatt	tcttgaaccg	tgccgaaac	gaagaagata	aactcagatc	tttaggacta	1140
	tgggaagtct	ctcatccatg	gcttaacatc	tttgtcccgg	ggtctcgaat	ccaagatttt	1200
	catgatgggt	ttattaatgg	ccttcttcta	aaccaaacct	caacttctgg	tgttactctc	1260
	ttctatccca	caaaccgaaa	caaatggaac	aaccgcgatg	caacgatgac	accggacgaa	1320
	gatgtttttt	atgtgatcgg	attactgcaa	tcagctgggt	gatctcaaaa	ttggcaagaa	1380
20	cttgaaaaac	tcaacgacaa	ggttattcag	ttttgtgaaa	actcgggaat	taagattaag	1440
	gaatatattga	tgcactatac	aagaaaagaa	gattgggtta	aacattttgg	accaaaatgg	1500
	gatgattttt	taagaaagaa	aattatgttt	gatcccaaaa	gactattgtc	tccaggacaa	1560
	gacatatatta	attaa					1575
25	<210> 29						
	<211> 1611						
	<212> DNA						
	<213> Arabidopsis thaliana						
30	<400> 29						
	atgacgtcaa	gctttcttct	cctgacgttc	gccatatgta	aactgatcat	agccgtgggt	60
	ctaaacgtgg	gccccagtga	gtcctctcgc	atcggagcca	tagatgtcga	cggccacttc	120
	accgtccacc	cttccgactt	agcctccgtc	tcctcagact	tcggtatgct	gaagtcacct	180
35	gaagagccat	tggcctgtct	tcatccatca	tcggccgaag	acgtggcacg	actcgtcaga	240
	acagcttacg	gttcagccac	ggcgtttccg	gtctcagccc	gaggccacgg	ccattccata	300
	aacggacaag	ccgcggcggg	gaggaacggt	gtgggtggtg	aaatgaacca	cggcgtaacc	360
	gggacgcccc	agccactcgt	ccgaccggat	gaaatgtatg	tggatgtatg	gggtggagag	420
	ttatgggtcg	atgtgttgaa	gaaaacgttg	gagcatggct	tagcaccaaa	atcatggacg	480
40	gattactttg	atctaaccgt	tggaggtaca	ctctccaatg	caggaatcag	tgggtcaagct	540
	tttcaccatg	gtcctcaaat	tagtaacgtc	cttgagctcg	acgttgtaac	tgggaaagga	600
	gaggtgatga	gatgctcaga	agaagagaac	acaaggctat	tccatggagt	tcttggtgga	660
	ttaggtcaat	ttgggatcat	cactcgagca	cgaatctctc	tcgaaccagc	tccccaaagg	720
	gtgagatgga	tacgggtatt	gtattcgagc	ttcaaagtgt	ttacggagga	ccaagagtac	780
45	ttaatctcaa	tgcattggtc	attaaagttt	gattacgtgg	aaggttttgt	gattgtggac	840
	gaaggactcg	tcaacaattg	gagatcttct	ttcttctctc	cacgtaacct	cgtcaagatc	900
	tcctctgtta	gttccaacgg	ctctgttttg	tattgccttg	agatcaccaa	gaactaccac	960
	gactccgact	ccgaaatcgt	tgatcaggaa	gttgagattc	tgatgaagaa	attgaatttc	1020
	ataccgacat	cgggtctttac	aacggattta	caatatgtgg	actttctcga	ccgggtacac	1080
50	aaggccgaat	tgaagctccg	gtccaagaat	ttatgggagg	ttccacacct	atgggtcaac	1140
	ctcttcgtgc	caaaatcaag	aatctctgac	ttcgataaag	gcgtttttcaa	gggcattttg	1200
	ggaaataaaa	caagtggccc	tattcttata	taccccatga	acaaagacaa	atgggacgag	1260
	aggagctcag	ccgtgacggc	ggatgaggaa	gttttctatc	tgggtggctct	attgagatca	1320
	gctttaaccg	acgggtgaaga	gacacagaag	ctagagtatc	tgaaagatca	gaaccgtcgg	1380
55	atcttggagt	tctgtgaaca	agccaagatc	aatgtgaagc	agtatcttcc	tcaccacgca	1440

```

acacaggaag agtgggtggc tcattttggg gacaagtggg atcggttcag aagcttaaag 1500
gctgagtttg atccgcgaca catactcgct actggtcaga gaatctttca aaacccatct 1560
ttgtctttgt ttccctccgc gtcgtcttct tcgtcagcgg cttcatggtg a 1611

```

5

```

<210> 30
<211> 1515
<212> DNA
<213> Arabidopsis thaliana

```

10

```

<400> 30
atgcttatag taagaagttt caccatcttg cttctcagct gcatagcctt taagttggct 60
tgctgcttct ctagcagcat ttcttctttg aaggcgcttc ccctagtagg ccatttggag 120
tttgaacatg tccatcacgc ctccaaagat ttgggaaatc gataccagtt gatccctttg 180
15 gcggtcttac atcccaaadc ggtaagcgac atcgctcaa cgatacgaca catctggatg 240
atgggcactc attcacagct tacagtggca gcgagaggtc gtggacattc actccaaggc 300
caagctcaaa caagacatgg aattgttata cacatggaat cactccatcc ccagaagctg 360
cagggtctaca gtgtggattc ccctgctcca tatgttgatg tgtctgggtg tgagctgttg 420
ataaacattt tgcagagac cctcaagtac gggcttgac caaaatcatg gacggattac 480
20 ctgcatttaa ctgtagggtg tactctgtcc aatgctggaa taagcggcca ggcattccga 540
catggaccac agatcagcaa tgttcatcaa ctggagattg tcacaggaaa aggcgagatc 600
ctaaactgta caaagaggca gaacagcgac ttatttaatg gtgttcttgg tggtttaggt 660
cagtttggca tcataacgag ggcaagaata gcattggaac cagcaccaac catggacca 720
gagcaactaa tatctgcccc ggccacaaa ttcgattaca tagaagggtt tgtgataata 780
25 aacaggacag gcctcctgaa cagctggagg ttgtctttca ccgcagaaga gcctttagaa 840
gcaagccaat tcaagtttga tgggaaggact ctgtattgtc tggagctagc caagtatttg 900
aagcaagata acaaagacgt aatcaaccag gaagtgaag aaacattatc agagctaagc 960
tacgtgacgt cgacactgtt tacaacggag gtagcatatg aagcattctt ggacagggt 1020
catgtgtctg aggtaaaact ccgatcgaaa gggcagtggg aggtgccaca tccatggctg 1080
30 aacctcctgg taccaagaag caaaatcaat gaatttgcaa gaggtgtatt tggaaacata 1140
ctaacggata caagcaacgg ccagtcacg gtctaccag tgaacaaatc aaagtgggac 1200
aatcaaacat cagcagtaac accggaggaa gaggtattct acctggtggc gatcctaaca 1260
tcggcatctc cagggtcggc aggaaggat ggagtagaag agatcttgag gcggaacaga 1320
agaatactgg aattcagtg agaagcagg ataggggtga agcagtatct gccacattac 1380
35 acgacaagag aagagtggag atcccatctt ggggacaagt ggggagaatt tgtgaggagg 1440
aatccagat atgatccatt ggcaattctt gcgcctggcc accgaatttt tcaaaaggca 1500
gtctcactac catga 1515

```

40

```

<210> 31
<211> 84
<212> DNA
<213> Arabidopsis thaliana

```

45

```

<400> 31
tcagcttcgg gactcgctct tctctatcca acaaaccgga ataatggga caatcgatg 60
tcggcgatga taccagagat cgat 84

```

50

```

<210> 32
<211> 28
<212> PRT
<213> Arabidopsis thaliana

```

55

```

<400> 32

```

Ser Ala Ser Gly Leu Ala Leu Leu Tyr Pro Thr Asn Arg Asn Lys Trp
 1 5 10 15

Asp Asn Arg Met Ser Ala Met Ile Pro Glu Ile Asp
 5 20 25

<210> 33

10 <211> 2814

<212> DNA

<213> Arabidopsis thaliana

<400> 33

15 atgaatcgta tgacgtcaag ctttcttctc ctgacgttcg ccatatgtaa actgatcata 60
 gccgtgggtc taaacgtggg cccagtgag ctectccgca tcggagccat agatgtcgac 120
 ggccacttca ccgtccaccc ttccgactta gcctccgtct cctcagactt cggtatgctg 180
 aagtcacctg aagagccatt ggccgtgctt catccatcat cggccgaaga cgtggcacga 240
 ctcgtcagaa cagcttacgg ttccagccacg gcgtttccgg tctcagcccg aggccacggc 300
 20 cattccataa acggacaagc cgcggcgggg aggaacggtg tgggtggtga aatgaaccac 360
 ggcgtaaccg ggacgcccac gccactcgtc cgaccggatg aaatgtatgt ggatgtatgg 420
 ggtggagagt tatgggtcga tgtgttgaa aaaacgttgg agcatggctt agcaccacaaa 480
 tcatggacgg attacttgta tctaaccgtt ggaggtacac tctccaatgc aggaatcagt 540
 ggtcaagctt ttcaccatgg tcttcaaatt agtaacgtcc ttgagctcga cgttgtaact 600
 25 ggttagtatt aaaacattca agttcatata ttttaaatgc ttttgtctga agttttacta 660
 ataacaagaa attgatacca aaaagtaggg aaaggagagg tgatgagatg ctcagaagaa 720
 gagaacacaa ggctattcca tggagtctct ggtggattag gtcaatttgg gatcatcact 780
 cgagcacgaa tctctctcga accagctccc caaagggtaa tattttttta atgactagct 840
 atcaaaaatc cctggcgggg ccatacgttg taatcttttt agtttttact gttgatggta 900
 30 ttttttatat attttggata ataaaaccct aaaatgggat atttgtatga cagggtgagat 960
 ggatacgggt attgtattcg agcttcaag tgtttacgga ggaccaagag tacttaactct 1020
 caatgcattg tcaattaaag tttgattacg tggaaaggtt tgtgattgtg gacgaaggac 1080
 tcgtcaacaa ttggagatct tctttcttct ctccacgtaa ccccgtaag atctctctctg 1140
 ttagttccaa cggctctgtt ttgtattgcc ttgagatcac caagaactac cacgactccg 1200
 35 actccgaaat cgttgatcag gtcactttca ttattcactt agaaaaaagc gatattttca 1260
 ttttttatat tgatgaatat ctggaaggat ttaacgctat gcgactattg ggaaatcatt 1320
 atgaaaaaat atttagttta tatgattgaa agtggctctc atagtatttt tgttgtgtcg 1380
 actttattat aacttaaat tggaaagaga catgaagaag aagccagaga ggatctacag 1440
 agatctagct tttccacctg aacttaataa tgcacattta tataattatt tttcttcttc 1500
 40 taaagtttag tttatcacta gcgaattaat catggttact aattaagtag tggacagggg 1560
 catggaccac tcaactacca aataatgatt cctctttact ctttaagttta attttaataa 1620
 aaccaactct actggaatct taacttatcc ttggttttgg taggctttta tagcaacacg 1680
 gtttttttaa ttttcttatt ccagattttg tatattaaat gtcgattttt tttctttttg 1740
 tttcaggaag ttgagattct gatgaagaaa ttgaatttca taccgacatc ggtctttaca 1800
 45 acggatttac aatatgtgga ctttctcgac cgggtacaca aggccgaatt gaagctccgg 1860
 tccaagaatt tatgggaggt tccacacca tggctcaacc tcttcgtgcc aaaatcaaga 1920
 atctctgact tcgataaagg cgttttcaag ggcatttttg gaaataaaac aagtggccct 1980
 attcttatct accccatgaa caaagacaag taagctctga cattaccatt gattactact 2040
 tctaaatttc ttctctagaa aaaagaataa aacgagtttt gcattgcatg catgcaaaagt 2100
 50 tacacttggt gggatttatt agtgggtcaa gaaaaaagt ttgtcaaaat tgaaaaaac 2160
 tagacacgtg tgatcagggg ttgtccgaaa aacgttgctc acatgtgcat cgaaccagct 2220
 aagattgaca acaacacttc gtcggctcgt atttctcttt ttgttttgtg accaaatccg 2280
 atgggtccaga ttgggtttat ttgtttttta gttcctagaa ctcatgggtg gtgggtccca 2340
 atcagattct cctagaccaa accgatctca acgaaccctc cgcacatcat tgattattac 2400
 55 attaatatag atattgtcgt tgctgacgtg tcgtaatttg atgttattgt cagatgggac 2460

gagaggagct cagccgtgac gccggatgag gaagttttct atctgggtggc tctattgaga 2520
 tcagcttttaa cggacgggtga agagacacag aagctagagt atctgaaaga tcagaaccgt 2580
 cggatccttgg agttctgtga acaagccaag atcaatgtga agcagtatct tcctcaccac 2640
 gcaacacagg aagagtgggt ggctcatttt ggggacaagt gggatcgggt cagaagctta 2700
 5 aaggctgagt ttgatccgcg acacatactc gctactggtc agagaatctt tcaaaaccca 2760
 tctttgtctt tgtttcctcc gtcgtcgtct tcttcgtcag cggcttcatt gtga 2814

<210> 34

10 <211> 1620

<212> DNA

<213> *Arabidopsis thaliana*

<400> 34

15 atgaatcgta tgacgtcaag ctttcttctc ctgacgttcg ccatatgtaa actgatcata 60
 gccgtgggtc taaacgtggg cccagtgag ctccctccgca tcggagccat agatgtcgac 120
 ggccacttca cgtccaccc ttccgactta gcctccgtct cctcagactt cggtagctg 180
 aagtcacctg aagagccatt ggccgtgctt catccatcat cggccgaaga cgtggcacga 240
 ctctgcagaa cagcttacgg ttccagccacg gcgtttccgg tctcagcccg aggccacggc 300
 20 cattccataa acggacaagc cgcggcgggg aggaacgggt tgggtggttga aatgaaccac 360
 ggcgtaaccg ggacgcccac gccactcgtc cgaccggatg aaatgtatgt ggatgtatgg 420
 ggtggagagt tatgggtcga tgtgttgaag aaaacgttgg agcatggctt agcaccacaaa 480
 tcatggacgg attacttgta tctaaccgtt ggaggtacac tctccaatgc aggaatcagt 540
 ggtcaagctt ttcaccatgg tctcacaatt agtaacgtcc ttgagctcga cgttgtaact 600
 25 gggaaaggag aggtgatgag atgctcagaa gaagagaaca caaggctatt ccatggagtt 660
 cttggtggat taggtcaatt tgggatcatc actcagacac gaatctctct cgaaccagct 720
 ccccaaaggg tgagatggat acgggtattg tattcgagct tcaaagtgtt tacggaggac 780
 caagagtact taatctcaat gcatgggtcaa ttaaagtttg attacgtgga aggttttgtg 840
 attgtggacg aaggactcgt caacaatttg agatcttctt tcttctctcc acgtaacccc 900
 30 gtcaagatct cctctgttag ttccaacggc tctgttttgt attgccttga gatcaccaag 960
 aactaccacg actccgactc cgaaatcggt gatcaggaag ttgagattct gatgaagaaa 1020
 ttgaatttca taccgacatc ggtctttaca acggatttac aatatgtgga ctttctcgac 1080
 cgggtacaca aggccgaatt gaagctccgg tccaagaatt tatgggaggt tccacaccca 1140
 tggctcaacc tcttcgtgcc aaaatcaaga atctctgact tcgataaagg cgttttcaag 1200
 35 ggcatttttg gaaataaaaac aagtggccct attcttatct accccatgaa caaagacaaa 1260
 tgggacgaga ggagctcagc cgtgacgccg gatgaggaag ttttctatct ggtggctcta 1320
 ttgagatcag ctttaacgga cgggtgaagag acacagaagc tagagtatct gaaagatcag 1380
 aaccgtcgga tcttggagtt ctgtgaacaa gccaaagatca atgtgaagca gtatcttctt 1440
 caccacgcaa cacaggaaga gtgggtggct cattttgggg acaagtggga tcggttcaga 1500
 40 agcttaaagg ctgagtttga tccgcgacac atactcgcta ctggtcagag aatctttcaa 1560
 aacccatctt tgtcttttgt tctccgctcg tcgtcttctt cgtcagcggc ttcattggtga 1620

<210> 35

45 <211> 539

<212> PRT

<213> *Arabidopsis thaliana*

<400> 35

50 Met Asn Arg Met Thr Ser Ser Phe Leu Leu Leu Thr Phe Ala Ile Cys
 1 5 10 15

Lys Leu Ile Ile Ala Val Gly Leu Asn Val Gly Pro Ser Glu Leu Leu
 20 25 30

55

Arg Ile Gly Ala Ile Asp Val Asp Gly His Phe Thr Val His Pro Ser
 35 40 45
 5 Asp Leu Ala Ser Val Ser Ser Asp Phe Gly Met Leu Lys Ser Pro Glu
 50 55 60
 Glu Pro Leu Ala Val Leu His Pro Ser Ser Ala Glu Asp Val Ala Arg
 65 70 75 80
 10 Leu Val Arg Thr Ala Tyr Gly Ser Ala Thr Ala Phe Pro Val Ser Ala
 85 90 95
 Arg Gly His Gly His Ser Ile Asn Gly Gln Ala Ala Ala Gly Arg Asn
 100 105 110
 15 Gly Val Val Val Glu Met Asn His Gly Val Thr Gly Thr Pro Lys Pro
 115 120 125
 Leu Val Arg Pro Asp Glu Met Tyr Val Asp Val Trp Gly Gly Glu Leu
 130 135 140
 Trp Val Asp Val Leu Lys Lys Thr Leu Glu His Gly Leu Ala Pro Lys
 145 150 155 160
 25 Ser Trp Thr Asp Tyr Leu Tyr Leu Thr Val Gly Gly Thr Leu Ser Asn
 165 170 175
 Ala Gly Ile Ser Gly Gln Ala Phe His His Gly Pro Gln Ile Ser Asn
 180 185 190
 30 Val Leu Glu Leu Asp Val Val Thr Gly Lys Gly Glu Val Met Arg Cys
 195 200 205
 Ser Glu Glu Glu Asn Thr Arg Leu Phe His Gly Val Leu Gly Gly Leu
 210 215 220
 Gly Gln Phe Gly Ile Ile Thr Arg Ala Arg Ile Ser Leu Glu Pro Ala
 225 230 235 240
 40 Pro Gln Arg Val Arg Trp Ile Arg Val Leu Tyr Ser Ser Phe Lys Val
 245 250 255
 Phe Thr Glu Asp Gln Glu Tyr Leu Ile Ser Met His Gly Gln Leu Lys
 260 265 270
 45 Phe Asp Tyr Val Glu Gly Phe Val Ile Val Asp Glu Gly Leu Val Asn
 275 280 285
 Asn Trp Arg Ser Ser Phe Phe Ser Pro Arg Asn Pro Val Lys Ile Ser
 290 295 300
 Ser Val Ser Ser Asn Gly Ser Val Leu Tyr Cys Leu Glu Ile Thr Lys
 305 310 315 320
 55 Asn Tyr His Asp Ser Asp Ser Glu Ile Val Asp Gln Glu Val Glu Ile

		325		330		335	
	Leu Met Lys Lys Leu Asn Phe Ile Pro Thr Ser Val Phe Thr Thr Asp						
		340		345		350	
5	Leu Gln Tyr Val Asp Phe Leu Asp Arg Val His Lys Ala Glu Leu Lys						
		355		360		365	
10	Leu Arg Ser Lys Asn Leu Trp Glu Val Pro His Pro Trp Leu Asn Leu						
		370		375		380	
	Phe Val Pro Lys Ser Arg Ile Ser Asp Phe Asp Lys Gly Val Phe Lys						
		385		390		395	
15	Gly Ile Leu Gly Asn Lys Thr Ser Gly Pro Ile Leu Ile Tyr Pro Met						
		405		410		415	
	Asn Lys Asp Lys Trp Asp Glu Arg Ser Ser Ala Val Thr Pro Asp Glu						
		420		425		430	
20	Glu Val Phe Tyr Leu Val Ala Leu Leu Arg Ser Ala Leu Thr Asp Gly						
		435		440		445	
	Glu Glu Thr Gln Lys Leu Glu Tyr Leu Lys Asp Gln Asn Arg Arg Ile						
25		450		455		460	
	Leu Glu Phe Cys Glu Gln Ala Lys Ile Asn Val Lys Gln Tyr Leu Pro						
		465		470		475	
30	His His Ala Thr Gln Glu Glu Trp Val Ala His Phe Gly Asp Lys Trp						
		485		490		495	
	Asp Arg Phe Arg Ser Leu Lys Ala Glu Phe Asp Pro Arg His Ile Leu						
		500		505		510	
35	Ala Thr Gly Gln Arg Ile Phe Gln Asn Pro Ser Leu Ser Leu Phe Pro						
		515		520		525	
	Pro Ser Ser Ser Ser Ser Ser Ala Ala Ser Trp						
40		530		535			
	<210> 36						
45	<211> 842						
	<212> DNA						
	<213> Arabidopsis thaliana						
	<400> 36						
50	aagcttaaat gacaatttag taccttgggt tgggtcatgat ttagagcgga acaaataac 60						
	catacatcaa acgaggatat acagagaaaa ttcatggaag tatggaattt agaggacaat 120						
	ttctcttctg ggctacaacg gaccggccca ttgcgtcatt taccagagg tatcgagttt 180						
	gtggactttt gatgccgcta gagactattg gcatcggatt gaaaaaatg tttacttcgt 240						
	tggttaacaat tttctgaatg caatattttc cttgtcatga atatttaaac ttgttattac 300						
55	tttcttttag cttaggtgtg gacaattatg gagtttactt caaacgagga agaattctaa 360						

acgctcgggt caggtctcga aaacaaacca actcacaatc ctgacttaat tgaggaaaac 420
aatgcaaaac cacatgcatg cttccatatt tctatcataa tcttataaga aaaaacacta 480
ctaagtgaag tgattctgta tatatataac caatgccttt tgttttgtga tattttatgt 540
atatataact attgactttt gtcactctat gatagtgtct cgggctcttg gcaaaccat 600
5 ttcaaagaaa agttaatgac tgtaattaat taatctgaag ctagaaacag aaccccgagg 660
taaaagaaaa agacagagca catgaagttt agtactttta tatatttaat atatcattct 720
ttctttattgc ttatctctaa agcaaaaact tccctaaacc ctaagccaaa ggactcagat 780
cgatgcagaa ccaagaaggc ttgttttgga ttgagagcc aaatgcaaag aaaaaaactc 840
tt 842

10